


































































































































































the 1990s, the number of people with a diagnosis of schizophrenia has increased in the United Kingdom (Meltzer 1998). The prevalence of schizophrenia in the United Kingdom is estimated to be 1.2% (Meltzer 1998).

There is a growing awareness of the need to improve the lives of people with a diagnosis of schizophrenia. The United Kingdom has a number of national strategies for mental health care, including the 1998 *Mental Health Act* (MHA) and the 1999 *Mental Health Strategy* (MHS). The MHA and MHS are both aimed at improving the lives of people with a diagnosis of schizophrenia. The MHA is aimed at improving the lives of people with a diagnosis of schizophrenia who are in the community. The MHS is aimed at improving the lives of people with a diagnosis of schizophrenia who are in hospital.

The MHA and MHS are both aimed at improving the lives of people with a diagnosis of schizophrenia. The MHA is aimed at improving the lives of people with a diagnosis of schizophrenia who are in the community. The MHS is aimed at improving the lives of people with a diagnosis of schizophrenia who are in hospital. The MHA and MHS are both aimed at improving the lives of people with a diagnosis of schizophrenia. The MHA is aimed at improving the lives of people with a diagnosis of schizophrenia who are in the community. The MHS is aimed at improving the lives of people with a diagnosis of schizophrenia who are in hospital.

The MHA and MHS are both aimed at improving the lives of people with a diagnosis of schizophrenia. The MHA is aimed at improving the lives of people with a diagnosis of schizophrenia who are in the community. The MHS is aimed at improving the lives of people with a diagnosis of schizophrenia who are in hospital. The MHA and MHS are both aimed at improving the lives of people with a diagnosis of schizophrenia. The MHA is aimed at improving the lives of people with a diagnosis of schizophrenia who are in the community. The MHS is aimed at improving the lives of people with a diagnosis of schizophrenia who are in hospital.

The MHA and MHS are both aimed at improving the lives of people with a diagnosis of schizophrenia. The MHA is aimed at improving the lives of people with a diagnosis of schizophrenia who are in the community. The MHS is aimed at improving the lives of people with a diagnosis of schizophrenia who are in hospital. The MHA and MHS are both aimed at improving the lives of people with a diagnosis of schizophrenia. The MHA is aimed at improving the lives of people with a diagnosis of schizophrenia who are in the community. The MHS is aimed at improving the lives of people with a diagnosis of schizophrenia who are in hospital.

The MHA and MHS are both aimed at improving the lives of people with a diagnosis of schizophrenia. The MHA is aimed at improving the lives of people with a diagnosis of schizophrenia who are in the community. The MHS is aimed at improving the lives of people with a diagnosis of schizophrenia who are in hospital. The MHA and MHS are both aimed at improving the lives of people with a diagnosis of schizophrenia. The MHA is aimed at improving the lives of people with a diagnosis of schizophrenia who are in the community. The MHS is aimed at improving the lives of people with a diagnosis of schizophrenia who are in hospital.

The MHA and MHS are both aimed at improving the lives of people with a diagnosis of schizophrenia. The MHA is aimed at improving the lives of people with a diagnosis of schizophrenia who are in the community. The MHS is aimed at improving the lives of people with a diagnosis of schizophrenia who are in hospital. The MHA and MHS are both aimed at improving the lives of people with a diagnosis of schizophrenia. The MHA is aimed at improving the lives of people with a diagnosis of schizophrenia who are in the community. The MHS is aimed at improving the lives of people with a diagnosis of schizophrenia who are in hospital.



