



UNIVERSITY OF PISA

DEPARTMENT OF COMPUTER SCIENCE

COMPUTATIONAL MATHEMATICS
WILDCARD PROJECT NR. 5 WITH MACHINE LEARNING
GROUP 35

Support Vector Machines

Author:

Donato Meoli
d.meoli@studenti.unipi.it

June, 2021

Contents

List of Figures	3
List of Tables	4
List of Algorithms	5
List of Theorems	6
1 Track	7
2 Abstract	7
3 Linear Support Vector Classifier	8
3.1 Hinge loss	9
3.1.1 Primal formulation	9
3.1.2 Wolfe Dual formulation	11
3.1.3 Lagrangian Dual formulation	12
3.2 Squared Hinge loss	14
3.2.1 Primal formulation	14
3.2.2 Wolfe Dual formulation	15
3.2.3 Lagrangian Dual formulation	15
4 Linear Support Vector Regression	17
4.1 Epsilon-insensitive loss	18
4.1.1 Primal formulation	18
4.1.2 Wolfe Dual formulation	18
4.1.3 Lagrangian Dual formulation	20
4.2 Squared Epsilon-insensitive loss	22
4.2.1 Primal formulation	22
4.2.2 Wolfe Dual formulation	23
4.2.3 Lagrangian Dual formulation	23
5 Nonlinear Support Vector Machines	25
5.1 Polynomial kernel	25
5.2 Gaussian kernel	26
5.3 Laplacian kernel	26
6 Optimization Methods	27
6.1 Gradient Descent for Primal formulations	30
6.1.1 Nonsmooth	31
6.1.2 Smooth	32
6.1.3 Momentum	33
6.2 Sequential Minimal Optimization for Wolfe Dual formulations	36
6.2.1 Classification	36
6.2.2 Regression	41
6.3 AdaGrad for Lagrangian Dual formulations	45
6.4 Losses properties	46

7	Experiments	48
7.1	Support Vector Classifier	48
7.1.1	Hinge loss	48
7.1.2	Squared Hinge loss	53
7.2	Support Vector Regression	56
7.2.1	Epsilon-insensitive loss	56
7.2.2	Squared Epsilon-insensitive loss	67
8	Conclusions	74
	References	75

List of Figures

1	Linear SVC hyperplane	9
2	Hinge loss with different optimization steps	10
3	Squared Hinge loss with different optimization steps	14
4	Linear SVR hyperplane	17
5	Epsilon-insensitive loss with different optimization steps	19
6	Squared Epsilon-insensitive loss with different optimization steps	22
7	Polynomial SVM hyperplanes	25
8	Gaussian SVM hyperplanes	26
9	Laplacian SVM hyperplanes	27
10	Polyak's and Nesterov's Momentum	33
11	SMO for two Lagrange multipliers	36
12	SGD Convergence for the Primal formulation of the \mathcal{L}_1 -SVC	49
13	AdaGrad convergence for the Lagrangian Dual formulation of the nonlinear \mathcal{L}_1 -SVC	52
14	SGD convergence for the Primal formulation of the \mathcal{L}_2 -SVC	53
15	AdaGrad convergence for the Lagrangian Dual formulation of the nonlinear \mathcal{L}_2 -SVC	55
16	SGD convergence for the Primal formulation of the \mathcal{L}_1 -SVR	58
17	AdaGrad convergence for the Lagrangian Dual formulation of the linear \mathcal{L}_1 -SVR	61
18	AdaGrad convergence for the Lagrangian Dual formulation of the Polynomial \mathcal{L}_1 -SVR	65
19	AdaGrad convergence for the Lagrangian Dual formulation of the Gaussian \mathcal{L}_1 -SVR	66
20	SGD convergence for the Primal formulation of the \mathcal{L}_2 -SVR	68
21	AdaGrad convergence for the Lagrangian Dual formulation of the linear \mathcal{L}_2 -SVR	70
22	AdaGrad convergence for the Lagrangian Dual formulation of the Polynomial \mathcal{L}_2 -SVR	72
23	AdaGrad convergence for the Lagrangian Dual formulation of the Gaussian \mathcal{L}_2 -SVR	73

List of Tables

1	Subgradient Descent convergence rates and iterations complexities	32
2	Gradient Descent convergence rates and iterations complexities	32
3	Nesterov's Accelerated Gradient Descent convergence rates and iterations complexities	34
4	SVM's objectives properties for primal formulations	46
5	SVM's objectives convergence rates for primal formulations	47
6	SVM's objectives convergence rate for Lagrangian dual formulations	47
7	Primal \mathcal{L}_1 -SVC results	48
8	Wolfe Dual linear \mathcal{L}_1 -SVC results	49
9	Lagrangian Dual linear \mathcal{L}_1 -SVC results	50
10	Wolfe Dual nonlinear \mathcal{L}_1 -SVC results	50
11	Lagrangian Dual nonlinear \mathcal{L}_1 -SVC results	51
12	Primal \mathcal{L}_2 -SVC results	53
13	Lagrangian Dual linear \mathcal{L}_2 -SVC results	54
14	Lagrangian Dual nonlinear \mathcal{L}_2 -SVC results	54
15	Primal \mathcal{L}_1 -SVR results	57
16	Wolfe Dual linear \mathcal{L}_1 -SVR results	59
17	Lagrangian Dual linear \mathcal{L}_1 -SVR results	60
18	Wolfe Dual nonlinear \mathcal{L}_1 -SVR formulation results	63
19	Lagrangian Dual nonlinear \mathcal{L}_1 -SVR results	64
20	Primal \mathcal{L}_2 -SVR results	67
21	Lagrangian Dual linear \mathcal{L}_2 -SVR results	69
22	Lagrangian Dual nonlinear \mathcal{L}_2 -SVR results	71

List of Algorithms

1	Gradient Descent	30
2	Stochastic Gradient Descent	30
3	Polyak's Accelerated Gradient Descent or Polyak Heavy-Ball method	33
4	Nesterov's Accelerated Gradient Descent or Nesterov Heavy-Ball method	34
5	Sequential Minimal Optimization for Classification	38
6	Sequential Minimal Optimization for Regression	43
7	AdaGrad	45

List of Theorems

1	Definition (Convexity)	27
2	Definition (Strict Convexity)	27
3	Definition (Strong Convexity)	28
4	Definition (L_f -Lipschitz continuity)	28
5	Definition (L-Lipschitz continuity)	28
6	Definition (Subgradient)	28
7	Theorem (L_f -Lipschitz continuity for convex functions)	29
8	Theorem (μ -strong convexity and L-Lipschitz continuity for convex functions)	29
9	Theorem (Stochastic Gradient Descent convergence for convex functions)	30
10	Theorem (Stochastic Gradient Descent convergence for strongly convex functions)	31
11	Theorem (Subgradient Descent convergence for convex functions with Polyak's stepsize)	31
12	Theorem (Subgradient Descent convergence for convex functions)	31
13	Theorem (Subgradient Descent convergence for strongly convex functions)	31
14	Theorem (Gradient Descent convergence for convex functions)	32
15	Theorem (Gradient Descent convergence for strongly convex functions)	32
16	Theorem (Gradient Descent convergence for convex quadratic functions)	32
17	Theorem (Polyak's Accelerated Gradient Descent convergence for convex quadratic functions)	33
18	Theorem (Nesterov's Accelerated Gradient Descent convergence for convex functions)	34
19	Theorem (Nesterov's Accelerated Gradient Descent convergence for strongly convex functions)	34

1 Track

(M1.1) is a *Support Vector Classifier (SVC)* with the *hinge* loss.

(A1.1.1) is a *momentum descent* approach [1, 2, 3], an *accelerated gradient* method for solving the SVC in its *primal* formulation.

(A1.1.2) is the *Sequential Minimal Optimization (SMO)* algorithm [4, 5], an ad hoc *active set* method for training a SVC in its *Wolfe dual* formulation with *linear*, *polynomial* and *gaussian* kernels.

(A1.1.3) is the *AdaGrad* algorithm [6], a *deflected subgradient* method for solving the SVC in its *Lagrangian dual* formulation with *linear*, *polynomial* and *gaussian* kernels.

(M1.2) is a *Support Vector Classifier (SVC)* with the *squared hinge* loss.

(A1.2.1) is a *momentum descent* approach [1, 2, 3], an *accelerated gradient* method for solving the SVC in its *primal* formulation.

(A1.2.2) is the *AdaGrad* algorithm [6], a *deflected subgradient* method for solving the SVC in its *Lagrangian dual* formulation with *linear*, *polynomial* and *gaussian* kernels.

(M2.1) is a *Support Vector Regression (SVR)* with the *epsilon-insensitive* loss.

(A2.1.1) is a *momentum descent* approach [1, 2, 3], an *accelerated gradient* method for solving the SVR in its *primal* formulation.

(A2.1.2) is the *Sequential Minimal Optimization (SMO)* algorithm [7, 8], an ad hoc *active set* method for training a SVR in its *Wolfe dual* formulation with *linear*, *gaussian* and *laplacian* kernels.

(A2.1.3) is the *AdaGrad* algorithm [6], a *deflected subgradient* method for solving the SVR in its *Lagrangian dual* formulation with *linear*, *gaussian* and *laplacian* kernels.

(M2.2) is a *Support Vector Regression (SVR)* with the *squared epsilon-insensitive* loss.

(A2.2.1) is a *momentum descent* approach [1, 2, 3], an *accelerated gradient* method for solving the SVR in its *primal* formulation.

(A2.2.2) is the *AdaGrad* algorithm [6], a *deflected subgradient* method for solving the SVR in its *Lagrangian dual* formulation with *linear*, *gaussian* and *laplacian* kernels.

2 Abstract

A *Support Vector Machine* is a learning model used both for *classification* and *regression* tasks whose goal is to construct a *maximum margin separator*, i.e., a decision boundary with the largest distance from the nearest training data points.

The aim of this report is to compare the *primal*, the *Wolfe dual* [9] and the *Lagrangian dual* formulations of this model in terms of *numerical precision*, *accuracy* and *complexity*.

Firstly, I will provide a detailed mathematical derivation of the model for all these formulations, then I will propose two algorithms to solve the optimization problem in case of *constrained* or *unconstrained* formulation of the problem, explaining their theoretical properties, i.e., *convergence* and *complexity*.

Finally, I will show some experiments for *linearly* and *nonlinearly* separable generated datasets to compare the performance of different *kernels*, also by comparing the *custom* results with *sklearn* SVM implementations, i.e., *liblinear* [10] and *libsvm* [11] implementations, and *cvxopt* [12] QP solver.

3 Linear Support Vector Classifier

Given n training points, where each input x_i has m attributes, i.e., is of dimensionality m , and is in one of two classes $y_i = \pm 1$, i.e., our training data is of the form:

$$\{(x_i, y_i), x_i \in \mathbb{R}^m, y_i = \pm 1, i = 1, \dots, n\} \quad (1)$$

For simplicity we first assume that data are (not fully) linearly separable in the input space x , meaning that we can draw a line separating the two classes when $m = 2$, a plane for $m = 3$ and, more in general, a hyperplane for an arbitrary m .

Support vectors are the examples closest to the separating hyperplane and the aim of support vector machines is to orientate this hyperplane in such a way as to be as far as possible from the closest members of both classes, i.e., we need to maximize this margin.

This hyperplane is represented by the equation $w^T x + b = 0$. So, we need to find w and b so that our training data can be described by:

$$\begin{aligned} w^T x_i + b &\geq +1 - \xi_i, \forall y_i = +1 \\ w^T x_i + b &\leq -1 + \xi_i, \forall y_i = -1 \\ \xi_i &\geq 0 \quad \forall_i \end{aligned} \quad (2)$$

where the positive slack variables ξ_i are introduced to allow misclassified points. In this way data points on the incorrect side of the margin boundary will have a penalty that increases with the distance from it.

These two equations can be combined into:

$$\begin{aligned} y_i(w^T x_i + b) &\geq 1 - \xi_i \quad \forall_i \\ \xi_i &\geq 0 \quad \forall_i \end{aligned} \quad (3)$$

The margin is equal to $\frac{1}{\|w\|}$ and maximizing it subject to the constraint in (3) while as we are trying to reduce the number of misclassifications is equivalent to finding:

$$\begin{aligned} \min_{w, b, \xi} \quad & \|w\| + C \sum_{i=1}^n \xi_i \\ \text{subject to} \quad & y_i(w^T x_i + b) \geq 1 - \xi_i \quad \forall_i \\ & \xi_i \geq 0 \quad \forall_i \end{aligned} \quad (4)$$

Minimizing $\|w\|$ is equivalent to minimizing $\frac{1}{2}\|w\|^2$, but in this form we will deal with a 1-strongly convex regularization term that has more desirable convergence properties. So we need to find:

$$\begin{aligned} \min_{w, b, \xi} \quad & \frac{1}{2}\|w\|^2 + C \sum_{i=1}^n \xi_i \\ \text{subject to} \quad & y_i(w^T x_i + b) \geq 1 - \xi_i \quad \forall_i \\ & \xi_i \geq 0 \quad \forall_i \end{aligned} \quad (5)$$

where the parameter C controls the trade-off between the slack variable penalty and the size of the margin.




Figure 1: Linear SVC hyperplane

3.1 Hinge loss

The *hinge* loss is defined as:

$$\mathcal{L}_1 = \max(0, 1 - y(w^T x + b)) \quad (6)$$

or, equivalently:

$$\mathcal{L}_1 = \begin{cases} 0 & \text{if } y(w^T x + b) \geq 1 \\ 1 - y(w^T x + b) & \text{otherwise} \end{cases} \quad (7)$$

and it is a nondifferentiable convex function due to its nonsmoothness in 1, but has a subgradient that is given by:

$$\partial_w \mathcal{L}_1 = \begin{cases} -yx & \text{if } y(w^T x + b) < 1 \\ 0 & \text{otherwise} \end{cases} \quad (8)$$

3.1.1 Primal formulation

The general primal unconstrained formulation takes the form:

$$\min_{w,b} \frac{1}{2} \|w\|^2 + C \sum_{i=1}^n \mathcal{L}(w, b; x_i, y_i) \quad (9)$$

where $\frac{1}{2} \|w\|^2$ is the *regularization term* and $\mathcal{L}(w, b; x_i, y_i)$ is the *loss function* associated with the observation (x_i, y_i) [13].

The quadratic optimization problem (5) can be equivalently formulated as:

$$\min_{w,b} \frac{1}{2} \|w\|^2 + C \sum_{i=1}^n \max(0, 1 - y_i(w^T x_i + b)) \quad (10)$$

where we make use of the *hinge* loss (6) or (7).

The above formulation penalizes slacks ξ linearly and is called \mathcal{L}_1 -SVC.




Figure 2: Hinge loss with different optimization steps

To simplify the notation and so also the design of the algorithms, the simplest approach to learn the bias term b is that of including that into the *regularization term*; so we can rewrite (9) as follows:

$$\min_{w,b} \frac{1}{2}(\|w\|^2 + b^2) + C \sum_{i=1}^n \mathcal{L}(w, b; x_i, y_i) \quad (11)$$

or, equivalently, by augmenting the weight vector w with the bias term b and each instance x_i with an additional dimension, i.e., with constant value equal to 1:

$$\begin{aligned} \min_w \quad & \frac{1}{2}\|\hat{w}\|^2 + C \sum_{i=1}^n \mathcal{L}(\hat{w}; \hat{x}_i, y_i) \\ \text{where} \quad & \hat{w}^T = [w^T, b] \\ & \hat{x}_i^T = [x_i^T, 1] \end{aligned} \quad (12)$$

with the advantages of having convex properties of the objective function useful for convergence analysis and the possibility to directly apply algorithms designed for models without the bias term.

In the specific case of the \mathcal{L}_1 -SVC the objective (10) become:

$$\min_{w,b} \frac{1}{2}(\|w\|^2 + b^2) + C \sum_{i=1}^n \max(0, 1 - y_i(w^T x_i + b)) \quad (13)$$

Note that in terms of numerical optimization the formulation (10) is not equivalent to (13) since in the first one the bias term b does not contribute to the *regularization term*, so the SVM formulation is based on an unregularized bias term b , as highlighted by the *statistical learning theory*. But, in machine learning sense, numerical experiments in [14] show that the accuracy does not vary much when the bias term b is embedded into the weight vector w .

3.1.2 Wolfe Dual formulation

To reformulate the (5) as a *Wolfe dual*, we need to allocate the Lagrange multipliers $\alpha_i, \mu_i \geq 0 \forall_i$:

$$\max_{\alpha, \mu} \min_{w, b, \xi} \mathcal{W}(w, b, \xi, \alpha, \mu) = \frac{1}{2} \|w\|^2 + C \sum_{i=1}^n \xi_i - \sum_{i=1}^n \alpha_i (y_i (w^T x_i + b) - 1 + \xi_i) - \sum_{i=1}^n \mu_i \xi_i \quad (14)$$

We wish to find the w , b and ξ_i which minimizes, and the α and μ which maximizes \mathcal{W} , provided $\alpha_i \geq 0, \mu_i \geq 0 \forall_i$. We can do this by differentiating \mathcal{W} wrt w and b and setting the derivatives to 0:

$$\frac{\partial \mathcal{W}}{\partial w} = w - \sum_{i=1}^n \alpha_i y_i x_i \Rightarrow w = \sum_{i=1}^n \alpha_i y_i x_i \quad (15)$$

$$\frac{\partial \mathcal{W}}{\partial b} = - \sum_{i=1}^n \alpha_i y_i \Rightarrow \sum_{i=1}^n \alpha_i y_i = 0 \quad (16)$$

$$\frac{\partial \mathcal{W}}{\partial \xi_i} = 0 \Rightarrow C = \alpha_i + \mu_i \quad (17)$$

Substituting (15) and (16) into (14) together with $\mu_i \geq 0 \forall_i$, which implies that $\alpha \leq C$, gives a new formulation being dependent on α . We therefore need to find:

$$\begin{aligned} \max_{\alpha} \mathcal{W}(\alpha) &= \sum_{i=1}^n \alpha_i - \frac{1}{2} \sum_{i,j} \alpha_i \alpha_j y_i y_j \langle x_i, x_j \rangle \\ &= \sum_{i=1}^n \alpha_i - \frac{1}{2} \sum_{i,j} \alpha_i Q_{ij} \alpha_j \text{ where } Q_{ij} = y_i y_j \langle x_i, x_j \rangle \\ &= \sum_{i=1}^n \alpha_i - \frac{1}{2} \alpha^T Q \alpha \text{ subject to } 0 \leq \alpha_i \leq C \forall_i, \sum_{i=1}^n \alpha_i y_i = 0 \end{aligned} \quad (18)$$

or, equivalently:

$$\begin{aligned} \min_{\alpha} \quad & \frac{1}{2} \alpha^T Q \alpha + q^T \alpha \\ \text{subject to} \quad & 0 \leq \alpha_i \leq C \forall_i \\ & y^T \alpha = 0 \end{aligned} \quad (19)$$

where $q^T = [1, \dots, 1]$.

By solving (19) we will know α and, from (15), we will get w , so we need to calculate b .

We know that any data point satisfying (16) which is a support vector x_s will have the form:

$$y_s (w^T x_s + b) = 1 \quad (20)$$

and, by substituting in (15), we get:

$$y_s \left(\sum_{m \in S} \alpha_m y_m \langle x_m, x_s \rangle + b \right) = 1 \quad (21)$$

where s denotes the set of indices of the support vectors and is determined by finding the indices i where $\alpha_i > 0$, i.e., nonzero Lagrange multipliers.

Multiplying through by y_s and then using $y_s^2 = 1$ from (2):

$$y_s^2 \left(\sum_{m \in S} \alpha_m y_m \langle x_m, x_s \rangle + b \right) = y_s \quad (22)$$

$$b = y_s - \sum_{m \in S} \alpha_m y_m \langle x_m, x_s \rangle \quad (23)$$

Instead of using an arbitrary support vector x_s , it is better to take an average over all of the support vectors in S :

$$b = \frac{1}{N_s} \sum_{s \in S} y_s - \sum_{m \in S} \alpha_m y_m \langle x_m, x_s \rangle \quad (24)$$

We now have the variables w and b that define our separating hyperplane's optimal orientation and hence our support vector machine. Each new point x' is classified by evaluating:

$$y' = \text{sign} \left(\sum_{i=1}^n \alpha_i y_i \langle x_i, x' \rangle + b \right) \quad (25)$$

From (19) we can notice that the equality constraint $y^T \alpha = 0$ arises from the stationarity condition $\partial_b \mathcal{W} = 0$. So, again, for simplicity, we can again consider the bias term b embedded into the weight vector. We report below the box-constrained dual formulation [14] that arises from the primal (11) or (12) where the bias term b is embedded into the weight vector w :

$$\begin{aligned} \min_{\alpha} \quad & \frac{1}{2} \alpha^T (Q + yy^T) \alpha + q^T \alpha \\ \text{subject to} \quad & 0 \leq \alpha_i \leq C \quad \forall_i \end{aligned} \quad (26)$$

3.1.3 Lagrangian Dual formulation

In order to relax the constraints in the *Wolfe dual* formulation (19) we define the problem as a *Lagrangian dual* relaxation by embedding them into objective function, so we need to allocate the Lagrange multipliers μ and $\lambda_+, \lambda_- \geq 0$:

$$\begin{aligned} \max_{\mu, \lambda_+, \lambda_-} \min_{\alpha} \mathcal{L}(\alpha, \mu, \lambda_+, \lambda_-) &= \frac{1}{2} \alpha^T Q \alpha + q^T \alpha - \mu^T (y^T \alpha) - \lambda_+^T (u - \alpha) - \lambda_-^T \alpha \\ &= \frac{1}{2} \alpha^T Q \alpha + (q - \mu y + \lambda_+ - \lambda_-)^T \alpha - \lambda_+^T u \\ \text{subject to} \quad & \lambda_+, \lambda_- \geq 0 \end{aligned} \quad (27)$$

where the upper bound $u^T = [C, \dots, C]$.

Taking the derivative of the Lagrangian \mathcal{L} wrt α and settings it to 0 gives:

$$\frac{\partial \mathcal{L}}{\partial \alpha} = 0 \Rightarrow Q\alpha + (q - \mu y + \lambda_+ - \lambda_-) = 0 \quad (28)$$

With α optimal solution of the linear system:

$$Q\alpha = -(q - \mu y + \lambda_+ - \lambda_-) \quad (29)$$

the gradient wrt μ , λ_+ and λ_- are:

$$\frac{\partial \mathcal{L}}{\partial \mu} = -y\alpha \quad (30)$$

$$\frac{\partial \mathcal{L}}{\partial \lambda_+} = \alpha - u \quad (31)$$

$$\frac{\partial \mathcal{L}}{\partial \lambda_-} = -\alpha \quad (32)$$

From (19) we can notice that the equality constraint $y^T \alpha = 0$ arises from the stationarity condition $\partial_b \mathcal{W} = 0$. So, again, for simplicity, we can again consider the bias term b embedded into the weight vector. In this way the dimensionality of (27) is reduced of 1/3 by removing the multipliers μ which was allocated to control the equality constraint $y^T \alpha = 0$, so we will end up solving exactly the problem (26).

$$\begin{aligned} \max_{\lambda_+, \lambda_-} \min_{\alpha} \mathcal{L}(\alpha, \lambda_+, \lambda_-) &= \frac{1}{2} \alpha^T (Q + yy^T) \alpha + q^T \alpha - \lambda_+^T (u - \alpha) - \lambda_-^T \alpha \\ &= \frac{1}{2} \alpha^T (Q + yy^T) \alpha + (q + \lambda_+ - \lambda_-)^T \alpha - \lambda_+^T u \end{aligned} \quad (33)$$

subject to $\lambda_+, \lambda_- \geq 0$

where, again, the upper bound $u^T = [C, \dots, C]$.

Now, taking the derivative of the Lagrangian \mathcal{L} wrt α and settings it to 0 gives:

$$\frac{\partial \mathcal{L}}{\partial \alpha} = 0 \Rightarrow (Q + yy^T) \alpha + (q + \lambda_+ - \lambda_-) = 0 \quad (34)$$

With α optimal solution of the linear system:

$$(Q + yy^T) \alpha = -(q + \lambda_+ - \lambda_-) \quad (35)$$

the gradient wrt λ_+ and λ_- are:

$$\frac{\partial \mathcal{L}}{\partial \lambda_+} = \alpha - u \quad (36)$$

$$\frac{\partial \mathcal{L}}{\partial \lambda_-} = -\alpha \quad (37)$$

3.2 Squared Hinge loss

The *squared hinge* loss is defined as:

$$\mathcal{L}_2 = \max(0, 1 - y(w^T x + b))^2 \quad (38)$$

or, equivalently:

$$\mathcal{L}_2 = \begin{cases} 0 & \text{if } y(w^T x + b) \geq 1 \\ (1 - y(w^T x + b))^2 & \text{otherwise} \end{cases} \quad (39)$$

It is a strictly convex function and its gradient is given by:

$$\nabla_w \mathcal{L}_2 = \begin{cases} -2 \max(0, 1 - y(w^T x + b))yx & \text{if } y(w^T x + b) < 1 \\ 0 & \text{otherwise} \end{cases} \quad (40)$$

3.2.1 Primal formulation

Since smoothed versions of objective functions may be preferred for optimization, we can reformulate (10) as:

$$\min_{w,b} \frac{1}{2} \|w\|^2 + C \sum_{i=1}^n \max(0, 1 - y_i(w^T x_i + b))^2 \quad (41)$$

where we make use of the *squared hinge* loss that quadratically penalized slacks ξ and is called \mathcal{L}_2 -SVC.

The \mathcal{L}_2 -SVC objective (41) can be rewritten in form (11) or (12) as:

$$\min_{w,b} \frac{1}{2} (\|w\|^2 + b^2) + C \sum_{i=1}^n \max(0, 1 - y_i(w^T x_i + b))^2 \quad (42)$$




Figure 3: Squared Hinge loss with different optimization steps

3.2.2 Wolfe Dual formulation

As done for the \mathcal{L}_1 -SVC we can derive the *Wolfe dual* formulation of the \mathcal{L}_2 -SVC by obtaining:

$$\begin{aligned} \min_{\alpha} \quad & \frac{1}{2} \alpha^T (Q + D) \alpha + q^T \alpha \\ \text{subject to} \quad & \alpha_i \geq 0 \quad \forall_i \\ & y^T \alpha = 0 \end{aligned} \quad (43)$$

or, alternatively, with the regularized bias term by obtaining:

$$\begin{aligned} \min_{\alpha} \quad & \frac{1}{2} \alpha^T (Q + yy^T + D) \alpha + q^T \alpha \\ \text{subject to} \quad & \alpha_i \geq 0 \quad \forall_i \end{aligned} \quad (44)$$

where the diagonal matrix $D_{ii} = \frac{1}{2C} \quad \forall_i$.

3.2.3 Lagrangian Dual formulation

In order to relax the constraints in the \mathcal{L}_2 -SVC *Wolfe dual* formulation (43) we define the problem as a *Lagrangian dual* relaxation by embedding them into objective function, so we need to allocate the Lagrange multipliers μ and $\lambda \geq 0$:

$$\begin{aligned} \max_{\mu, \lambda} \min_{\alpha} \mathcal{L}(\alpha, \mu, \lambda) &= \frac{1}{2} \alpha^T (Q + D) \alpha + q^T \alpha - \mu^T (y^T \alpha) - \lambda^T \alpha \\ &= \frac{1}{2} \alpha^T (Q + D) \alpha + (q - \mu y - \lambda)^T \alpha \\ \text{subject to} \quad & \lambda \geq 0 \end{aligned} \quad (45)$$

Taking the derivative of the Lagrangian \mathcal{L} wrt α and settings it to 0 gives:

$$\frac{\partial \mathcal{L}}{\partial \alpha} = 0 \Rightarrow (Q + D) \alpha + (q - \mu y - \lambda) = 0 \quad (46)$$

With α optimal solution of the linear system:

$$(Q + D) \alpha = -(q - \mu y - \lambda) \quad (47)$$

the gradient wrt μ and λ are:

$$\frac{\partial \mathcal{L}}{\partial \mu} = -y \alpha \quad (48)$$

$$\frac{\partial \mathcal{L}}{\partial \lambda} = -\alpha \quad (49)$$

From (19) we can notice that the equality constraint $y^T \alpha = 0$ arises from the stationarity condition $\partial_b \mathcal{W} = 0$. So, again, for simplicity, we can again consider the bias term b embedded into the weight vector. In this way the dimensionality of (45) is reduced of 1/3 by removing the multipliers μ which was allocated to control the equality constraint $y^T \alpha = 0$, so we will end up solving exactly the problem (44).

$$\begin{aligned} \max_{\lambda} \min_{\alpha} \mathcal{L}(\alpha, \lambda) &= \frac{1}{2} \alpha^T (Q + yy^T + D) \alpha + q^T \alpha - \lambda^T \alpha \\ &= \frac{1}{2} \alpha^T (Q + yy^T + D) \alpha + (q - \lambda)^T \alpha \\ \text{subject to} \quad & \lambda \geq 0 \end{aligned} \quad (50)$$

where, again, the upper bound $u^T = [C, \dots, C]$.

Now, taking the derivative of the Lagrangian \mathcal{L} wrt α and settings it to 0 gives:

$$\frac{\partial \mathcal{L}}{\partial \alpha} = 0 \Rightarrow (Q + yy^T + D)\alpha + (q - \lambda) = 0 \quad (51)$$

With α optimal solution of the linear system:

$$(Q + yy^T + D)\alpha = - (q - \lambda) \quad (52)$$

the gradient wrt λ is:

$$\frac{\partial \mathcal{L}}{\partial \lambda} = -\alpha \quad (53)$$

4 Linear Support Vector Regression

In the case of regression the goal is to predict a real-valued output for y' so that our training data is of the form:

$$\{(x_i, y_i), x \in \mathbb{R}^m, y_i \in \mathbb{R}, i = 1, \dots, n\} \quad (54)$$

The regression SVM use a loss function that not allocating a penalty if the predicted value y'_i is less than a distance ϵ away from the actual value y_i , i.e., if $|y_i - y'_i| \leq \epsilon$, where $y'_i = w^T x_i + b$. The region bound by $y'_i \pm \epsilon \forall_i$ is called an ϵ -insensitive tube. The output variables which are outside the tube are given one of two slack variable penalties depending on whether they lie above, ξ^+ , or below, ξ^- , the tube, provided $\xi^+ \geq 0$ and $\xi^- \geq 0 \forall_i$:

$$\begin{aligned} y_i &\leq y'_i + \epsilon + \xi^+ \forall_i \\ y_i &\geq y'_i - \epsilon - \xi^- \forall_i \\ \xi_i^+, \xi_i^- &\geq 0 \forall_i \end{aligned} \quad (55)$$

The objective function for SVR can then be written as:

$$\begin{aligned} \min_{w, b, \xi^+, \xi^-} \quad & \frac{1}{2} \|w\|^2 + C \sum_{i=1}^n (\xi_i^+ + \xi_i^-) \\ \text{subject to} \quad & y_i - w^T x_i - b \leq \epsilon + \xi_i^+ \forall_i \\ & w^T x_i + b - y_i \leq \epsilon + \xi_i^- \forall_i \\ & \xi_i^+, \xi_i^- \geq 0 \forall_i \end{aligned} \quad (56)$$



Figure 4: Linear SVR hyperplane

4.1 Epsilon-insensitive loss

The *epsilon-insensitive* loss is defined as:

$$\mathcal{L}_\epsilon^1 = \max(0, |y - (w^T x + b)| - \epsilon) \quad (57)$$

or, equivalently:

$$\mathcal{L}_\epsilon^1 = \begin{cases} 0 & \text{if } |y - (w^T x + b)| \leq \epsilon \\ |y - (w^T x + b)| - \epsilon & \text{otherwise} \end{cases} \quad (58)$$

As the *hinge* loss, also the *epsilon-insensitive* loss is a nondifferentiable convex function due to its nonsmoothness in $\pm\epsilon$, but has a subgradient that is given by:

$$\partial_w \mathcal{L}_\epsilon^1 = \begin{cases} \frac{y - (w^T x + b)}{|y - (w^T x + b)|} x & \text{if } |y - (w^T x + b)| \geq \epsilon \\ 0 & \text{otherwise} \end{cases} \quad (59)$$

4.1.1 Primal formulation

The general primal unconstrained formulation takes the same form of (9).

The quadratic optimization problem (56) can be equivalently formulated as:

$$\min_{w,b} \frac{1}{2} \|w\|^2 + C \sum_{i=1}^n \max(0, |y_i - (w^T x_i + b)| - \epsilon) \quad (60)$$

where we make use of the *epsilon-insensitive* loss (57) or (58).

The above formulation penalizes slacks ξ linearly and is called \mathcal{L}_1 -SVR.

The \mathcal{L}_1 -SVR objective (60) can be rewritten in form (11) or (12) as:

$$\min_{w,b} \frac{1}{2} (\|w\|^2 + b^2) + C \sum_{i=1}^n \max(0, |y_i - (w^T x_i + b)| - \epsilon) \quad (61)$$

4.1.2 Wolfe Dual formulation

To reformulate the (56) as a *Wolfe dual*, we introduce the Lagrange multipliers $\alpha_i^+, \alpha_i^-, \mu_i^+, \mu_i^- \geq 0 \forall i$:

$$\begin{aligned} \max_{\alpha^+, \alpha^-, \mu^+, \mu^-} \min_{w, b, \xi^+, \xi^-} \mathcal{W}(w, b, \xi^+, \xi^-, \alpha^+, \alpha^-, \mu^+, \mu^-) = & \frac{1}{2} \|w\|^2 + C \sum_{i=1}^n (\xi_i^+ + \xi_i^-) - \sum_{i=1}^n (\mu_i^+ \xi_i^+ + \mu_i^- \xi_i^-) \\ & - \sum_{i=1}^n \alpha_i^+ (\epsilon + \xi_i^+ + y'_i - y_i) - \sum_{i=1}^n \alpha_i^- (\epsilon + \xi_i^- - y'_i + y_i) \end{aligned} \quad (62)$$

Substituting for y_i , differentiating wrt w, b, ξ^+, ξ^- and setting the derivatives to 0 gives:

$$\frac{\partial \mathcal{W}}{\partial w} = w - \sum_{i=1}^n (\alpha_i^+ - \alpha_i^-) x_i \Rightarrow w = \sum_{i=1}^n (\alpha_i^+ - \alpha_i^-) x_i \quad (63)$$

$$\frac{\partial \mathcal{W}}{\partial b} = - \sum_{i=1}^n (\alpha_i^+ - \alpha_i^-) \Rightarrow \sum_{i=1}^n (\alpha_i^+ - \alpha_i^-) = 0 \quad (64)$$

$$\frac{\partial \mathcal{W}}{\partial \xi_i^+} = 0 \Rightarrow C = \alpha_i^+ + \mu_i^+ \quad (65)$$




Figure 5: Epsilon-insensitive loss with different optimization steps

$$\frac{\partial \mathcal{W}}{\partial \xi_i^-} = 0 \Rightarrow C = \alpha_i^- + \mu_i^- \quad (66)$$

Substituting (63) and (64) in, we now need to maximize \mathcal{W} wrt α_i^+ and α_i^- , where $\alpha_i^+ \geq 0$, $\alpha_i^- \geq 0 \forall i$:

$$\max_{\alpha^+, \alpha^-} \mathcal{W}(\alpha^+, \alpha^-) = \sum_{i=1}^n y_i (\alpha_i^+ - \alpha_i^-) - \epsilon \sum_{i=1}^n (\alpha_i^+ + \alpha_i^-) - \frac{1}{2} \sum_{i,j} (\alpha_i^+ - \alpha_i^-) \langle x_i, x_j \rangle (\alpha_j^+ - \alpha_j^-) \quad (67)$$

Using $\mu_i^+ \geq 0$ and $\mu_i^- \geq 0$ together with (63) and (64) means that $\alpha_i^+ \leq C$ and $\alpha_i^- \leq C$. We therefore need to find:

$$\begin{aligned} \min_{\alpha^+, \alpha^-} & \quad \frac{1}{2} (\alpha^+ - \alpha^-)^T K (\alpha^+ - \alpha^-) + \epsilon e^T (\alpha^+ + \alpha^-) - y^T (\alpha^+ - \alpha^-) \\ \text{subject to} & \quad 0 \leq \alpha_i^+, \alpha_i^- \leq C \forall i \\ & \quad e^T (\alpha^+ - \alpha^-) = 0 \end{aligned} \quad (68)$$

where $e^T = [1, \dots, 1]$.

We can write the (68) in a standard quadratic form as:

$$\begin{aligned} \min_{\alpha} & \quad \frac{1}{2} \alpha^T Q \alpha - q^T \alpha \\ \text{subject to} & \quad 0 \leq \alpha_i \leq C \forall i \\ & \quad e^T \alpha = 0 \end{aligned} \quad (69)$$

where the Hessian matrix $Q = \begin{bmatrix} K & -K \\ -K & K \end{bmatrix}$, $\alpha = \begin{bmatrix} \alpha^+ \\ \alpha^- \end{bmatrix}$, $q = \begin{bmatrix} -y \\ y \end{bmatrix} + \epsilon$, and $e = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$.

Each new predictions y' can be found using:

$$y' = \sum_{i=1}^n (\alpha_i^+ - \alpha_i^-) \langle x_i, x' \rangle + b \quad (70)$$

A set S of support vectors x_s can be created by finding the indices i where $0 \leq \alpha \leq C$ and $\xi_i^+ = 0$ or $\xi_i^- = 0$. This gives us:

$$b = y_s - \epsilon - \sum_{m \in S} (\alpha_m^+ - \alpha_m^-) \langle x_m, x_s \rangle \quad (71)$$

As before it is better to average over all the indices i in S :

$$b = \frac{1}{N_s} \sum_{s \in S} y_s - \epsilon - \sum_{m \in S} (\alpha_m^+ - \alpha_m^-) \langle x_m, x_s \rangle \quad (72)$$

From (69) we can notice that the equality constraint $e^T \alpha = 0$ arises from the stationarity condition $\partial_b \mathcal{W} = 0$. So, again, for simplicity, we can again consider the bias term b embedded into the weight vector. We report below the box-constrained dual formulation [14] that arises from the primal (11) or (12) where the bias term b is embedded into the weight vector w :

$$\begin{aligned} \min_{\alpha} \quad & \frac{1}{2} \alpha^T (Q + ee^T) \alpha + q^T \alpha \\ \text{subject to} \quad & 0 \leq \alpha_i \leq C \quad \forall_i \end{aligned} \quad (73)$$

4.1.3 Lagrangian Dual formulation

In order to relax the constraints in the *Wolfe dual* formulation (68) we define the problem as a *Lagrangian dual* relaxation by embedding them into objective function, so we need to allocate the Lagrange multipliers μ and $\lambda_+, \lambda_- \geq 0$:

$$\begin{aligned} \max_{\mu, \lambda_+, \lambda_-} \min_{\alpha} \mathcal{L}(\alpha, \mu, \lambda_+, \lambda_-) &= \frac{1}{2} \alpha^T Q \alpha + q^T \alpha - \mu^T (e^T \alpha) - \lambda_+^T (u - \alpha) - \lambda_-^T \alpha \\ &= \frac{1}{2} \alpha^T Q \alpha + (q - \mu e + \lambda_+ - \lambda_-)^T \alpha - \lambda_+^T u \\ \text{subject to} \quad & \lambda_+, \lambda_- \geq 0 \end{aligned} \quad (74)$$

where the upper bound $u^T = [C, \dots, C]$.

Taking the derivative of the Lagrangian \mathcal{L} wrt α and settings it to 0 gives:

$$\frac{\partial \mathcal{L}}{\partial \alpha} = 0 \Rightarrow Q\alpha + (q - \mu e + \lambda_+ - \lambda_-) = 0 \quad (75)$$

With α optimal solution of the linear system:

$$Q\alpha = -(q - \mu e + \lambda_+ - \lambda_-) \quad (76)$$

the gradient wrt μ , λ_+ and λ_- are:

$$\frac{\partial \mathcal{L}}{\partial \mu} = -e\alpha \quad (77)$$

$$\frac{\partial \mathcal{L}}{\partial \lambda_+} = \alpha - u \quad (78)$$

$$\frac{\partial \mathcal{L}}{\partial \lambda_-} = -\alpha \quad (79)$$

From (69) we can notice that the equality constraint $e^T \alpha = 0$ arises from the stationarity condition $\partial_b \mathcal{W} = 0$. So, again, for simplicity, we can again consider the bias term b embedded into the weight vector. In this way

the dimensionality of (74) is reduced of 1/3 by removing the multipliers μ which was allocated to control the equality constraint $e^T \alpha = 0$, so we will end up solving exactly the problem (73).

$$\begin{aligned} \max_{\lambda_+, \lambda_-} \min_{\alpha} \mathcal{L}(\alpha, \lambda_+, \lambda_-) &= \frac{1}{2} \alpha^T (Q + ee^T) \alpha + q^T \alpha - \lambda_+^T (u - \alpha) - \lambda_-^T \alpha \\ &= \frac{1}{2} \alpha^T (Q + ee^T) \alpha + (q + \lambda_+ - \lambda_-)^T \alpha - \lambda_+^T u \\ \text{subject to} \quad &\lambda_+, \lambda_- \geq 0 \end{aligned} \tag{80}$$

where, again, the upper bound $u^T = [C, \dots, C]$.

Now, taking the derivative of the Lagrangian \mathcal{L} wrt α and settings it to 0 gives:

$$\frac{\partial \mathcal{L}}{\partial \alpha} = 0 \Rightarrow (Q + ee^T) \alpha + (q + \lambda_+ - \lambda_-) = 0 \tag{81}$$

With α optimal solution of the linear system:

$$(Q + ee^T) \alpha = -(q + \lambda_+ - \lambda_-) \tag{82}$$

the gradient wrt λ_+ and λ_- are:

$$\frac{\partial \mathcal{L}}{\partial \lambda_+} = \alpha - u \tag{83}$$

$$\frac{\partial \mathcal{L}}{\partial \lambda_-} = -\alpha \tag{84}$$

4.2 Squared Epsilon-insensitive loss

The *squared epsilon-insensitive* loss is defined as:

$$\mathcal{L}_\epsilon^2 = \max(0, |y - (w^T x + b)| - \epsilon)^2 \quad (85)$$

or, equivalently:

$$\mathcal{L}_\epsilon^2 = \begin{cases} 0 & \text{if } |y - (w^T x + b)| \leq \epsilon \\ (|y - (w^T x + b)| - \epsilon)^2 & \text{otherwise} \end{cases} \quad (86)$$

As the *squared hinge* loss, also the *squared epsilon-insensitive* loss is a strictly convex function and its gradient is given by:

$$\nabla_w \mathcal{L}_\epsilon^2 = \begin{cases} 2 \operatorname{sign}(y - (w^T x + b))(|y - (w^T x + b)| - \epsilon)x & \text{if } |y - (w^T x + b)| \geq \epsilon \\ 0 & \text{otherwise} \end{cases} \quad (87)$$

4.2.1 Primal formulation

To provide a continuously differentiable function the optimization problem (60) can be formulated as:

$$\min_{w,b} \frac{1}{2} \|w\|^2 + C \sum_{i=1}^n \max(0, |y_i - (w^T x_i + b)| - \epsilon)^2 \quad (88)$$

where we make use of the *squared epsilon-insensitive* loss that quadratically penalized slacks ξ and is called \mathcal{L}_2 -SVR.

The \mathcal{L}_2 -SVR objective (88) can be rewritten in form (11) or (12) as:

$$\min_{w,b} \frac{1}{2} (\|w\|^2 + b^2) + C \sum_{i=1}^n \max(0, |y_i - (w^T x_i + b)| - \epsilon)^2 \quad (89)$$




Figure 6: Squared Epsilon-insensitive loss with different optimization steps

4.2.2 Wolfe Dual formulation

As done for the \mathcal{L}_1 -SVR we can derive the *Wolfe dual* formulation of the \mathcal{L}_2 -SVR by obtaining:

$$\begin{aligned} \min_{\alpha} \quad & \frac{1}{2} \alpha^T (Q + D) \alpha + q^T \alpha \\ \text{subject to} \quad & \alpha_i \geq 0 \quad \forall_i \\ & e^T \alpha = 0 \end{aligned} \quad (90)$$

or, alternatively, with the regularized bias term by obtaining:

$$\begin{aligned} \min_{\alpha} \quad & \frac{1}{2} \alpha^T (Q + ee^T + D) \alpha + q^T \alpha \\ \text{subject to} \quad & \alpha_i \geq 0 \quad \forall_i \end{aligned} \quad (91)$$

where the diagonal matrix $D_{ii} = \frac{1}{2C} \quad \forall_i$.

4.2.3 Lagrangian Dual formulation

In order to relax the constraints in the \mathcal{L}_2 -SVR *Wolfe dual* formulation (90) we define the problem as a *Lagrangian dual* relaxation by embedding them into objective function, so we need to allocate the Lagrange multipliers μ and $\lambda \geq 0$:

$$\begin{aligned} \max_{\mu, \lambda} \min_{\alpha} \mathcal{L}(\alpha, \mu, \lambda) &= \frac{1}{2} \alpha^T (Q + D) \alpha + q^T \alpha - \mu^T (e^T \alpha) - \lambda^T \alpha \\ &= \frac{1}{2} \alpha^T (Q + D) \alpha + (q - \mu e - \lambda)^T \alpha \\ \text{subject to} \quad & \lambda \geq 0 \end{aligned} \quad (92)$$

Taking the derivative of the Lagrangian \mathcal{L} wrt α and settings it to 0 gives:

$$\frac{\partial \mathcal{L}}{\partial \alpha} = 0 \Rightarrow (Q + D) \alpha + (q - \mu e - \lambda) = 0 \quad (93)$$

With α optimal solution of the linear system:

$$(Q + D) \alpha = -(q - \mu e - \lambda) \quad (94)$$

the gradient wrt μ and λ are:

$$\frac{\partial \mathcal{L}}{\partial \mu} = -e \alpha \quad (95)$$

$$\frac{\partial \mathcal{L}}{\partial \lambda} = -\alpha \quad (96)$$

From (69) we can notice that the equality constraint $e^T \alpha = 0$ arises from the stationarity condition $\partial_b \mathcal{W} = 0$. So, again, for simplicity, we can again consider the bias term b embedded into the weight vector. In this way the dimensionality of (92) is reduced of 1/3 by removing the multipliers μ which was allocated to control the equality constraint $e^T \alpha = 0$, so we will end up solving exactly the problem (91).

$$\begin{aligned} \max_{\lambda} \min_{\alpha} \mathcal{L}(\alpha, \lambda) &= \frac{1}{2} \alpha^T (Q + ee^T + D) \alpha + q^T \alpha - \lambda^T \alpha \\ &= \frac{1}{2} \alpha^T (Q + ee^T + D) \alpha + (q - \lambda)^T \alpha \\ \text{subject to} \quad & \lambda \geq 0 \end{aligned} \quad (97)$$

where, again, the upper bound $u^T = [C, \dots, C]$.

Now, taking the derivative of the Lagrangian \mathcal{L} wrt α and settings it to 0 gives:

$$\frac{\partial \mathcal{L}}{\partial \alpha} = 0 \Rightarrow (Q + ee^T + D)\alpha + (q - \lambda) = 0 \quad (98)$$

With α optimal solution of the linear system:

$$(Q + ee^T + D)\alpha = - (q - \lambda) \quad (99)$$

the gradient wrt λ is:

$$\frac{\partial \mathcal{L}}{\partial \lambda} = -\alpha \quad (100)$$

5 Nonlinear Support Vector Machines

When applying our SVC to *linearly separable* data in (18), we have started by creating a matrix Q from the dot product of our input variables:

$$Q_{ij} = y_i y_j k(x_i, x_j) \quad (101)$$

or, a matrix K from the dot product of our input variables in the SVR case (68):

$$K_{ij} = k(x_i, x_j) \quad (102)$$

where $k(x_i, x_j)$ is an example of a family of functions called *kernel functions*.

For any positive definite kernel function k (a so called Mercer kernel), it is guaranteed that there exists a mapping ϕ into a Hilbert space \mathcal{H} , such that:

$$k(x_i, x_j) = \langle \phi(x_i), \phi(x_j) \rangle = \phi(x_i)^T \phi(x_j) \quad (103)$$

where $\langle \cdot, \cdot \rangle$ denotes the inner product in the Hilbert space and $\phi(\cdot)$ is the identity function.

The reason that this *kernel trick* is useful is that there are many classification/regression problems that are nonlinearly separable/regressable in the *input space*, which might be in a higher dimensionality *feature space* given a suitable mapping $x \rightarrow \phi(x)$.

5.1 Polynomial kernel

The *polynomial* kernel is defined as:

$$k(x_i, x_j) = (\gamma \langle x_i, x_j \rangle + r)^d \quad (104)$$

where γ define how far the influence of a single training example reaches (low values meaning ‘far’ and high values meaning ‘close’).




Figure 7: Polynomial SVM hyperplanes

5.2 Gaussian kernel


The *gaussian* kernel is defined as:

$$k(x_i, x_j) = \exp\left(-\frac{\|x_i - x_j\|_2^2}{2\sigma^2}\right) \quad (105)$$


or, equivalently:

$$k(x_i, x_j) = \exp(-\gamma\|x_i - x_j\|_2^2) \quad (106)$$

where $\gamma = \frac{1}{2\sigma^2}$ define how far the influence of a single training example reaches (low values meaning ‘far’ and high values meaning ‘close’).



(a) Gaussian SVC hyperplane



(b) Gaussian SVR hyperplane

Figure 8: Gaussian SVM hyperplanes

5.3 Laplacian kernel

The *laplacian* kernel is defined as:

$$k(x_i, x_j) = \exp\left(-\frac{\|x_i - x_j\|_1}{2\sigma^2}\right) \quad (107)$$

or, equivalently:

$$k(x_i, x_j) = \exp(-\gamma\|x_i - x_j\|_1) \quad (108)$$

where $\gamma = \frac{1}{2\sigma^2}$ define how far the influence of a single training example reaches (low values meaning ‘far’ and high values meaning ‘close’).




Figure 9: Laplacian SVM hyperplanes

6 Optimization Methods

In order to explain the *convergence rates* of the following optimization methods, we need to introduce some preliminary definitions about *convexity* and the *Lipschitz continuity* of a function [15].

Definition 1 (Convexity).

- (i) We say that a function $f : \mathbb{R}^m \rightarrow \mathbb{R}$ is convex if:

$$(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y) \quad \forall x, y \in \mathbb{R}^m, \lambda \in [0, 1]$$

- (ii) We say that a differentiable function $f : \mathbb{R}^m \rightarrow \mathbb{R}$ is convex if:

$$f(y) \geq f(x) + \langle \nabla f(x), y - x \rangle \quad \forall x, y \in \mathbb{R}^m$$

- (iii) We say that a twice differentiable function $f : \mathbb{R}^m \rightarrow \mathbb{R}$ is convex iff:

$$\nabla^2 f(x) \succeq 0 \quad \forall x \in \mathbb{R}^m$$

i.e., the Hessian matrix is *positive semidefinite*.

Definition 2 (Strict Convexity).

- (i) We say that a function $f : \mathbb{R}^m \rightarrow \mathbb{R}$ is strictly convex if:

$$(\lambda x + (1 - \lambda)y) < \lambda f(x) + (1 - \lambda)f(y) \quad \forall x, y \in \mathbb{R}^m, x \neq y, \lambda \in (0, 1)$$

- (ii) We say that a differentiable function $f : \mathbb{R}^m \rightarrow \mathbb{R}$ is strictly convex if:

$$f(y) > f(x) + \langle \nabla f(x), y - x \rangle \quad \forall x, y \in \mathbb{R}^m, x \neq y$$

- (iii) We say that a twice differentiable function $f : \mathbb{R}^m \rightarrow \mathbb{R}$ is strictly convex iff:

$$\nabla^2 f(x) \succ 0 \quad \forall x \in \mathbb{R}^m$$

i.e., the Hessian matrix is *positive definite*.

Definition 3 (Strong Convexity). We say that a function $f : \mathbb{R}^m \rightarrow \mathbb{R}$ is μ -strongly convex if the function:

$$g(x) = f(x) - \frac{\mu}{2} \|x\|^2$$

is convex for any $\mu > 0$. If f is differentiable this is also equivalent to:

$$f(y) \geq f(x) + \langle \nabla f(x), y - x \rangle + \frac{\mu}{2} \|y - x\|^2 \quad \forall x, y \in \mathbb{R}^m$$

and, if f is a twice differentiable function then f is μ -strongly convex iff:

$$\nabla^2 g(x) \succ 0 \quad \forall x \in \mathbb{R}^m$$

i.e., the Hessian matrix is *positive definite*, which is:

$$\nabla^2 f(x) \succeq \mu I \quad \forall x \in \mathbb{R}^m$$

i.e., all the eigenvalues of the Hessian matrix are lowerbounded by μI .

Definition 4 (L_f -Lipschitz continuity). We say that a function $f : \mathbb{R}^m \rightarrow \mathbb{R}$ is L_f -Lipschitz continuous if:

$$|f(x) - f(y)| \leq L_f \|x - y\| \quad \forall x, y \in \mathbb{R}^m$$

meaning that f is bounded above and below by a linear function.

Intuitively, L is a measure of how fast the function can change.

Finally, we say that a function $f : \mathbb{R}^m \rightarrow \mathbb{R}$ is locally L_f -Lipschitz continuous if for every x in \mathbb{R}^m there exists a neighborhood U of x such that f restricted to U is L_f -Lipschitz continuous. Every convex function is locally L_f -Lipschitz continuous.

Definition 5 (L-Lipschitz continuity). We say that a function $f : \mathbb{R}^m \rightarrow \mathbb{R}$ is L-Lipschitz gradient continuous if f is differentiable and:

$$\|\nabla f(x) - \nabla f(y)\| \leq L \|x - y\| \quad \forall x, y \in \mathbb{R}^m$$

that is equivalent to:

$$|f(y) - f(x) - \langle \nabla f(x), y - x \rangle| \leq \frac{L}{2} \|y - x\|^2 \quad \forall x, y \in \mathbb{R}^m$$

meaning that f is bounded above and below by a quadratic function.

Also, if f is a twice differentiable function this is equivalent to:

$$\nabla^2 f(x) \preceq LI \quad \forall x \in \mathbb{R}^m$$

i.e., all the eigenvalues of the Hessian matrix are upperbounded by L .

Note that if f is a μ -strongly convex function, we give the following Hessian bounds:

$$0 \prec \mu I \preceq \nabla^2 f(x) \preceq LI \quad \forall x \in \mathbb{R}^m$$

i.e., all the eigenvalues of the Hessian matrix are lowerbounded by μI and upperbounded by L .

Finally, we say that a function $f : \mathbb{R}^m \rightarrow \mathbb{R}$ is locally L-Lipschitz gradient continuous if for every x in \mathbb{R}^m there exists a neighborhood U of x such that f restricted to U is L-Lipschitz gradient continuous.

Definition 6 (Subgradient). Given a function $f : \mathbb{R}^m \rightarrow \mathbb{R}$ and $x \in \mathbb{R}^m$, we define a subgradient $g \in \mathbb{R}^m$ at x to be any point satisfying:

$$f(y) \geq f(x) + \langle g, y - x \rangle \quad \forall y \in \mathbb{R}^m$$

Subgradients always exist for convex function.

Theorem 7 (L_f -Lipschitz continuity for convex functions). *Let $f : \mathbb{R}^m \rightarrow \mathbb{R}$ be a convex function and let K be a closed and bounded set contained in the relative interior of the domain of f , i.e., $K \subset \mathbb{R}^m$. Then f is L_f -Lipschitz continuous on K , i.e.,:*

$$|f(x) - f(y)| \leq L_f \|x - y\| \quad \forall x, y \in K$$

In particular, f is bounded on K .

Proof. Let x and y be any two points in the set K . Since $\partial f(x)$ is nonempty, by using the subgradient inequality 6, it follows that:

$$f(y) \geq f(x) + \langle g, y - x \rangle \quad \forall g \in \partial f(x)$$

implying that:

$$f(x) - f(y) \leq \|g\| \|x - y\| \quad \forall g \in \partial f(x)$$

By definition, the set $\cup_{x \in K} \partial f(x)$ is nonempty and bounded, so that for some constant $L > 0$, we have:

$$\|g\| \leq L_f \quad \forall g \in \partial f(x) \quad \forall x \in K$$

and therefore:

$$f(x) - f(y) \leq L_f \|x - y\|$$

By exchanging the roles of x and y , we similarly obtain:

$$f(y) - f(x) \leq L_f \|x - y\|$$

and by combining the preceding two relations, we see that:

$$|f(x) - f(y)| \leq L_f \|x - y\|$$

showing that f is L_f -Lipschitz continuous over K . □

Note that this proof shows how to determine the Lipschitz constant L_f : it is the maximum subgradient norm, over all subgradients in $\cup_{x \in K} \partial f(x)$.

Strong convexity and L -Lipschitz continuity are related by Fenchel duality according to the following theorem, which proof is given in [16].

Theorem 8 (μ -strong convexity and L -Lipschitz continuity for convex functions). *A function f and its Fenchel dual f^* satisfy the following assertions:*

(i) *if f is μ -strongly convex, then f^* is $\frac{1}{\mu}$ -Lipschitz continuous.*

(ii) *if f is convex and L -Lipschitz continuous, then f^* is $\frac{1}{L}$ -strongly convex.*

Note that since f is convex and its epigraph is a closed convex set, $f^* = f$.

6.1 Gradient Descent for Primal formulations

The Gradient Descent algorithm is the simplest *first-order optimization* method that exploits the orthogonality of the gradient wrt the level sets to take a descent direction. In particular, it performs the following iterations:

Algorithm 1 Gradient Descent

Require: Function f to minimize

Require: Learning rate or step size $\alpha > 0$

function GRADIENTDESCENT(f, α)

 Initialize weight vector x_0

$t = 0$

while *not_convergence* **do**

$x_{t+1} = x_t - \alpha \partial f(x_t)$

▷ if f is differentiable then $\partial f(x_t) = \nabla f(x_t)$

$t = t + 1$

end while

return x_t

end function

Gradient Descent is based on full gradients, since at each iteration we compute the average gradient on the whole dataset:

$$\partial f(x) = \frac{1}{n} \sum_{i=1}^n \partial f_i(x)$$

The downside is that every step is very computationally expensive, $\mathcal{O}(nm)$ per iteration, where n is the number of samples in our dataset and m is the number of dimensions.

Since *Gradient Descent* becomes impractical when dealing with large datasets we introduce a stochastic version, called *Stochastic Gradient Descent*, which does not use the whole set of examples to compute the gradient at every step. By doing so, we can reduce computation all the way down to $\mathcal{O}(m)$ per iteration.

Algorithm 2 Stochastic Gradient Descent

Require: Function f to minimize

Require: Learning rate or step size $\alpha > 0$

Require: Batch size k

function STOCHASTICGRADIENTDESCENT(f, α, k)

 Initialize weight vector x_0

$t \leftarrow 0$

while *not_convergence* **do**

 Sample $(i_1, \dots, i_k) \sim \mathcal{U}^k(1, \dots, n)$

$x_{t+1} \leftarrow x_t - \alpha \frac{1}{k} \sum_{j=1}^k \partial f_{i_j}(x_t)$

▷ if f is differentiable then $\partial f_{i_j}(x_t) = \nabla f_{i_j}(x_t)$

$t \leftarrow t + 1$

end while

return x_t

end function

Note that in expectation, we converge like GD, since $\mathbb{E}_{i \sim \mathcal{U}(1, \dots, n)}[\partial f_i(x_t)] = \partial f(x_t)$, therefore, the expected iterate of SGD converges to the optimum.

Now, consider the SGD algorithm introduced previously but where each iteration is projected into the ball $\mathcal{B}(0, R)$ with radius $R > 0$ fixed. So, the following lower bounds on convergence rates are given.

Theorem 9 (Stochastic Gradient Descent convergence for convex functions). *Let $f : \mathbb{R}^m \rightarrow \mathbb{R}$ be a L -Lipschitz continuous convex function and assume that exists $b > 0$ satisfying:*

$$\|f_i(x)\| \leq b \quad \forall x \in \mathcal{B}(0, R)$$

Besides, assume that all minima of f belong to $\mathcal{B}(0, R)$. Then the Stochastic Gradient Descent with step size $\alpha = \frac{2R}{b\sqrt{k}}$ satisfies:

$$\mathbb{E} \left[f \left(\frac{1}{k} \sum_{t=1}^k x_t \right) \right] - f(x^*) \leq \frac{3Rb}{\sqrt{k}}$$

Theorem 10 (Stochastic Gradient Descent convergence for strongly convex functions). *Let $f : \mathbb{R}^m \rightarrow \mathbb{R}$ be a L -Lipschitz continuous, μ -strongly convex function and assume that exists $b > 0$ satisfying:*

$$\|f_i(x)\| \leq b \quad \forall x \in \mathcal{B}(0, R)$$

Besides, assume that all minima of f belong to $\mathcal{B}(0, R)$. Then the Stochastic Gradient Descent with step size $\alpha = \frac{2}{\mu(k+1)}$ satisfies:

$$\mathbb{E} \left[f \left(\frac{2}{k(k+1)} \sum_{t=1}^k t x_{t-1} \right) \right] - f(x^*) \leq \frac{2b^2}{\mu(k+1)}$$

SGD's convergence rate for L -Lipschitz continuous convex functions is $\mathcal{O}\left(\frac{1}{\sqrt{t}}\right)$ and $\mathcal{O}\left(\frac{1}{t}\right)$ for L -Lipschitz continuous and strongly convex functions. More iterations are needed to reach the same accuracy as GD, but the iterations are far cheaper.

6.1.1 Nonsmooth

First, consider a nonsmooth, i.e., nondifferentiable, convex function. So, the following lower bounds on convergence rates are given.

Theorem 11 (Subgradient Descent convergence for convex functions with Polyak's stepsize). *Let $f : \mathbb{R}^m \rightarrow \mathbb{R}$ be a L_f -Lipschitz continuous convex function. Then the Subgradient Descent with Polyak's step size $\alpha_t = \frac{f(x_t) - f(x^*)}{\|g_t\|^2}$ satisfies:*

$$f(x_t) - f(x^*) \leq \frac{L\|x_0 - x^*\|^2}{\sqrt{t+1}}$$

Unfortunately, Polyak's stepsize rule requires knowledge of $f(x^*)$, which is often unknown a priori, so we might often need simpler rule for setting stepsizes.

Theorem 12 (Subgradient Descent convergence for convex functions). *Let $f : \mathbb{R}^m \rightarrow \mathbb{R}$ be a L_f -Lipschitz continuous convex function. Then the Subgradient Descent with step size $\alpha_t = \frac{1}{\sqrt{t}}$ satisfies:*

$$f(x_t) - f(x^*) \leq \frac{\|x_0 - x^*\|^2 + L^2 \log t}{\sqrt{t}}$$

Theorem 13 (Subgradient Descent convergence for strongly convex functions). *Let $f : \mathbb{R}^m \rightarrow \mathbb{R}$ be a L_f -Lipschitz continuous and μ -strongly convex function. Then the Subgradient Descent with step size $\alpha_t = \frac{2}{\mu(t+1)}$ satisfies:*

$$f(x_t) - f(x^*) \leq \frac{2L^2}{\mu} \frac{1}{t+1}$$

In summary, the following *convergence rates* and *iterations complexities* are given:

Table 1: Subgradient Descent convergence rates and iterations complexities

	stepsize rule	convergence rate	iteration complexity
convex and L_f -Lipschitz	$\alpha = \frac{1}{\sqrt{t}}$	$\mathcal{O}\left(\frac{1}{\sqrt{t}}\right)$	$\mathcal{O}\left(\frac{1}{\epsilon^2}\right)$
strongly convex and L_f -Lipschitz	$\alpha = \frac{1}{t}$	$\mathcal{O}\left(\frac{1}{t}\right)$	$\mathcal{O}\left(\frac{1}{\epsilon}\right)$

Among algorithms that only use subgradient, these *convergence rates* cannot be further improved.

6.1.2 Smooth

Now, consider a smooth, i.e., differentiable, convex function. So, the following lower bounds on convergence rates are given.

Theorem 14 (Gradient Descent convergence for convex functions). *Let $f : \Re^m \rightarrow \Re$ be a L -Lipschitz continuous convex function. Then the Gradient Descent with step size $\alpha = 1/L$ satisfies:*

$$f(x_t) - f(x^*) \leq \frac{L\|x_0 - x^*\|^2}{2t}$$

Theorem 15 (Gradient Descent convergence for strongly convex functions). *Let $f : \Re^m \rightarrow \Re$ be a L -Lipschitz continuous and μ -strongly convex function. Then the Gradient Descent with step size $\alpha = 1/L$ satisfies:*

$$\begin{aligned} f(x_t) - f(x^*) &\leq \left(1 - \frac{\mu}{L}\right)^t \|f(x_0) - f(x^*)\|^2 \\ &= \left(1 - \frac{1}{\kappa}\right)^t \|f(x_0) - f(x^*)\|^2 \end{aligned}$$

where $\kappa = L/\mu$.

Theorem 16 (Gradient Descent convergence for convex quadratic functions). *Let $f : \Re^m \rightarrow \Re$ be a L -Lipschitz continuous and μ -strongly convex quadratic function. Then the Gradient Descent with step size $\alpha = \frac{2}{L + \mu}$ and momentum $\beta = \max\{|1 - \alpha\mu|, |1 - \alpha L|\}$ satisfies:*

$$\|x_t - x^*\| = \left(\frac{\kappa - 1}{\kappa + 1}\right)^t \|x_0 - x^*\|$$

where $\kappa = L/\mu$.

In summary, the following *convergence rates* and *iterations complexities* are given:

Table 2: Gradient Descent convergence rates and iterations complexities

	stepsize rule	convergence rate	iteration complexity
convex and L-Lipschitz	$\alpha = \frac{1}{L}$	$\mathcal{O}\left(\frac{1}{t}\right)$	$\mathcal{O}\left(\frac{1}{\epsilon}\right)$
strongly convex and L-Lipschitz	$\alpha = \frac{1}{L}$	$\mathcal{O}\left(\left(1 - \frac{1}{\kappa}\right)^t\right)$	$\mathcal{O}\left(\kappa \log \frac{1}{\epsilon}\right)$

6.1.3 Momentum

To mitigate the pathological zig-zagging by speeding up the *convergence rate* of the SGD method, we introduce two accelerated methods [1] and [2, 3] that exploits information from the history, i.e., past iterates, to add some inertia, i.e., the momentum, to yield smoother trajectory.

In the Polyak's method [1] the velocity vector v_t is calculated by applying the β momentum to the previous v_{t-1} displacement, and subtracting the gradient step to x_t .




Figure 10: Polyak's and Nesterov's Momentum

Algorithm 3 Polyak's Accelerated Gradient Descent or Polyak Heavy-Ball method

Require: Function f to minimize

Require: Learning rate or step size $\alpha > 0$

Require: Momentum $\beta \in [0, 1]$

function POLYAKACCELERATEDGRADIENTDESCENT(f, α, β)

 Initialize weight vector $x_1 \leftarrow x_0$ and velocity vector $v_0 \leftarrow 0$

$t \leftarrow 1$

while *not_convergence* **do**

$v_t = \beta v_{t-1} + \alpha \nabla f(x_t)$

$x_{t+1} = x_t - v_t$

$t \leftarrow t + 1$

end while

return x_t

end function

Theorem 17 (Polyak's Accelerated Gradient Descent convergence for convex quadratic functions). *Let $f : \mathbb{R}^m \rightarrow \mathbb{R}$ be a L -Lipschitz continuous and μ -strongly convex quadratic function. Then the Polyak's Accelerated Gradient Descent with step size $\alpha = \frac{4}{(\sqrt{L} + \sqrt{\mu})^2}$ and momentum $\beta = \max \{|1 - \sqrt{\alpha\mu}|, |1 - \sqrt{\alpha L}|\}^2$ satisfies:*

$$\|x_t - x^*\| = \left(\frac{\sqrt{\kappa} - 1}{\sqrt{\kappa} + 1} \right)^t \|x_0 - x^*\|$$

where $\kappa = L/\mu$.

Leveraging the idea of momentum introduced by Polyak, Nesterov introduced a slightly altered update rule that has been shown to converge not only for quadratic functions, but for general convex functions. In the Nesterov's method [2], instead, the velocity vector v_t is calculated by applying the β momentum to the previous v_{t-1} displacement, and subtracting the gradient step to $x_t + \beta v_{t-1}$, which is the point where the momentum term leads from x_t .

Algorithm 4 Nesterov's Accelerated Gradient Descent or Nesterov Heavy-Ball method

Require: Function f to minimize
Require: Learning rate $\alpha > 0$
Require: Momentum $\beta \in [0, 1)$
function NESTEROVACCELERATEDGRADIENTDESCENT(f, α, β)
 Initialize weight vector $x_1 \leftarrow x_0$ and velocity vector $v_0 \leftarrow 0$
 $t \leftarrow 1$
 while *not_convergence* **do**
 $\hat{x}_t \leftarrow x_t + \beta v_{t-1}$
 $v_t \leftarrow \beta v_{t-1} + \alpha \nabla f(\hat{x}_t)$
 $x_{t+1} \leftarrow x_t - v_t$
 $t \leftarrow t + 1$
 end while
 return x_t
end function

Comparing the algorithm 3 with the algorithm 4, we can see that Polyak's method evaluates the gradient before adding momentum, whereas Nesterov's algorithm evaluates it after applying momentum, which intuitively brings us closer to the minimum x^* , as showb in figure 10.

Theorem 18 (Nesterov's Accelerated Gradient Descent convergence for convex functions). *Let $f : \mathbb{R}^m \rightarrow \mathbb{R}$ be a L -Lipschitz continuous convex function. Then the Nesterov's Accelerated Gradient Descent with step size $\alpha = 1/L$ and momentum $\beta_{t+1} = t/(t+3)$ satisfies:*

$$f(x_t) - f(x^*) \leq \frac{2L\|x_0 - x^*\|^2}{(t+1)^2}$$

Theorem 19 (Nesterov's Accelerated Gradient Descent convergence for strongly convex functions). *Let $f : \mathbb{R}^m \rightarrow \mathbb{R}$ be a L -Lipschitz continuous and μ -strongly convex function. Then the Nesterov's Accelerated Gradient Descent with step size $\alpha = 1/L$ and momentum $\beta = \frac{\sqrt{\kappa} - 1}{\sqrt{\kappa} + 1}$ satisfies:*

$$\begin{aligned} f(x_t) - f(x^*) &\leq \left(1 - \sqrt{\frac{\mu}{L}}\right)^t \left(f(x_0) - f(x^*) + \frac{\mu\|x_0 - x^*\|^2}{2}\right) \\ &= \left(1 - \frac{1}{\sqrt{\kappa}}\right)^t \left(f(x_0) - f(x^*) + \frac{\mu\|x_0 - x^*\|^2}{2}\right) \end{aligned}$$

where $\kappa = L/\mu$.

In summary, the following *convergence rates* and *iterations complexities* are given:

Table 3: Nesterov's Accelerated Gradient Descent convergence rates and iterations complexities

	stepsize rule	convergence rate	iteration complexity
convex and L-Lipschitz	$\alpha = \frac{1}{L}$	$\mathcal{O}\left(\frac{1}{t^2}\right)$	$\mathcal{O}\left(\frac{1}{\sqrt{\epsilon}}\right)$
strongly convex and L-Lipschitz	$\alpha = \frac{1}{L}$	$\mathcal{O}\left(\left(1 - \frac{1}{\sqrt{\kappa}}\right)^t\right)$	$\mathcal{O}\left(\sqrt{\kappa} \log \frac{1}{\epsilon}\right)$

Note that in case of L -Lipschitz continuous and strongly convex functions, Nesterov's momentum gives the acceleration that we had with Polyak's momentum for quadratic functions. This is great because we get the guarantee for a more general class of functions, but these *convergence rates* cannot be further improved only using first-order information.

6.2 Sequential Minimal Optimization for Wolfe Dual formulations

The *Sequential Minimal Optimization (SMO)* [4] method is the most popular approach for solving the SVM QP problem without any extra Q matrix storage required by common QP methods. The advantage of SMO lies in the fact that it performs a series of two-point optimizations since we deal with just one equality constraint, so the Lagrange multipliers can be solved analytically.

6.2.1 Classification

At each iteration, SMO chooses two α_i to jointly optimize, let α_1 and α_2 , finds the optimal values for these multipliers and update the SVM to reflect these new values. In order to solve for two Lagrange multipliers, SMO first computes the constraints over these and then solves for the constrained minimum. Since there are only two multipliers, the box-constraints cause the Lagrange multipliers to lie within a box, while the linear equality constraint causes the Lagrange multipliers to lie on a diagonal line inside the box. So, the constrained minimum must lie there as shown in 11.



Figure 11: SMO for two Lagrange multipliers

In case of classification the ends of the diagonal line segment, i.e., the lower and upper bounds, can be expressed as follow if the target $y_1 \neq y_2$:

$$\begin{aligned} L &= \max(0, \alpha_2 - \alpha_1) \\ H &= \min(C, C + \alpha_2 - \alpha_1) \end{aligned} \quad (109)$$

or, alternatively, if the target $y_1 = y_2$:

$$\begin{aligned} L &= \max(0, \alpha_2 + \alpha_1 - C) \\ H &= \min(C, \alpha_2 + \alpha_1) \end{aligned} \quad (110)$$

The second derivative of the objective quadratic function along the diagonal line can be expressed as:

$$\eta = K(x_1, x_1) + K(x_2, x_2) - 2K(x_1, x_2) \quad (111)$$

that will be grather than zero if the kernel matrix will be positive definite, so there will be a minimum along the linear equality constraints that will be:

$$\alpha_2^{new} = \alpha_2 + \frac{y_2(E_1 - E_2)}{\eta} \quad (112)$$

where $E_i = y_i - y'_i$ is the error on the i -th training example and y'_i is the output of the SVC for the same.

Then, the box-constrained minimum is found by clipping the unconstrained minimum to the ends of the line segment:

$$\alpha_2^{new,clipped} = \begin{cases} H & \text{if } \alpha_2^{new} \geq H \\ \alpha_2^{new} & \text{if } L < \alpha_2^{new} < H \\ L & \text{if } \alpha_2^{new} \leq L \end{cases} \quad (113)$$

Finally, the value of α_1 is computed from the new clipped α_2 as:

$$\alpha_1^{new} = \alpha_1 + s(\alpha_2 - \alpha_2^{new,clipped}) \quad (114)$$

where $s = y_1 y_2$.

Since the *Karush-Kuhn-Tucker* conditions are necessary and sufficient conditions for optimality of a positive definite QP problem and the KKT conditions for the classification problem (19) are:

$$\begin{aligned} \alpha_i &= 0 \Leftrightarrow y_i y'_i \geq 1 \\ 0 < \alpha_i < C &\Leftrightarrow y_i y'_i = 1 \\ \alpha_i &= C \Leftrightarrow y_i y'_i \leq 1 \end{aligned} \quad (115)$$

the steps described above will be iterate as long as there will be an example that violates them.

After optimizing α_1 and α_2 , we select the threshold b such that the KKT conditions are satisfied for x_1 and x_2 . If, after optimization, α_1 is not at the bounds, i.e., $0 < \alpha_1 < C$, then the following threshold b_{up} is valid, since it forces the SVC to output y_1 when the input is x_1 :

$$b_{up} = E_1 + y_1(\alpha_1^{new} - \alpha_1)K(x_1, x_1) + y_2(\alpha_2^{new,clipped} - \alpha_2)K(x_1, x_2) + b \quad (116)$$

similarly, the following threshold b_{low} is valid if $0 < \alpha_2 < C$:

$$b_{low} = E_2 + y_1(\alpha_1^{new} - \alpha_1)K(x_1, x_2) + y_2(\alpha_2^{new,clipped} - \alpha_2)K(x_2, x_2) + b \quad (117)$$

If, after optimization, both $0 < \alpha_1 < C$ and $0 < \alpha_2 < C$ then both these thresholds are valid, and they will be equal; else, if both α_1 and α_2 are at the bounds, i.e., $\alpha_1 = 0$ or $\alpha_1 = C$ and $\alpha_2 = 0$ or $\alpha_2 = C$, then all the thresholds between b_{up} and b_{low} satisfy the KKT conditions, so we choose the threshold to be halfway in between b_{up} and b_{low} . This gives the complete equation for b :

$$b = \begin{cases} b_{up} & \text{if } 0 < \alpha_1 < C \\ b_{low} & \text{if } 0 < \alpha_2 < C \\ \frac{b_{up} + b_{low}}{2} & \text{otherwise} \end{cases} \quad (118)$$

Algorithm 5 Sequential Minimal Optimization for Classification

Require: Training examples matrix $X \in \mathbb{R}^{n \times m}$
Require: Training target vector $y \in \pm 1^n$
Require: Kernel matrix $K \in \mathbb{R}^{n \times n}$
Require: Regularization parameter $C > 0$
Require: Tolerance value tol for stopping criterion

function SMOCLASSIFIER(X, y, K, C, tol)
 Initialize the Lagrange multipliers vector $\alpha \in \mathbb{R}^n, \alpha \leftarrow 0$
 Initialize the empty set $I0 \leftarrow \{i : 0 < \alpha_i < C\}$
 Initialize the set $I1 \leftarrow \{i : y_i = +1, \alpha_i = 0\}$ to contain all the indices of the training examples of class +1
 Initialize the empty set $I2 \leftarrow \{i : y_i = -1, \alpha_i = C\}$
 Initialize the empty set $I3 \leftarrow \{i : y_i = +1, \alpha_i = C\}$
 Initialize the set $I4 \leftarrow \{i : y_i = -1, \alpha_i = 0\}$ to contain all the indices of the training examples of class -1
 Initialize $b_{up} \leftarrow -1$
 Initialize $b_{low} \leftarrow +1$
 Initialize the error cache vector $errors \in \mathbb{R}^n, errors \leftarrow 0$
while $num_changed > 0$ **or** $examine_all = True$ **do**
 $num_changed \leftarrow 0$
 $examine_all \leftarrow True$
 if $examine_all = True$ **then**
 for $i \leftarrow 0$ to n **do** ▷ loop over all training examples
 $num_changed \leftarrow num_changed + EXAMINEEXAMPLE(i)$
 end for
 else
 for i in $I0$ **do** ▷ loop over examples where α_i are not already at their bounds
 $num_changed \leftarrow num_changed + EXAMINEEXAMPLE(i)$
 if $b_{up} > b_{low} - 2tol$ **then** ▷ check if optimality on $I0$ is attained
 $num_changed \leftarrow 0$
 break
 end if
 end for
 end if
 if $examine_all = True$ **then**
 $examine_all \leftarrow False$
 else if $num_changed = 0$ **then**
 $examine_all \leftarrow True$
 end if
end while
 Compute b by (118)
return α, b
end function

Require: $i2$ -th Lagrange multiplier

```
function EXAMINEEXAMPLE( $i2$ )  
  if  $i2$  in  $I0$  then  
     $E_2 \leftarrow errors_{i2}$   
  else  
    Compute  $E_2$   
     $errors_{i2} \leftarrow E_2$   
    Update  $(b_{low}, i_{low})$  or  $(b_{up}, i_{up})$  using  $(E_2, i2)$   
  end if  
  if optimality is attained using current  $b_{low}$  and  $b_{up}$  then  
    return 0  
  else  
    Find an index  $i1$  to do joint optimization with  $i2$   
    if TAKESTEP( $i1, i2$ ) = True then  
      return 1  
    else  
      return 0  
    end if  
  end if  
end function
```


Require: $i1$ -th Lagrange multiplier

Require: $i2$ -th Lagrange multiplier

function TAKESTEP($i1, i2$)

if $i1 = i2$ **then**
 return False

end if

 Compute L and H using (109) or (110)

if $L = H$ **then**

return False

end if

 Compute η by (111)

\triangleright we assume that $\eta > 0$, i.e., the kernel matrix K is positive definite

if $\eta < 0$ **then**

 Choose $\alpha_2^{new,clipped}$ between L and H according to the largest value of the objective function at these points

else

 Compute α_2^{new} by (112)

 Compute $\alpha_2^{new,clipped}$ by (113)

end if

if changes in $\alpha_2^{new,clipped}$ are larger than some eps **then**

 Compute α_1^{new} by (114)

 Update $\alpha_2^{new,clipped}$ and α_1^{new}

for i in $I0$ **do**

 Update $errors_i$ using new Lagrange multipliers

end for

 Update α using new Lagrange multipliers

 Update $I0, I1, I2, I3$ and $I4$

 Update $errors_{i1}$ and $errors_{i2}$

for i in $I0 \cup \{i1, i2\}$ **do**

 Compute (i_{low}, b_{low}) by $b_{low} = \max\{errors_i : i \in I0 \cup I3 \cup I4\}$

 Compute (i_{up}, b_{up}) by $b_{up} = \min\{errors_i : i \in I0 \cup I1 \cup I2\}$

end for

return True

else

return False

end if

end function

6.2.2 Regression

In case of regression the bounds and the new multipliers $\alpha_1^{+,new}$ and $\alpha_2^{+,new}$ can be expressed as follows if $(\alpha_1^+ > 0 \text{ or } (\alpha_1^- = 0 \text{ and } E_1 - E_2 > 0))$ and $(\alpha_2^+ > 0 \text{ or } (\alpha_2^- = 0 \text{ and } E_1 - E_2 < 0))$:

$$\begin{aligned} L &= \max(0, \gamma - C) \\ H &= \min(C, \gamma) \end{aligned} \quad (119)$$

$$\alpha_2^{+,new} = \alpha_2^+ - \frac{E_1 - E_2}{\eta} \quad (120)$$

$$\alpha_1^{+,new} = \alpha_1^+ - (\alpha_2^{+,new,clipped} - \alpha_2^+) \quad (121)$$

or, if $(\alpha_1^+ > 0 \text{ or } (\alpha_1^- = 0 \text{ and } E_1 - E_2 > 2\epsilon))$ and $(\alpha_2^- > 0 \text{ or } (\alpha_2^+ = 0 \text{ and } E_1 - E_2 > 2\epsilon))$:

$$\begin{aligned} L &= \max(0, -\gamma) \\ H &= \min(C, -\gamma + C) \end{aligned} \quad (122)$$

$$\alpha_2^{-,new} = \alpha_2^- + \frac{(E_1 - E_2) - 2\epsilon}{\eta} \quad (123)$$

$$\alpha_1^{+,new} = \alpha_1^+ + (\alpha_2^{-,new,clipped} - \alpha_2^-) \quad (124)$$

or, if $(\alpha_1^- > 0 \text{ or } (\alpha_1^+ = 0 \text{ and } E_1 - E_2 < -2\epsilon))$ and $(\alpha_2^+ > 0 \text{ or } (\alpha_2^- = 0 \text{ and } E_1 - E_2 < -2\epsilon))$:

$$\begin{aligned} L &= \max(0, \gamma) \\ H &= \min(C, C + \gamma) \end{aligned} \quad (125)$$

$$\alpha_2^{+,new} = \alpha_2^+ - \frac{(E_1 - E_2) + 2\epsilon}{\eta} \quad (126)$$

$$\alpha_1^{-,new} = \alpha_1^- + (\alpha_2^{+,new,clipped} - \alpha_2^+) \quad (127)$$

or, finally, if $(\alpha_1^- > 0 \text{ or } (\alpha_1^+ = 0 \text{ and } E_1 - E_2 < 0))$ and $(\alpha_2^- > 0 \text{ or } (\alpha_2^+ = 0 \text{ and } E_1 - E_2 > 0))$:

$$\begin{aligned} L &= \max(0, -\gamma - C) \\ H &= \min(C, -\gamma) \end{aligned} \quad (128)$$

$$\alpha_2^{-,new} = \alpha_2^- + \frac{E_1 - E_2}{\eta} \quad (129)$$

$$\alpha_1^{-,new} = \alpha_1^- - (\alpha_2^{-,new,clipped} - \alpha_2^-) \quad (130)$$

where $\gamma = \alpha_1^+ - \alpha_1^- + \alpha_2^+ - \alpha_2^-$. Note that η and $\alpha_2^{+,new,clipped}$ or $\alpha_2^{-,new,clipped}$ are identical to (111) and (113) respectively.

The KKT conditions for the regression problem (68) are:

$$\begin{aligned} \alpha_i^+ - \alpha_i^- &= 0 \Leftrightarrow |y_i - y'_i| < \epsilon \\ -C < \alpha_i^+ - \alpha_i^- < C &\Leftrightarrow |y_i - y'_i| = \epsilon \\ \alpha_i^+ + \alpha_i^- &= C \Leftrightarrow |y_i - y'_i| > \epsilon \end{aligned} \quad (131)$$

so, the steps described above will be iterate as long as there will be an example that violates them.

In case of regression we select the threshold b as follows:

$$b_{up} = E_1 + ((\alpha_1^+ - \alpha_1^-) - (\alpha_1^{+,new} - \alpha_1^{-,new}))K(x_1, x_1) + ((\alpha_2^+ - \alpha_2^-) - (\alpha_2^{+,new,clipped} - \alpha_2^{-,new,clipped}))K(x_1, x_2) + b \quad (132)$$

$$b_{low} = E_2 + ((\alpha_1^+ - \alpha_1^-) - (\alpha_1^{+,new} - \alpha_1^{-,new}))K(x_1, x_2) + ((\alpha_2^+ - \alpha_2^-) - (\alpha_2^{+,new,clipped} - \alpha_2^{-,new,clipped}))K(x_2, x_2) + b \quad (133)$$

$$b = \begin{cases} b_{up} & \text{if } 0 < \alpha_1^+, \alpha_1^- < C \\ b_{low} & \text{if } 0 < \alpha_2^+, \alpha_2^- < C \\ \frac{b_{up} + b_{low}}{2} & \text{otherwise} \end{cases} \quad (134)$$

The improvements described in [5, 8] for classification and regression respectively are about the definition of subsets of multipliers to efficiently update them at each iteration by separating the multipliers at the bounds from those who can be further minimized.

Algorithm 6 Sequential Minimal Optimization for Regression

Require: Training examples matrix $X \in \mathbb{R}^{n \times m}$
Require: Training target vector $y \in \mathbb{R}^n$
Require: Kernel matrix $K \in \mathbb{R}^{n \times n}$
Require: Regularization parameter $C > 0$
Require: Epsilon-tube value $\epsilon \geq 0$ within which no penalty is associated in the epsilon-insensitive loss function
Require: Tolerance value tol for stopping criterion

function SMOREGRESSION($X, y, K, C, \epsilon, tol$)
 Initialize the Lagrange multipliers vector $\alpha^+ \in \mathbb{R}^n, \alpha^+ \leftarrow 0$
 Initialize the Lagrange multipliers vector $\alpha^- \in \mathbb{R}^n, \alpha^- \leftarrow 0$
 Initialize the empty set $I0 \leftarrow \{i : 0 < \alpha_i^+, \alpha_i^- < C\}$
 Initialize the set $I1 \leftarrow \{i : \alpha_i^+ = 0, \alpha_i^- = 0\}$ to contain all the indices of the training examples
 Initialize the empty set $I2 \leftarrow \{i : \alpha_i^+ = 0, \alpha_i^- = C\}$
 Initialize the empty set $I3 \leftarrow \{i : \alpha_i^+ = C, \alpha_i^- = 0\}$
 Initialize $i_{up} \leftarrow 0$ ▷ or any other target index i_{up} from the training examples
 Initialize $i_{low} \leftarrow 0$ ▷ or any other target index i_{low} from the training examples
 Initialize $b_{up} \leftarrow y_{i_{up}} + \epsilon$
 Initialize $b_{low} \leftarrow y_{i_{low}} - \epsilon$
 Initialize the error cache vector $errors \in \mathbb{R}^n, errors \leftarrow 0$
while $num_changed > 0$ **or** $examine_all = True$ **do**
 $num_changed \leftarrow 0$
 $examine_all \leftarrow True$
 if $examine_all = True$ **then**
 for $i \leftarrow 0$ to n **do** ▷ loop over all training examples
 $num_changed \leftarrow num_changed + EXAMINEEXAMPLE(i)$
 end for
 else
 for i in $I0$ **do** ▷ loop over examples where α_i^+ and α_i^- are not already at their bounds
 $num_changed \leftarrow num_changed + EXAMINEEXAMPLE(i)$
 if $b_{up} > b_{low} - 2tol$ **then** ▷ check if optimality on $I0$ is attained
 $num_changed \leftarrow 0$
 break
 end if
 end for
 end if
 if $examine_all = True$ **then**
 $examine_all \leftarrow False$
 else if $num_changed = 0$ **then**
 $examine_all \leftarrow True$
 end if
end while
 Compute b by (134)
return α^+, α^-, b
end function

Require: $i1$ -th Lagrange multiplier

Require: $i2$ -th Lagrange multiplier

function TAKESTEP($i1, i2$)

if $i1 = i2$ **then**

return False

end if

$finished = False$

while not $finished$ **do**

 Compute L and H using (119), (122), (125) or (128)

if $L < H$ **then**

 Compute η by (111) \triangleright we assume that $\eta > 0$, i.e., the kernel matrix K is positive definite

if $\eta < 0$ **then**

 Choose $\alpha_2^{+,new,clipped}$ or $\alpha_2^{-,new,clipped}$ between L and H according to the largest value of the objective function at these points

else

 Compute $\alpha_2^{+,new}$ or $\alpha_2^{-,new}$ using (120), (126) or (123), (129) respectively

 Compute $\alpha_2^{+,new,clipped}$ or $\alpha_2^{-,new,clipped}$ by (113)

end if

 Compute $\alpha_1^{+,new}$ or $\alpha_1^{-,new}$ using (121), (124) or (127), (130) respectively

if changes in $\alpha_2^{+,new,clipped}$, $\alpha_2^{-,new,clipped}$, $\alpha_1^{+,new}$ or $\alpha_1^{-,new}$ are larger than some eps **then**

 Update $\alpha_2^{+,new,clipped}$, $\alpha_2^{-,new,clipped}$, $\alpha_1^{+,new}$ or $\alpha_1^{-,new}$

end if

else

$finished = True$

end if

end while

if changes in $\alpha_2^{+,new,clipped}$, $\alpha_2^{-,new,clipped}$, $\alpha_1^{+,new}$ or $\alpha_1^{-,new}$ are larger than some eps **then**

for i in $I0$ **do**

 Update $errors_i$ using new Lagrange multipliers

end for

 Update α^+ and α^- using new Lagrange multipliers

 Update $I0, I1, I2$ and $I3$

 Update $errors_{i1}$ and $errors_{i2}$

for i in $I0 \cup \{i1, i2\}$ **do**

 Compute (i_{low}, b_{low}) by $b_{low} = \max\{errors_i : i \in I0 \cup I1 \cup I2\}$

 Compute and (i_{up}, b_{up}) by $b_{up} = \min\{errors_i : i \in I0 \cup I1 \cup I3\}$

end for

return True

else

return False

end if

end function

6.3 AdaGrad for Lagrangian Dual formulations

Due to the sparsity of the weight vector of the *Lagrangian dual*, i.e., the Lagrange multipliers, we might end up in a situation where some components of the gradient are very small and others large. This, in terms of *conditioning number*, i.e., $\kappa = L/\mu \gg 1$, means that the level sets of f are ellipsoid, i.e., we are dealing with an ill-conditioned problem. So, given a learning rate, a standard gradient descent approach might end up in a situation where it decreases too quickly the small weights or too slowly the large ones.

Another method, that is usually deprecated in ML applications due to its increased computational complexity, is Newton's method. Newton's method favors a much faster *convergence rate*, i.e., number of iterations, at the cost of being more expensive per iteration. For convex problems, the recursion is similar to the gradient descent algorithm:

$$x_{t+1} = x_t - \alpha H^{-1} \nabla f(x_t)$$

where α is often close to one (damped-Newton) or one, and H^{-1} denotes the Hessian of f at the current point, i.e., $\nabla^2 f(x_t)$.

The above suggest a general rule in optimization: find any preconditioner, in convex optimization it has to be positive semidefinite, that improves the performance of gradient descent in terms of iterations, but without wasting too much time to compute that preconditioner. The above result into:

$$x_{t+1} = x_t - \alpha P^{-1} \nabla f(x_t)$$

where P is the preconditioner. This idea is the basis of the BFGS quasi-Newton method.

The *AdaGrad* [6] algorithm is just a variant of preconditioned gradient descent, where P is selected to be a diagonal preconditioner matrix and is updated using the gradient information, in particular it is the diagonal approximation of the inverse of the square roots of gradient outer products, until the k -th iteration. The above lead to the algorithm:

Algorithm 7 AdaGrad

Require: Function f to minimize

Require: Learning rate or step size $\alpha > 0$

Require: Offset $\epsilon > 0$ to ensures not divide by 0

function ADAGRAD(f, α, ϵ)

Initialize weight vector x_0 and the squared accumulated gradients vector $s_t \leftarrow 0$

$t = 1$

while *not_convergence* **do**

$g_t \leftarrow \partial f(x_t)$

\triangleright if f is differentiable then $\partial f(x_t) = \nabla f(x_t)$

$s_t \leftarrow s_{t-1} + g_t^2$

$x_{t+1} \leftarrow x_t - \alpha P_t^{-1} g_t = x_t - \frac{\alpha}{\sqrt{s_t + \epsilon}} \odot g_t$ where $P_t \leftarrow \text{diag}(s_t + \epsilon)^{1/2}$

$t \leftarrow t + 1$

end while

return x_t

end function

In practical terms, *AdaGrad* addresses the problem of the sparse optimal by adaptively scaling the learning rate for each dimension with the magnitude of the gradients. Coordinates that routinely correspond to large gradients are scaled down significantly, whereas others with small gradients receive a much more gentle treatment.

6.4 Losses properties

Several losses and objectives have been presented in section 3 and 4. In our experiments, we will consider the following.

For what about the loss functions, two of them are nonsmooth convex functions, i.e., the *hinge* and the *epsilon-insensitive* losses for *classification* and *regression* tasks respectively, and linearly penalizes the misclassified points, i.e., \mathcal{L}_1 -SVM, meanwhile, their two *squared* versions are smooth, i.e., \mathcal{L}_2 -SVM, and quadratically penalizes the misclassified points.

Also, both the *margin-based* losses, i.e., the *hinge* and the *squared hinge* losses, are L_f -Lipschitz continuous; meanwhile, among the *distance-based* losses, the *epsilon-insensitive* loss is L_f -Lipschitz continuous but the *squared epsilon-insensitive* is not L_f -Lipschitz continuous, however it is convex and for this reason is locally L_f -Lipschitz continuous.

Also the regularization term, i.e., $\frac{1}{2}\|w\|^2$, is not L_f -Lipschitz continuous since it becomes arbitrarily steep as w approaches infinity, but it is strictly convex and for this reason is locally L_f -Lipschitz continuous. Clearly, its gradient, i.e., w , is not bounded since, again, they go to infinity as w goes to infinity, so this function is not L -Lipschitz continuous.

Since for our purposes, we need to show that our \mathcal{L}_1 -SVM objectives are L_f -Lipschitz continuous and the \mathcal{L}_2 -SVM objectives are L -Lipschitz continuous for the applicability of the convergence theorems, we will use the theorem 7 and 8 respectively.

In general, if the objective function of a quadratic programming problem is strictly convex, i.e., the associated Hessian matrix is positive definite, the solution is unique. Meanwhile, if the objective function is convex, there may be cases where the solution is nonunique.

Assume that the hard margin SVM has a solution, i.e., the given problem is separable in the feature space. Then, since the objective function of the primal problem is $\frac{1}{2}\|w\|^2$, which is strongly convex, the primal problem has a unique solution for w and b .

Since the \mathcal{L}_1 -SVM linearly penalizes the misclassified points, the primal objective function is convex. Likewise, the Hessian matrix of the dual objective function is positive semidefinite. Thus the primal and dual solutions may be nonunique. Meanwhile, the objective function of the primal problem for the \mathcal{L}_2 -SVM is strictly convex, due to the quadratic penalization of the misclassified points. Therefore, w and b are uniquely determined if we solve the primal or dual problem.

In summary, the following properties for the SVM's objectives are given:

Table 4: SVM's objectives properties for primal formulations

objective	smooth	Lipschitz continuous	convexity
\mathcal{L}_1 -SVC (13)	no	L_f -Lipschitz	convex
\mathcal{L}_2 -SVC (42)	yes	L-Lipschitz	strongly convex
\mathcal{L}_1 -SVR (61)	no	L_f -Lipschitz	convex
\mathcal{L}_2 -SVR (89)	yes	L-Lipschitz	strongly convex

And, according to the theoretical analysis, the following *convergence rates* are given for the primal and *Lagrangian dual* formulations respectively:

Table 5: SVM's objectives convergence rates for primal formulations

objective	SGD convergence rate	Polyak SGD convergence rate	Nesterov SGD convergence rate
\mathcal{L}_1 -SVM (13, 61)	$\mathcal{O}\left(\frac{m}{\sqrt{t}}\right)$	$\mathcal{O}\left(\frac{m}{\sqrt{t}}\right)$	$\mathcal{O}\left(\frac{m}{\sqrt{t}}\right)$
\mathcal{L}_2 -SVM (42, 89)	$\mathcal{O}\left(\frac{m}{t}\right)$	$\mathcal{O}\left(\frac{m}{t}\right)$	$\mathcal{O}\left(\frac{m}{t^2}\right)$

Table 6: SVM's objectives convergence rate for Lagrangian dual formulations

objective	AdaGrad convergence rate
\mathcal{L}_1 -SVM (27, 74) or (33, 80)	$\mathcal{O}\left(\frac{nm}{\sqrt{t}}\right)$
\mathcal{L}_2 -SVM (45, 92) or (50, 97)	$\mathcal{O}\left(\frac{nm}{t}\right)$

To improve the practical convergence of all the *Lagrangian dual* formulations we adding a strictly convex augmentation term, so by considering a general quadratic optimization problem subject to linear constraints, i.e., equality and inequality constraints, defined as:

$$\begin{aligned}
& \min_{\alpha} \quad \frac{1}{2} \alpha^T Q \alpha + q^T \alpha \\
& \text{subject to} \quad A\alpha = b \\
& \quad \quad \quad G\alpha \leq h \\
& \quad \quad \quad lb \leq \alpha \leq ub
\end{aligned} \tag{135}$$

or, equivalently:

$$\begin{aligned}
& \min_{\alpha} \quad \frac{1}{2} \alpha^T Q \alpha + q^T \alpha \\
& \text{subject to} \quad A\alpha = b \\
& \quad \quad \quad \hat{G}\alpha \leq \hat{h}
\end{aligned} \tag{136}$$

where $\hat{G} = \begin{bmatrix} G \\ -I \\ I \end{bmatrix}$ and $\hat{h} = [h \quad -lb \quad ub]$; we give the following *augmented Lagrangian dual*:

$$\begin{aligned}
& \max_{\mu, \lambda} \min_{\alpha} \quad \frac{1}{2} \alpha^T Q \alpha + q^T \alpha + \mu^T (A\alpha - b) + \lambda^T (\hat{G}\alpha - \hat{h}) + \frac{\rho}{2} \|A\alpha - b\|^2 + \frac{\rho}{2} \|\hat{G}\alpha - \hat{h}\|^2 \\
& \text{subject to} \quad \lambda \geq 0
\end{aligned} \tag{137}$$

with $\rho > 0$.

7 Experiments

The following experiments refer to *linearly* and *nonlinearly* separable generated datasets of size 100. All the training times refer to running on a laptop with an Intel i7-6700HQ (8) @ 3.500GHz and 31.2 GB of memory.

The Python source code is available at: github.com/dmeoli/optiml.

7.1 Support Vector Classifier

Below experiments are about the SVC for which I tested different values for the regularization hyperparameter C , i.e., from *soft* to *hard margin*, and in case of nonlinearly separable data also different *kernel functions* mentioned above.

The experiments about SVCs are available at:

github.com/dmeoli/optiml/blob/master/notebooks/optimization/CM_SVC_report_experiments.ipynb.


7.1.1 Hinge loss

Primal formulation The experiments results shown in 7 referred to *Stochastic Gradient Descent* algorithm are obtained with α , i.e., the *learning rate* or *step size*, setted to 0.02 and β , i.e., the *momentum*, equal to 0.5. The optimization process is stopped if after 5 iterations the function value does not improve by at least $1e-8$.

Table 7: Primal \mathcal{L}_1 -SVC results

solver	momentum	C	fit_time	accuracy	n_iter	n_sv
sgd	none	0.1	7.062483	0.975	7196	36
		1.0	3.250063	0.985	4092	15
		10.0	0.795309	0.980	947	10
	polyak	0.1	4.450933	0.975	3589	37
		1.0	1.794078	0.985	2098	16
		10.0	0.367355	0.980	467	10
	nesterov	0.1	3.677246	0.975	4344	37
		1.0	1.788335	0.985	2115	16
		10.0	0.414720	0.980	472	10
liblinear	-	0.1	0.007178	0.980	31	37
		1.0	0.004740	0.985	332	16
		10.0	0.002364	0.985	1183	7

The results provided from the *custom* implementation, i.e., the SGD with different momentum settings, are strongly similar to those of *sklearn* implementation, i.e., *liblinear* [10] implementation, in terms of *accuracy* score. More training data points are selected as *support vectors* from the SGD solver but it always requires lower iterations, i.e., epochs, to achieve the same *numerical precision*. *Polyak* and *Nesterov* momentums always perform lower iterations as expected from the theoretical analysis of the convergence rate.

Figure 12: SGD Convergence for the Primal formulation of the \mathcal{L}_1 -SVC

Linear Dual formulations The experiments results shown in 9 are obtained with α , i.e., the *learning rate* or *step size*, setted to 1 for the *AdaGrad* algorithm. Note that the *unreg.bias* and *reg.bias* duals refers to the augmented dual formulations (137) of the problems (19) and (26) respectively with ρ equals to 1. The optimization process is stopped if the primal-dual weight vector does not change by at least $1e-8$ between two consecutive iterations.

Table 8: Wolfe Dual linear \mathcal{L}_1 -SVC results

		fit_time	accuracy	n_iter	n_sv
solver	C				
smo	0.1	0.158289	0.985	33	38
	1.0	0.209209	0.980	62	17
	10.0	0.352789	0.980	295	10
libsvm	0.1	0.007530	0.985	37	38
	1.0	0.008367	0.985	243	17
	10.0	0.003456	0.985	194	10
cvxopt	0.1	0.079530	0.985	9	38
	1.0	0.073912	0.980	10	17
	10.0	0.071737	0.980	10	11

For what about the linear *Wolfe dual* formulation we can immediately notice as higher *regularization hyperparameter* C makes the model harder, so the *custom* implementation of the SMO algorithm and also the *sklearn* implementation, i.e., *libsvm* [11] implementation, needs to perform more iterations to achieve the same *numerical precision*; meanwhile the *cvxopt* [12] seems to be insensitive to the increasing complexity of the model. The results in terms of *accuracy* and number of *support vectors* are strongly similar to each others.

Table 9: Lagrangian Dual linear \mathcal{L}_1 -SVC results

		fit_time	accuracy	n_iter	n_sv
dual	C				
reg_bias	0.1	101.691994	0.985	86461	37
	1.0	150.951210	0.980	89394	17
	10.0	156.155644	0.980	94941	10
unreg_bias	0.1	302.375478	0.985	212154	38
	1.0	157.811984	0.980	94542	17
	10.0	167.588434	0.980	97199	10

For what about the linear *Lagrangian dual* formulation we can see as it seems to be insensitive to the increasing complexity of the model in terms of number of *iterations* but it tends to select many training data points as *support vectors*.

Nonlinear Dual formulations The experiments results shown in 10 and 11 are obtained with d and r hyperparameters equal to 3 and 1 respectively for the *polynomial* kernel; *gamma* is setted to ‘scale’ for both *polynomial* and *gaussian* kernels. The experiments results shown in 11 are obtained with α , i.e., the *learning rate* or *step size*, setted to 1 for the *AdaGrad* algorithm. Note that the *unreg_bias* and *reg_bias* duals refers to the augmented dual formulations (137) of the problems (19) and (26) respectively with ρ equals to 1. The optimization process is stopped if the primal-dual weight vector does not change by at least $1e-8$ between two consecutive iterations.


Table 10: Wolfe Dual nonlinear \mathcal{L}_1 -SVC results

			fit_time	accuracy	n_iter	n_sv
solver	kernel	C				
smo	gaussian	0.1	0.878962	1.0000	65	222
		1.0	0.860773	1.0000	76	48
		10.0	0.525582	1.0000	29	13
	poly	0.1	1.096614	0.8675	121	142
		1.0	0.999406	0.6825	143	30
		10.0	0.680124	0.9475	65	10
libsvm	gaussian	0.1	0.011811	1.0000	131	222
		1.0	0.007152	1.0000	252	50
		10.0	0.003799	1.0000	134	13
	poly	0.1	0.022929	1.0000	210	143
		1.0	0.010823	1.0000	233	30
		10.0	0.003304	1.0000	118	10
cvxopt	gaussian	0.1	0.426778	1.0000	10	222
		1.0	0.568489	1.0000	10	49
		10.0	0.557587	1.0000	10	14
	poly	0.1	0.487677	0.8575	10	143
		1.0	0.693849	0.6775	10	31
		10.0	0.610578	0.9475	10	10

Table 11: Lagrangian Dual nonlinear \mathcal{L}_1 -SVC results

dual	kernel	C	fit_time	accuracy	n_iter	n_sv
reg_bias	poly	0.1	676.281088	0.8575	132326	143
		1.0	872.864681	0.6775	224601	31
		10.0	650.963683	0.9475	140304	10
	rbf	0.1	330.997649	1.0000	59727	222
		1.0	2300.525620	1.0000	500000	50
		10.0	1059.727331	1.0000	185902	13
unreg_bias	poly	0.1	1255.429093	0.8600	274706	143
		1.0	691.678106	0.6850	153439	30
		10.0	695.467444	0.9475	147822	10
	rbf	0.1	790.552171	1.0000	166776	222
		1.0	2449.294145	1.0000	500000	50
		10.0	1790.555752	1.0000	318882	13

The same considerations made for the previous linear *Wolfe dual* and *Lagrangian dual* formulations are confirmed also in the nonlinearly separable case. In this setting the complexity of the model coming with higher C regularization values seems to be not paying a tradeoff in terms of the number of *iterations* of the algorithm and, moreover, the *reg_bias Lagrangian dual* formulation seems to perform better wrt the *unreg_bias* formulation, both tends to select even more training data points as *support vectors*.

Figure 13: AdaGrad convergence for the Lagrangian Dual formulation of the nonlinear \mathcal{L}_1 -SVC

7.1.2 Squared Hinge loss

Primal formulation The experiments results shown in 12 referred to *Stochastic Gradient Descent* algorithm are obtained with α , i.e., the *learning rate* or *step size*, setted to 0.02 and β , i.e., the *momentum*, equal to 0.5. The optimization process is stopped if after 5 iterations the function value does not improve by at least $1e-8$.

Table 12: Primal \mathcal{L}_2 -SVC results

solver	momentum	C	fit_time	accuracy	n_iter	n_sv
sgd	none	0.1	4.478252	0.980	5661	49
		1.0	4.060005	0.980	5165	25
		10.0	2.564839	0.980	1895	19
	polyak	0.1	2.756118	0.980	3287	49
		1.0	2.298601	0.980	2827	25
		10.0	0.023973	0.975	10	20
	nesterov	0.1	2.714415	0.980	3291	49
		1.0	2.367481	0.980	2835	25
		10.0	0.731884	0.980	849	19
liblinear	-	0.1	0.003543	0.980	52	46
		1.0	0.005556	0.980	563	25
		10.0	0.008653	0.980	3129	19

Again, the results provided from the *custom* implementation, i.e., the SGD with different momentum settings, are strongly similar to those of *sklearn* implementation, i.e., *liblinear* [10] implementation, in terms of *accuracy* score. More training data points are selected as *support vectors* from the SGD solver but it always requires even lower iterations, i.e., epochs, to achieve the same *numerical precision*. *Polyak* and *Nesterov* momentums always perform lower iterations as expected from the theoretical analysis of the convergence rate.




Figure 14: SGD convergence for the Primal formulation of the \mathcal{L}_2 -SVC

Linear Dual formulations The experiments results shown in 13 are obtained with α , i.e., the *learning rate* or *step size*, setted to 1 for the *AdaGrad* algorithm. Note that the *unreg.bias* and *reg.bias* duals refers to the augmented dual formulations (137) of the problems (43) and (44) respectively with ρ equals to 1. The optimization process is stopped if the primal-dual weight vector does not change by at least $1e-8$ between two consecutive iterations.

Table 13: Lagrangian Dual linear \mathcal{L}_2 -SVC results

dual	C	fit_time	accuracy	n_iter	n_sv
reg_bias	0.1	15.297997	0.98	8582	46
	1.0	82.010417	0.98	44372	25
	10.0	401.840846	0.98	199427	19
unreg_bias	0.1	17.460816	0.98	9010	47
	1.0	78.865659	0.98	45217	25
	10.0	432.222420	0.98	211808	19


For what about the linear *Lagrangian dual* formulation we can see as it seems to be insensitive to the increasing complexity of the model in terms of number of *iterations* but it tends to select many training data points as *support vectors*.

Nonlinear Dual formulations The experiments results shown in 14 are obtained with d and r hyperparameters equal to 3 and 1 respectively for the *polynomial* kernel; γ is setted to ‘*scale*’ for both *polynomial* and *gaussian* kernels. The experiments results shown in 11 are obtained with α , i.e., the *learning rate* or *step size*, setted to 1 for the *AdaGrad* algorithm. Note that the *unreg.bias* and *reg.bias* duals refers to the augmented dual formulations (137) of the problems (43) and (44) respectively with ρ equals to 1. The optimization process is stopped if the primal-dual weight vector does not change by at least $1e-8$ between two consecutive iterations.

Table 14: Lagrangian Dual nonlinear \mathcal{L}_2 -SVC results

dual	kernel	C	fit_time	accuracy	n_iter	n_sv
reg_bias	poly	0.1	39.962172	0.8550	14277	233
		1.0	264.053112	0.6950	69608	80
		10.0	539.714278	0.7300	128384	16
	rbf	0.1	5.142467	1.0000	2377	345
		1.0	54.398084	1.0000	15288	130
		10.0	421.391269	1.0000	102765	33
unreg_bias	poly	0.1	45.310120	0.8625	14966	234
		1.0	283.749638	0.6950	73114	80
		10.0	573.433257	0.7300	135428	16
	rbf	0.1	11.287864	1.0000	4916	344
		1.0	99.375446	1.0000	27865	130
		10.0	703.632651	1.0000	172343	33

The same considerations made for the previous linear *Lagrangian dual* formulations are confirmed also in the nonlinearly separable case. In this setting the complexity of the model coming with higher C regularization values seems to be not paying a tradeoff in terms of the number of *iterations* of the algorithm and, moreover, the *reg.bias Lagrangian dual* formulation seems to perform better wrt the *unreg.bias* formulation, both tends to select even more training data points as *support vectors*.

Figure 15: AdaGrad convergence for the Lagrangian Dual formulation of the nonlinear \mathcal{L}_2 -SVC

7.2 Support Vector Regression

Below experiments are about the SVR for which I tested different values for regularization hyperparameter C , i.e., from *soft* to *hard margin*, the ϵ penalty value and in case of nonlinearly separable data also different *kernel functions* mentioned above.

The experiments about SVRs are available at:

`github.com/dmeoli/optimpl/blob/master/notebooks/optimization/CM_SVR_report_experiments.ipynb`.


7.2.1 Epsilon-insensitive loss

Primal formulation The experiments results shown in 15 referred to *Stochastic Gradient Descent* algorithm are obtained with α , i.e., the *learning rate* or *step size*, setted to 0.02 and β , i.e., the *momentum*, equal to 0.4. The optimization process is stopped if after 5 iterations the function value does not improve by at least $1e-8$.

Table 15: Primal \mathcal{L}_1 -SVR results

				fit_time	r2	n_iter	n_sv
solver	momentum	C	epsilon				
sgd	none	1	0.1	13.207407	0.954298	16161	100
			0.2	9.523358	0.954544	12267	99
			0.3	11.594773	0.955424	13665	99
		10	0.1	0.690761	0.983893	806	98
			0.2	0.722907	0.983891	884	98
			0.3	0.756686	0.983884	958	97
		100	0.1	0.121053	0.984034	85	97
			0.2	0.088140	0.984047	96	98
			0.3	0.156799	0.984056	109	98
	polyak	1	0.1	7.895537	0.954321	9874	100
			0.2	5.816864	0.954549	7400	99
			0.3	6.935173	0.955424	8200	99
		10	0.1	0.489926	0.983893	487	97
			0.2	0.503936	0.983891	535	98
			0.3	0.483785	0.983885	569	98
		100	0.1	0.090085	0.984030	48	98
			0.2	0.108489	0.984046	56	98
			0.3	0.114756	0.984055	61	97
	nesterov	1	0.1	8.996545	0.954310	9785	100
			0.2	6.001156	0.954546	7382	99
			0.3	7.318327	0.955424	8198	99
		10	0.1	0.803194	0.983892	489	97
			0.2	0.457146	0.983890	533	97
			0.3	0.561535	0.983884	579	98
		100	0.1	0.097744	0.984031	61	98
			0.2	0.120314	0.984047	58	98
			0.3	0.107943	0.984057	62	98
liblinear	-	1	0.1	0.019717	0.954684	12	100
			0.2	0.001534	0.955112	10	99
			0.3	0.001302	0.955415	10	97
		10	0.1	0.002194	0.983893	57	99
			0.2	0.001483	0.983890	69	98
			0.3	0.001968	0.983906	142	97
		100	0.1	0.001972	0.984023	980	97
			0.2	0.006431	0.984028	1340	97
			0.3	0.002328	0.984051	2886	97

The results provided from the *custom* implementation, i.e., the SGD with different momentum settings, are strongly similar to those of *sklearn* implementation, i.e., *liblinear* [10] implementation, in terms of $r2$ score, except in case of C regularization hyperparameter equals to 1 for which those of SGD are lower. Moreover, the SGD solver always requires lower iterations, i.e., epochs, for higher C regularization values, i.e., for C equals to 10 or 100, to achieve the same *numerical precision*. Again, *Polyak* and *Nesterov* momentums always perform lower iterations as expected from the theoretical analysis of the convergence rate. The results in terms of *support vectors* are strongly similar to each others.

Figure 16: SGD convergence for the Primal formulation of the \mathcal{L}_1 -SVR

Linear Dual formulations The experiments results shown in 17 are obtained with α , i.e., the *learning rate* or *step size*, setted to 1 for the *AdaGrad* algorithm. Note that the *unreg-bias* and *reg-bias* duals refers to the augmented dual formulations (137) of the problems (69) and (73) respectively with ρ equals to 1. The optimization process is stopped if the primal-dual weight vector does not change by at least $1e-8$ between two consecutive iterations.

Table 16: Wolfe Dual linear \mathcal{L}_1 -SVR results


solver	C	epsilon	fit_time	r2	n_iter	n_sv
smo	1	0.1	0.049352	0.954396	10	100
		0.2	0.026155	0.954546	15	100
		0.3	0.090050	0.955429	13	99
	10	0.1	0.201986	0.983893	44	99
		0.2	0.091500	0.983893	48	99
		0.3	0.084329	0.983893	41	99
	100	0.1	0.826075	0.984071	623	98
		0.2	0.409304	0.984088	157	98
		0.3	0.521488	0.984103	334	98
libsvm	1	0.1	0.009594	0.954393	79	100
		0.2	0.005095	0.954543	82	100
		0.3	0.029558	0.955424	78	99
	10	0.1	0.033350	0.983892	206	99
		0.2	0.005173	0.983890	219	99
		0.3	0.009250	0.983885	216	99
	100	0.1	0.019970	0.984028	2239	98
		0.2	0.006288	0.984041	1189	98
		0.3	0.003961	0.984051	1366	98
cvxopt	1	0.1	0.122198	0.954685	9	100
		0.2	0.152970	0.954849	9	100
		0.3	0.066905	0.955429	10	100
	10	0.1	0.144911	0.983893	9	100
		0.2	0.056698	0.983893	8	100
		0.3	0.045109	0.983893	8	100
	100	0.1	0.094987	0.984071	9	100
		0.2	0.070957	0.984088	9	100
		0.3	0.095619	0.984103	8	100

For what about the linear *Wolfe dual* formulation we can immediately notice as higher *regularization hyperparameter* C and lower ϵ values makes the model harder, so the *custom* implementation of the SMO algorithm and also the *sklearn* implementation, i.e., *libsvm* [11] implementation, needs to perform more iterations to achieve the same *numerical precision*; meanwhile, again, the *cvxopt* [12] seems to be insensitive to the increasing complexity of the model. The results in terms of $r2$ and number of *support vectors* are strongly similar to each others.

Table 17: Lagrangian Dual linear \mathcal{L}_1 -SVR results

dual	C	epsilon	fit_time	r2	n_iter	n_sv
reg_bias	1	0.1	61.665806	0.954685	30114	100
		0.2	49.460681	0.954845	29829	100
		0.3	47.498286	0.955429	29479	99
	10	0.1	71.959409	0.983893	35304	99
		0.2	74.538032	0.983893	35372	99
		0.3	81.454717	0.983893	34782	99
	100	0.1	330.733877	0.984071	128527	98
		0.2	385.085790	0.984088	165358	98
		0.3	712.218703	0.984103	309382	98
unreg_bias	1	0.1	69.608550	0.954396	35767	100
		0.2	179.243881	0.954546	101652	100
		0.3	1409.724278	0.955429	637390	99
	10	0.1	217.236034	0.983893	85088	99
		0.2	196.254325	0.983893	85973	99
		0.3	229.680320	0.983893	92590	99
	100	0.1	1800.660170	0.984071	785943	98
		0.2	2009.937387	0.984088	847207	98
		0.3	2332.760692	0.984103	969237	98

For what about the linear *Lagrangian dual* formulation we can see as it seems to be insensitive to the increasing complexity of the model in terms of number of *iterations* and require many *iterations* wrt the *Wolfe dual* formulation.

Figure 17: AdaGrad convergence for the Lagrangian Dual formulation of the linear \mathcal{L}_1 -SVR

Nonlinear Dual formulations The experiments results shown in 18 and 19 are obtained with d and r hyperparameters both equal to 3 for the *polynomial* kernel; γ is setted to ‘*scale*’ for both *polynomial* and *gaussian* kernels. The experiments results shown in 11 are obtained with α , i.e., the *learning rate* or *step size*, setted to 1 for the *AdaGrad* algorithm. Note that the *unreg-bias* and *reg-bias* duals refers to the augmented dual formulations (137) of the problems (69) and (73) respectively with ρ equals to 1. The optimization process is stopped if the primal-dual weight vector does not change by at least $1e-8$ between two consecutive iterations.


Table 18: Wolfe Dual nonlinear \mathcal{L}_1 -SVR formulation results


solver	kernel	C	epsilon	fit_time	r2	n_iter	n_sv
smo	gaussian	1	0.1	0.134567	0.988249	61	17
			0.2	0.098772	0.924439	18	7
			0.3	0.058488	0.882880	17	5
		10	0.1	0.628905	0.989828	289	18
			0.2	0.073629	0.924770	27	6
			0.3	0.034380	0.883067	13	5
		100	0.1	10.473704	0.899765	4835	17
			0.2	0.087404	0.924770	27	6
			0.3	0.023395	0.883067	13	5
	laplacian	1	0.1	0.190778	0.972858	23	23
			0.2	0.103656	0.942216	21	13
			0.3	0.073561	0.866739	17	9
		10	0.1	0.308953	0.989399	19	22
			0.2	0.110892	0.941932	17	13
			0.3	0.087622	0.866472	13	9
		100	0.1	0.237147	0.989399	19	22
			0.2	0.083979	0.941932	17	13
			0.3	0.080471	0.866472	13	9
libsvm	gaussian	1	0.1	0.009212	0.990088	96	17
			0.2	0.007524	0.977763	36	7
			0.3	0.002327	0.945601	24	5
		10	0.1	0.006896	0.990493	616	18
			0.2	0.008819	0.980673	39	6
			0.3	0.002239	0.945601	24	5
		100	0.1	0.010395	0.990496	9854	18
			0.2	0.002149	0.980673	39	6
			0.3	0.001906	0.945601	24	5
	laplacian	1	0.1	0.019776	0.990050	47	23
			0.2	0.006970	0.969067	28	13
			0.3	0.005510	0.924296	22	9
		10	0.1	0.002124	0.990777	47	23
			0.2	0.007089	0.969103	31	13
			0.3	0.002430	0.924237	22	9
		100	0.1	0.005186	0.990777	47	23
			0.2	0.002676	0.969103	31	13
			0.3	0.005486	0.924237	22	9
cvxopt	gaussian	1	0.1	0.106479	0.988117	10	17
			0.2	0.060167	0.924679	10	7
			0.3	0.048079	0.883386	10	5
		10	0.1	0.126220	0.989956	10	18
			0.2	0.085939	0.925595	10	6
			0.3	0.109721	0.883386	10	5
		100	0.1	0.097486	0.990216	10	40
			0.2	0.076151	0.925595	10	6
			0.3	0.095276	0.883386	10	5
	laplacian	1	0.1	0.094410	0.977836	9	24
			0.2	0.096828	0.942110	9	13
			0.3	0.145580	0.866633	9	9
		10	0.1	0.112691	0.984378	10	24
			0.2	0.133206	0.942110	10	13
			0.3	0.098581	0.866633	10	9
		100	0.1	0.071575	0.984378	10	24
			0.2	0.086546	0.955697	10	14
			0.3	0.076675	0.888440	10	10

Table 19: Lagrangian Dual nonlinear \mathcal{L}_1 -SVR results

dual	kernel	C	epsilon	fit_time	r2	n_iter	n_sv
reg_bias	poly	1	0.1	3799.855584	0.759735	4345854	45
			0.2	4101.699981	-0.593191	3979752	14
			0.3	3736.716269	0.950884	3667706	12
		10	0.1	3227.341450	0.889292	4273682	43
			0.2	3066.053793	0.932512	3833638	17
			0.3	2973.237179	0.405079	3666349	14
		100	0.1	5389.059721	-0.454737	7000000	43
			0.2	3120.464042	0.932512	3833638	17
			0.3	5343.459601	0.405079	3666349	14
	rbf	1	0.1	47.461531	0.988828	53076	19
			0.2	19.048212	0.924431	17252	7
			0.3	10.817777	0.894204	13637	6
		10	0.1	517.461881	0.989716	636053	18
			0.2	14.603766	0.916999	17749	7
			0.3	12.477623	0.921671	12840	10
		100	0.1	4853.336714	0.980079	5920433	18
			0.2	13.425201	0.916999	17749	7
			0.3	10.280788	0.921671	12840	10
unreg_bias	poly	1	0.1	3861.455070	0.878100	4300698	43
			0.2	3443.375626	-1.808568	3718479	15
			0.3	2860.392630	0.944686	3569608	12
		10	0.1	3349.425094	0.845000	4372132	45
			0.2	3226.271276	0.849851	3940737	15
			0.3	3054.851898	0.931490	3624333	13
		100	0.1	5480.802903	-0.461495	7000000	43
			0.2	3184.266593	0.849851	3940737	15
			0.3	3287.755984	0.931490	3624333	13
	rbf	1	0.1	65.452607	0.987761	74401	20
			0.2	35.227437	0.903375	31217	9
			0.3	19.617716	0.911973	25336	7
		10	0.1	482.568146	0.988548	598918	19
			0.2	24.508186	0.929011	31397	7
			0.3	20.715078	0.911975	24386	7
		100	0.1	4407.457835	0.886070	5388026	19
			0.2	24.151294	0.929011	31397	7
			0.3	19.131887	0.911975	24386	7

The same considerations made for the previous linear *Wolfe dual* and *Lagrangian dual* formulations are confirmed also in the nonlinearly separable case. In this setting, the complexity of the model coming with higher C regularization hyperparameters and lower ϵ values pays a larger tradeoff in terms of the number of *iterations* of the algorithm.

Figure 18: AdaGrad convergence for the Lagrangian Dual formulation of the Polynomial \mathcal{L}_1 -SVR

Figure 19: AdaGrad convergence for the Lagrangian Dual formulation of the Gaussian \mathcal{L}_1 -SVR


7.2.2 Squared Epsilon-insensitive loss

Primal formulation The experiments results shown in 20 referred to *Stochastic Gradient Descent* algorithm are obtained with α , i.e., the *learning rate* or *step size*, setted to 0.02 and β , i.e., the *momentum*, equal to 0.4. The optimization process is stopped if after 5 iterations the function value does not improve by at least $1e-8$.

Table 20: Primal \mathcal{L}_2 -SVR results

solver	momentum	C	epsilon	fit_time	r2	n_iter	n_sv
sgd	none	1	0.1	2.610538	0.984109	3283	100
			0.2	2.536275	0.984109	3294	100
			0.3	2.622454	0.984109	3321	98
		10	0.1	0.373240	0.984133	409	98
			0.2	0.373901	0.984133	410	98
			0.3	0.406998	0.984133	411	98
		100	0.1	0.086858	0.984133	47	98
			0.2	0.064771	0.984133	47	98
			0.3	0.034043	0.984133	47	98
	polyak	1	0.1	1.642638	0.984109	2011	100
			0.2	1.685593	0.984109	2018	100
			0.3	1.856611	0.984109	2035	98
		10	0.1	0.233500	0.984133	241	98
			0.2	0.233849	0.984133	242	98
			0.3	0.260583	0.984133	243	98
		100	0.1	0.062701	0.984133	40	98
			0.2	0.062025	0.984133	40	98
			0.3	0.062099	0.984133	40	98
	nesterov	1	0.1	2.522627	0.984109	2015	100
			0.2	1.593459	0.984109	2022	100
			0.3	1.696347	0.984109	2039	98
		10	0.1	0.224427	0.984133	247	98
			0.2	0.231077	0.984133	248	98
			0.3	0.251711	0.984133	248	98
		100	0.1	0.041334	0.984133	27	98
			0.2	0.036721	0.984133	27	98
			0.3	0.057086	0.984133	27	98
liblinear	-	1	0.1	0.002193	0.984109	84	100
			0.2	0.003775	0.984109	84	100
			0.3	0.006009	0.984109	84	98
		10	0.1	0.004759	0.984133	778	98
			0.2	0.005728	0.984133	773	98
			0.3	0.005014	0.984133	773	98
		100	0.1	0.055547	0.984133	7296	99
			0.2	0.062751	0.984133	7434	98
			0.3	0.038260	0.984133	7262	98

Again, the results provided from the *custom* implementation, i.e., the SGD with different momentum settings, are strongly similar to those of *sklearn* implementation, i.e., *liblinear* [10] implementation, in terms of $r2$ score. SGD solver always requires even lower iterations, i.e., epochs, for higher C regularization values, i.e., for C equals to 10 or 100, to achieve the same *numerical precision*. *Polyak* and *Nesterov* momentums always perform lower iterations as expected from the theoretical analysis of the convergence rate.


Figure 20: SGD convergence for the Primal formulation of the \mathcal{L}_2 -SVR

Linear Dual formulations The experiments results shown in 21 are obtained with α , i.e., the *learning rate* or *step size*, setted to 1 for the *AdaGrad* algorithm. Note that the *unreg.bias* and *reg.bias* duals refers to the augmented dual formulations (137) of the problems (90) and (91) respectively with ρ equals to 1. The optimization process is stopped if the primal-dual weight vector does not change by at least $1e-8$ between two consecutive iterations.

Table 21: Lagrangian Dual linear \mathcal{L}_2 -SVR results

dual	C	epsilon	fit_time	r2	n_iter	n_sv
reg_bias	1	0.1	29.455724	0.984109	12602	100
		0.2	28.844870	0.984109	12550	100
		0.3	30.448152	0.984109	12510	98
	10	0.1	211.842157	0.984133	100000	98
		0.2	88.650141	0.984133	100000	98
		0.3	87.400583	0.984133	100000	98
	100	0.1	88.560439	0.984131	100000	100
		0.2	84.623098	0.984131	100000	100
		0.3	87.405402	0.984132	100000	100
unreg_bias	1	0.1	29.886016	0.984109	14743	100
		0.2	35.391258	0.984109	14681	100
		0.3	34.577396	0.984109	14691	98
	10	0.1	158.556563	0.984133	100000	98
		0.2	89.270932	0.984133	100000	98
		0.3	89.312740	0.984133	100000	98
	100	0.1	85.335563	0.984131	100000	100
		0.2	86.922623	0.984132	100000	100
		0.3	85.433142	0.984132	100000	100

For what about the linear *Lagrangian dual* formulation we can see as it seems to be insensitive to the increasing complexity of the model in terms of number of *iterations* and require many *iterations* wrt the *Wolfe dual* formulation.


Figure 21: AdaGrad convergence for the Lagrangian Dual formulation of the linear \mathcal{L}_2 -SVR


Nonlinear Dual formulations The experiments results shown in 22 are obtained with d and r hyperparameters both equal to 3 for the *polynomial* kernel; γ is setted to ‘scale’ for both *polynomial* and *gaussian* kernels. The experiments results shown in 22 are obtained with α , i.e., the *learning rate* or *step size*, setted to 1 for the *AdaGrad* algorithm. Note that the *unreg-bias* and *reg-bias* duals refers to the augmented dual formulations (137) of the problems (90) and (91) respectively with ρ equals to 1. The optimization process is stopped if the primal-dual weight vector does not change by at least $1e-8$ between two consecutive iterations.

Table 22: Lagrangian Dual nonlinear \mathcal{L}_2 -SVR results

dual	kernel	C	epsilon	fit_time	r2	n.iter	n_sv
reg-bias	poly	1	0.1	97.551234	0.981708	100000	100
			0.2	99.213337	0.975801	100000	97
			0.3	96.057328	0.948619	100000	96
		10	0.1	96.669691	0.968303	100000	99
			0.2	96.580167	0.967125	100000	99
			0.3	97.984428	0.974149	100000	98
		100	0.1	96.924278	0.965041	100000	100
			0.2	98.486066	0.980911	100000	96
			0.3	95.471666	0.903538	100000	94
	rbf	1	0.1	9.849164	0.980790	8467	38
			0.2	10.063439	0.945653	8563	28
			0.3	10.447014	0.886380	8627	23
		10	0.1	65.058718	0.988575	53056	22
			0.2	45.359936	0.957657	36215	11
			0.3	46.770068	0.900641	37412	8
		100	0.1	122.264825	0.989799	100000	19
			0.2	120.171495	0.966485	100000	10
			0.3	122.602509	0.920652	100000	7
unreg-bias	poly	1	0.1	96.043177	0.981739	100000	99
			0.2	95.672288	0.976645	100000	91
			0.3	96.231995	0.971742	100000	99
		10	0.1	96.942310	0.968902	100000	100
			0.2	97.522340	0.981894	100000	100
			0.3	95.140821	0.800723	100000	95
		100	0.1	96.468312	0.965794	100000	100
			0.2	98.143747	0.977436	100000	96
			0.3	95.971270	0.975918	100000	96
	rbf	1	0.1	19.112700	0.980790	16609	38
			0.2	19.546136	0.945647	16818	28
			0.3	20.166675	0.886373	16970	23
		10	0.1	121.256781	0.988575	99391	22
			0.2	83.407763	0.957660	67123	11
			0.3	87.705072	0.900795	69138	8
		100	0.1	124.393783	0.989783	100000	19
			0.2	119.844592	0.966486	100000	10
			0.3	124.659568	0.920584	100000	7

The same considerations made for the previous linear *Lagrangian dual* formulations are confirmed also in the nonlinearly separable case. In this setting, the complexity of the model coming with higher C regularization hyperparameters and lower ϵ values pays a larger tradeoff in terms of the number of *iterations* of the algorithm.

Figure 22: AdaGrad convergence for the Lagrangian Dual formulation of the Polynomial \mathcal{L}_2 -SVR

Figure 23: AdaGrad convergence for the Lagrangian Dual formulation of the Gaussian \mathcal{L}_2 -SVR

8 Conclusions

For what about the SVM formulations, it is known, in general, that the *primal formulation*, is suitable for large linear training since the complexity of the model grows with the number of features or, more in general, when the number of examples n is much larger than the number of features m , i.e., $n \gg m$; meanwhile the *dual formulation*, is more suitable in case the number of examples n is less than the number of features m , i.e., $n < m$, since the complexity of the model is dominated by the number of examples, or more in general when the training data are not linearly separable in the input space.

From all these experiments we can see as all the *custom* implementations underperforms all the others, i.e., both *cvxopt* [12] and *sklearn* implementations, i.e., *liblinear* [10] and *libsvm* [11] implementations, in terms of *time* obviously due to the different core implementation languages, i.e., Python and C respectively.

In the *primal* formulations the *liblinear* [10] implementation uses an optimization method called *Coordinate Gradient Descent* which minimizes one coordinate at a time.

Meanwhile, for what about the *Wolfe dual* formulations we can notice as *cvxopt* [12] underperforms the *sklearn* implementation, i.e., *libsvm* [11] implementation, in terms of *time* since it is a general-purpose QP solver and it does not exploit the structure of the problem, as SMO does. An interesting consideration can be made about the number of *iterations* of *custom* SMO implementation wrt that in *libsvm* which seems to be always lower thanks to the improvements described in [5, 8] for classification and regression respectively.

Finally, in the *Lagrangian dual* formulations, we can see as fitting the intercept in an explicit way, i.e., by adding Lagrange multipliers to control the equality constraint always get lower scores wrt the *Lagrangian dual* of the same problem with the bias term embedded into the weight matrix.

References

- [1] Boris T Polyak. Some methods of speeding up the convergence of iteration methods. *Ussr computational mathematics and mathematical physics*, 4(5):1–17, 1964.
- [2] Yurii Nesterov. Introductory lectures on convex programming volume i: Basic course. *Lecture notes*, 3(4):5, 1998.
- [3] Yurii E Nesterov. A method for solving the convex programming problem with convergence rate $O(1/k^2)$. In *Dokl. akad. nauk Sssr*, volume 269, pages 543–547, 1983.
- [4] John Platt. Sequential minimal optimization: A fast algorithm for training support vector machines. 1998.
- [5] S. Sathiya Keerthi, Shirish Krishnaji Shevade, Chiranjib Bhattacharyya, and Karuturi Radha Krishna Murthy. Improvements to Platt’s SMO algorithm for SVM classifier design. *Neural computation*, 13(3):637–649, 2001.
- [6] John Duchi, Elad Hazan, and Yoram Singer. Adaptive subgradient methods for online learning and stochastic optimization. *Journal of machine learning research*, 12(7), 2011.
- [7] Gary William Flake and Steve Lawrence. Efficient SVM regression training with SMO. *Machine Learning*, 46(1):271–290, 2002.
- [8] SK Shevade, SS Keerthi, C Bhattacharyya, and KRK Murthy. Improvements to SMO algorithm for SVM regression (Tech. Rep. No. CD-99-16). *Singapore: Control Division Department of Mechanical and Production Engineering*, 1999.
- [9] Tristan Fletcher. Support vector machines explained. *Tutorial paper.*, Mar, page 28, 2009.
- [10] Rong-En Fan, Kai-Wei Chang, Cho-Jui Hsieh, Xiang-Rui Wang, and Chih-Jen Lin. LIBLINEAR: A library for large linear classification. *the Journal of machine Learning research*, 9:1871–1874, 2008.
- [11] Chih-Chung Chang and Chih-Jen Lin. LIBSVM: a library for support vector machines. *ACM transactions on intelligent systems and technology (TIST)*, 2(3):1–27, 2011.
- [12] Lieven Vandenbergh. The CVXOPT linear and quadratic cone program solvers. *Online: <http://cvxopt.org/documentation/coneprog.pdf>*, 2010.
- [13] Veronica Piccialli and Marco Sciandrone. Nonlinear optimization and support vector machines. *4OR*, 16(2):111–149, 2018.
- [14] Chih-Wei Hsu and Chih-Jen Lin. A simple decomposition method for support vector machines. *Machine Learning*, 46(1):291–314, 2002.
- [15] Stephen Boyd, Stephen P Boyd, and Lieven Vandenbergh. *Convex optimization*. Cambridge university press, 2004.
- [16] Xingyu Zhou. On the fenchel duality between strong convexity and lipschitz continuous gradient. *arXiv preprint arXiv:1803.06573*, 2018.