Crossover SNP−by−PCNM interactions grouped by PCNM context

Top crossover examples within each PCNM across bacteria and fungi

Allele A C G T

PCNM12 | PCNM5 | PCNM3 | PCNM17

JHI−Hv50k−2016−274673
Fungi | SpringPan | OTU_739
interaction P = 0.011
slope(C)=−0.14, slope(T)=4.2

JHI−Hv50k−2016−27968
Bacteria | SpringPan | ASV16927
interaction P = 0.012
slope(A)=0.85, slope(G)=−1.5

JHI−Hv50k−2016−201999
Bacteria | SpringPan | ASV137553
interaction P = 0.00054
slope(A)=−0.23, slope(G)=2.6

JHI−Hv50k−2016−27968
Bacteria | SpringPan | ASV68333
interaction P = 0.05
slope(A)=0.86, slope(G)=−0.2

JHI−Hv50k−2016−201999
Fungi | SpringPan | OTU_4625
interaction P = 0.027
slope(A)=3.7, slope(G)=−4.4

JHI−Hv50k−2016−27968
Bacteria | SpringPan | ASV137553
interaction P = 0.015
slope(A)=0.78, slope(G)=−1.3

JHI−Hv50k−2016−201999
Bacteria | SpringPan | ASV16927
interaction P = 0.0012
slope(A)=−0.27, slope(G)=2.6

JHI−Hv50k−2016−201999
Fungi | SpringPan | OTU_3665
interaction P = 0.028
slope(A)=1.7, slope(G)=−2

JHI−Hv50k−2016−27968
Bacteria | SpringPan | ASV68333
interaction P = 0.031
slope(A)=0.63, slope(G)=−0.5

JHI−Hv50k−2016−201999
Fungi | SpringPan | OTU_3662
interaction P = 0.043
slope(A)=1.3, slope(G)=−1.1

rCLR biomarker abunda

PCNM value