

Ranking: Consistency and Rates

Dany Haddad

July 2020

1 Introduction

This is a work in progress document discussing the consistency of common convex surrogate functions for ranking as well as the convergence rate of the corresponding empirical risk. First we summarize the main points of Duchi et al. in this regard [5, 6] focusing on the uniform convergence guarantees. Then we discuss some relevant research directions, in particular, we are interested in achieving fast convergence rates for ranking applications.¹

2 Literature Review

2.1 The Asymptotics of Ranking Algorithms

2.1.1 Overview

The authors first introduce the ranking problem and describe a pairwise loss function, equivalent to the pairwise 0-1 ranking loss. They then define several forms of consistency for ranking problems and unify them under some conditions (theorem 1). Surprisingly, common surrogates for the pairwise 0-1 ranking loss function are shown to not be consistent.

The next section discusses how aggregating the pairwise preferences into lists allows us to define consistent and tractable surrogate loss functions. The key idea is that the surrogates must be order preserving.²

Given that our new surrogate loss functions consist of an aggregation step, it is natural to consider the aggregation in batches. In particular, the authors describe the loss function as a sum of U-statistics. However, this seems mostly to simplify the analysis rather than to leverage the fact that U-statistics

¹I also point out confusion and questions I have in the footnotes.

²The authors describe inconsistency as arising due to a lack of complete preference information. This point is not entirely clear to me since the proof of inconsistency seems to depend on the variability in preference information rather than the lack of information. See Ravikumar et al. [1] for a discussion on how a lack of normalization leads to “non-robustness”.

are UMVUEs.³ The authors then prove a ULLN for the risk functional they define and show that it is consistent for the underlying listwise losses.

I still have not spent sufficient time on the experimental section to provide an overview.

2.2 Preliminaries

The ranking problem can be described generally as ordering a list of m items for each query q in the query space Q (which we assume is countable). This is equivalent to providing a real valued score for each item in the list.

Duchi et al. [6] consider a general setup where the item preference information is received in an arbitrary form and then transformed into a problem specific structure space by an aggregation function. Here we will focus on the case where the preference information Y_i is a matrix of pairwise preferences and the aggregation function s is either a simple average, or a transformation into a list of item scores. We also consider the case where s performs no aggregation at all. Consider the 0-1 pairwise ranking loss:

$$L(\alpha, A) = \sum_{i < j} A_{ij} 1(\alpha_i \leq \alpha_j) + \sum_{i > j} A_{ij} 1(\alpha_i < \alpha_j) \quad (1)$$

where $A = s(\{Y_i\}_{i=1}^k)$. Note that this is different from the typical 0-1 pairwise ranking loss in that the error $\alpha_i = \alpha_j$ is penalized only once.⁴

Assumption A. The distribution of preference information μ_q^k converges weakly to some limiting distribution μ_q for every query q where k is the amount of preference aggregation. So, for $S_k = s(\{Y_i\}_{i=1}^k) \sim \mu_q^k$ as $k \rightarrow \infty$:

$$\mu_q^k \rightarrow^d \mu_q \quad (2)$$

Assumption B. The loss function L is bounded in $[0, 1]$ and continuous.

Given the loss function L we define the loss over a distribution of preference information μ as:

$$\ell(\alpha, \mu) = \int L(\alpha, s) d\mu(s) \quad (3)$$

For the case where the preference information Y_i is a matrix of pairwise preferences and the aggregation function is the average, then this reduces to:

³The authors mention that they introduce the U-statistics based risk functional to be able to analyze the setup without overly detailed knowledge of the surrogate loss. But they already assume the surrogate is Lipschitz continuous and bounded. What more information would we need to conduct the analysis without taking the U-statistics route?

⁴They mention that this is a technical detail, but it's not clear to me where in the proofs it is important. Most likely it has to do with the inconsistency results.

$ell(alpha, \mu) = L(\alpha, Y_{i,j}^\mu)$ where $Y_{i,j}^\mu$ is the average preference of item i over item j according to the distribution μ .

We define the ranking risk of our scoring function $f : Q \rightarrow R^m$ as:

$$R(f) = \sum_q^Q p_q \ell(f(q), \mu_q) \quad (4)$$

Where p_q is the prior probability of query q . Without loss of generality, let p_q be decreasing with the query index q .

Similarly, we define the surrogate loss and surrogate risk as:

$$\ell_\varphi(\alpha, \mu) = \int \varphi(\alpha, s) d\mu(s) \quad (5)$$

$$R_\varphi(f) = \sum_q p_q \ell_\varphi(f(q), \mu_q) \quad (6)$$

2.3 Inconsistency Results

2.4 Proving a ULLN for the U-statistics Based Risk Functional

Define the U-statistics based empirical risk:

$$\hat{R}_{n,\varphi}(f) = \frac{1}{n} \sum_q \hat{n}_q \frac{1}{\binom{\hat{n}_q}{k}} \sum_{I_q \in \mathcal{I}_q} \varphi(f(q), s(\{Y_i\}_{I_q})) \quad (7)$$

where \hat{n}_q is the relative frequency of query q in a sample of size n and \mathcal{I}_q is the set of all combinations of indexes of samples corresponding to query q .

We include conditions I and II as well as assumptions D, E and F from Duchi et al. [6] in the statement of theorem 4.

Theorem 4. For $\phi \in [0, B_n]$ bounded and L_n -lipshitz over \mathcal{F}_n , if there exists some $\rho, \beta > 0$ such that:

$$p_q = O(q^{-\beta-1}) \quad (8)$$

and the expected surrogate loss at aggregation level k satisfies for all f, k, q, n :

$$\left| \mathbb{E}_q \varphi(f(q), s(\{Y_i\}^k)) - \lim_{k' \rightarrow \infty} \mathbb{E}_q \varphi(f(q), s(\{Y_i\}^{k'})) \right| = O(B_n k^{-\rho}) \quad (9)$$

with $B_n/k_n^\rho = o(1)$ and $k_n B_n^{(1+\beta)/\beta} = o(n)$ and in addition we have the following constrain on the covering numbers:

$$k_n B_n \sqrt{\log \left(N \left(\frac{\epsilon}{4L_n} \right) \right)} = o(\sqrt{n}) \quad (10)$$

then we have the uniform convergence guarantee:

$$\sup_{f \in \mathcal{F}_n} \left| \hat{R}_{n,\varphi}(f) - R_\varphi(f) \right| \rightarrow^p 0 \quad (11)$$

The proof for this uniform law of large numbers can be broken down by decomposing the uniform deviation into three terms and controlling each of them:

$$\sup_{f \in \mathcal{F}_n} \left| \hat{R}_{n,\varphi}(f) - R_\varphi(f) \right| \leq \sup_{f \in \mathcal{F}_n} \left| \hat{R}_{n,\varphi}(f) - R_{n,\varphi}(f) \right| + \sup_{f \in \mathcal{F}_n} |R_{n,\varphi}(f) - R_\varphi(f)| \quad (12)$$

where we denote :

$$R_{\varphi,n} = \sum_q \sum_{l=0}^n l \mathbb{P}_n(\hat{n}_q = l) \mathbb{E}(\varphi(f(q), s(\{Y_i\}^{\min(l,k)})) | Q = q) \quad (13)$$

The first term (I) on the RHS of inequality 12 is the deviation of the U-statistic based estimate $\hat{R}_{n,\varphi}(f)$ from it's expectation $R_{n,\varphi}(f)$. The second term (II) is the deviation of $R_{n,\varphi}(f)$ from it's limit $R_\varphi(f)$ as $n, k \rightarrow \infty$. The second term is controlled by lemma 8 of the supplementary material [6] while the first term is controlled by lemma 10.

Lemma 8 uses the conditions on the the boundedness of ϕ and the aggregation level k_n as well as a covering number argument to give a deterministic bound on the second term above.

Lemma 10 shows that the first term is $o_p(1)$ by arguing that the risk functional $\hat{R}_{n,\varphi}(f)$ satisfies a bounded differences inequality. Applying McDiarmid's inequality as well as a union bound involving the covering number gives the following bound, yielding the claim:

$$\mathbb{P} \left(\sup_{f \in \mathcal{F}_n} \left| \hat{R}_{n,\varphi}(f) - R_{n,\varphi}(f) \right| \geq \epsilon \right) \leq 2 \exp \left(\log N \left(\frac{\epsilon}{4L_n}, n \right) - \frac{n\epsilon^2}{32k^2 B_n^2} \right) \quad (14)$$

where $N \left(\frac{\epsilon}{4L_n}, n \right)$ gives the covering number of \mathcal{F}_n .⁵

⁵My understanding is that this gives a rate worse than $O_p(1/\sqrt{n})$ since the concentration inequality involves the covering number which is dependent on ϵ . Unless for the function class under consideration this can be bound uniformly independent of *epsilon*, this is worse than the typical rate.

2.4.1 Potential improvements

The U-statistics formulation is totally ignored in controlling term (I), leading to a rate slower than $O_p(1/\sqrt{n})$. For the case of bipartite ranking, Clemencon et al. [4] derive fast rates on the convergence of the empirical risk minimizer to the optimal optimizer. The approach involves making a low noise assumption (bound variance of the risk in terms of the expectation) and applying a Hoeffding decomposition of the U-statistic based risk. The decomposed terms are controlled by applying the decoupling technique for degenerate U-statistics.

However, their results are more relevant in a model selection context since they deal directly with the non-convex pairwise 0-1 ranking loss. Clemencon et al. [4] also mention (in Remark 7) that it would be interesting to derive fast rates for convex surrogates, but that it would be quite technical. They point to Blanchard et al. [3] for a similar derivation for the binary classification case.

Bartlett et al. [2] derive fast rates for convex surrogates in the context of binary classification. They also demonstrate via simulation that their derived rates are representative. Clemencon et al. [4] show how the results from this work transfer over to the case of bipartite ranking but do not derive fast rates.

Questions

- What is the next step in deriving fast rates for the case of bipartite ranking?
- Is the case of listwise ranking with aggregation (as framed by Duchi) simpler?
- How can we unify the notions of aggregation from Duchi and normalization from Ravikumar?
- Why hasn't there been further work on this topic in the past 10 years?
- Is the model selection problem more interesting? In particular for the case of listwise ranking.

References

- [1] On NDCG Consistency of Listwise Ranking Methods.
- [2] Peter L Bartlett, Michael I Jordan, and Jon D McAuliffe. Convexity, Classification, and Risk Bounds. *Journal of the American Statistical Association*, 101(473):138–156, 2006.
- [3] Gilles Blanchard, Gabor Lugosi, and Nicola Vayatis. On the rates of convergence of regularized boosting classifiers.

- [4] Stéphan Cléménçon, Gábor Lugosi, and Nicolas Vayatis. Ranking and Empirical Minimization of U-statistics. *The Annals of Statistics*, 36(2):844–874, 2008.
- [5] John Duchi. On the Consistency of Ranking Algorithms.
- [6] John C. Duchi, Lester Mackey, and Michael I. Jordan. The asymptotics of ranking algorithms. *The Annals of Statistics*, 41(5):2292–2323, 2013.