

ROB 101 - Fall 2021

# Triangular Systems of Equations: Forward and Back Substitution

September 8, 2021



- ▶ Equations that have special structure are often much easier to solve
- ▶ Some examples to show this.

- ▶ Recognize triangular systems of linear equations and distinguish those with a unique answer.
- ▶ Learn that the determinant of a square triangular matrix is the product of the terms on its diagonal.
- ▶ How to use forward and back substitution.

## Example of Lower Triangular (Linear) Systems

This is an example of a square system of linear equations that is *lower triangular*.

$$3x_1 = 6$$

$$2x_1 - x_2 = -2$$

$$x_1 - 2x_2 + 3x_3 = 2.$$

## Example of Lower Triangular (Linear) Systems

When we write the system as  $Ax = b$ , in the lower triangular case we have

$$\begin{array}{rcl} 3x_1 & = & 6 \\ 2x_1 - x_2 & = & -2 \\ x_1 - 2x_2 + 3x_3 & = & 2 \end{array} \iff \underbrace{\begin{bmatrix} 3 & 0 & 0 \\ 2 & -1 & 0 \\ 1 & -2 & 3 \end{bmatrix}}_A \underbrace{\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}}_x = \underbrace{\begin{bmatrix} 6 \\ -2 \\ 2 \end{bmatrix}}_b.$$

$$A = \begin{bmatrix} 3 & 0 & 0 \\ 2 & -1 & 0 \\ 1 & -2 & 3 \end{bmatrix}$$

- ▶ All terms above the diagonal of the matrix  $A$  are zero.
- ▶ More precisely, the condition is  $a_{ij} = 0$  for all  $j > i$ .
- ▶ Such matrices are called *lower-triangular*.

## Determinant of a Lower Triangular Matrix

### Fact

*The matrix determinant of a square lower triangular matrix is equal to the product of the elements on the diagonal.*

$$A = \begin{bmatrix} 3 & 0 & 0 \\ 2 & -1 & 0 \\ 1 & -2 & 3 \end{bmatrix} \implies \det(A) = 3 \cdot (-1) \cdot 3 = -9 \neq 0.$$

# Determinant of a Lower Triangular Matrix

## Fact

*The matrix determinant of a square lower triangular matrix is equal to the product of the elements on the diagonal.*

$$A = \begin{bmatrix} 3 & 0 & 0 \\ 2 & -1 & 0 \\ 1 & -2 & 3 \end{bmatrix} \implies \det(A) = 3 \cdot (-1) \cdot 3 = -9 \neq 0.$$

```
In [1]: using LinearAlgebra  
  
A = [3 0 0; 2 -1 0; 1 -2 3];  
det(A)
```

```
Out[1]: -9.0
```



## Lower Triangular Systems and Forward Substitution

We will solve the previous example using a method called *forward substitution*.

$$\begin{array}{rcl} 3x_1 & = & 6 \\ 2x_1 - x_2 & = & -2 \\ x_1 - 2x_2 + 3x_3 & = & 2 \end{array} \iff \underbrace{\begin{bmatrix} 3 & 0 & 0 \\ 2 & -1 & 0 \\ 1 & -2 & 3 \end{bmatrix}}_A \underbrace{\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}}_x = \underbrace{\begin{bmatrix} 6 \\ -2 \\ 2 \end{bmatrix}}_b.$$

## Lower Triangular Systems and Forward Substitution

Because we have ordered the variables as  $x_1, x_2, x_3$ , we isolate  $x_1$  in the first equation,  $x_2$  in the second equation, and  $x_3$  in the third equation by moving the other variables to the right-hand side,

$$3x_1 = 6$$

$$-x_2 = -2 - 2x_1$$

$$3x_3 = 2 - x_1 + 2x_2.$$

## Lower Triangular Systems and Forward Substitution

Next, we substitute in, working from top to bottom:

$$x_1 = \frac{1}{3}6 = 2$$

$$x_2 = 2 + 2x_1 = 6$$

$$x_3 = \frac{1}{3}(2 - x_1 + 2x_2) = \frac{1}{3}(12) = 4.$$

## Upper Triangular Systems

This is an example of a square *upper triangular* system of equations.

$$\begin{array}{rcl} x_1 + 3x_2 + 2x_3 & = & 6 \\ 2x_2 + x_3 & = & -2 \\ 3x_3 & = & 4, \end{array} \iff \underbrace{\begin{bmatrix} 1 & 3 & 2 \\ 0 & 2 & 1 \\ 0 & 0 & 3 \end{bmatrix}}_A \underbrace{\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}}_x = \underbrace{\begin{bmatrix} 6 \\ -2 \\ 4 \end{bmatrix}}_b.$$

$$A = \begin{bmatrix} 1 & 3 & 2 \\ \mathbf{0} & 2 & 1 \\ \mathbf{0} & \mathbf{0} & 3 \end{bmatrix}$$

- ▶ All terms below the diagonal of the matrix  $A$  are zero.
- ▶ More precisely, the condition is  $a_{ij} = 0$  for  $i > j$ .
- ▶ Such matrices are called *upper-triangular*.

# Determinant of an Upper Triangular Matrix

## Fact

*The matrix determinant of a square upper triangular matrix is equal to the product of the elements on the diagonal.*

$$A = \begin{bmatrix} 1 & 3 & 2 \\ 0 & 2 & 1 \\ 0 & 0 & 3 \end{bmatrix} \implies \det(A) = 1 \cdot 2 \cdot 3 = 6 \neq 0.$$

# Determinant of an Upper Triangular Matrix

## Fact

*The matrix determinant of a square upper triangular matrix is equal to the product of the elements on the diagonal.*

$$A = \begin{bmatrix} 1 & 3 & 2 \\ 0 & 2 & 1 \\ 0 & 0 & 3 \end{bmatrix} \implies \det(A) = 1 \cdot 2 \cdot 3 = 6 \neq 0.$$

```
In [2]: A = [1 3 2; 0 2 1; 0 0 3];  
        det(A)
```

```
Out[2]: 6.0
```

We solve the upper triangular systems using a method called *back substitution*.

$$\begin{array}{rcl} x_1 + 3x_2 + 2x_3 & = & 6 \\ 2x_2 + x_3 & = & -2 \\ 3x_3 & = & 4, \end{array} \iff \underbrace{\begin{bmatrix} 1 & 3 & 2 \\ 0 & 2 & 1 \\ 0 & 0 & 3 \end{bmatrix}}_A \underbrace{\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}}_x = \underbrace{\begin{bmatrix} 6 \\ -2 \\ 4 \end{bmatrix}}_b.$$



We develop a solution by starting at the bottom and work our way to the top via back substitution.

$$x_1 = 6 - (3x_2 + 2x_3)$$

$$x_2 = \frac{1}{2}(-2 - x_3)$$

$$x_3 = \frac{4}{3},$$

Next, we do back substitution, from bottom to top, sequentially plugging in numbers from the previous equations.

$$x_1 = 6 - (3x_2 + 2x_3) = 6 - \left(3 \cdot \left(-\frac{5}{3}\right) + 2 \cdot \frac{4}{3}\right) = \frac{11}{3}$$

$$x_2 = \frac{1}{2} \cdot (-2 - x_3) = \frac{1}{2} \cdot \left(-2 - \frac{4}{3}\right) = -\frac{5}{3}$$

$$x_3 = \frac{4}{3}.$$

## General Cases of Triangular Systems

The general form of a lower triangular system with a non-zero determinant is

$$\begin{aligned}a_{11}x_1 &= b_1 & (a_{11} \neq 0) \\a_{21}x_1 + a_{22}x_2 &= b_2 & (a_{22} \neq 0) \\&\vdots = \vdots \\a_{n1}x_1 + a_{n2}x_2 + a_{n3}x_3 + \cdots + a_{nn}x_n &= b_n & (a_{nn} \neq 0)\end{aligned}$$

## General Cases of Triangular Systems

and the solution proceeds from top to bottom, like this

$$x_1 = \frac{b_1}{a_{11}} \quad (a_{11} \neq 0)$$

$$x_2 = \frac{b_2 - a_{21}x_1}{a_{22}} \quad (a_{22} \neq 0)$$

$$\vdots = \vdots$$

$$x_n = \frac{b_n - a_{n1}x_1 - a_{n2}x_2 - \cdots - a_{n,n-1}x_{n-1}}{a_{nn}} \quad (a_{nn} \neq 0).$$

## General Cases of Triangular Systems

The general form of an upper triangular system with a non-zero determinant is

$$a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \cdots + a_{1n}x_n = b_1 \quad (a_{11} \neq 0)$$

$$a_{22}x_2 + a_{23}x_3 + \cdots + a_{2n}x_n = b_2 \quad (a_{22} \neq 0)$$

$$a_{33}x_3 + \cdots + a_{3n}x_n = b_3 \quad (a_{33} \neq 0)$$

$$\vdots = \vdots$$

$$a_{nn}x_n = b_n \quad (a_{nn} \neq 0),$$

## General Cases of Triangular Systems

and the solution proceeds from bottom to top, like this,

$$x_1 = \frac{b_1 - a_{12}x_2 - \cdots - a_{1n}x_n}{a_{11}} \quad (a_{11} \neq 0)$$

$$x_2 = \frac{b_2 - a_{23}x_3 - \cdots - a_{2n}x_n}{a_{22}} \quad (a_{22} \neq 0)$$

$$\vdots = \vdots \quad \quad \quad \vdots$$

$$x_{n-1} = \frac{b_{n-1} - a_{n-1,n}x_n}{a_{n-1,n-1}} \quad (a_{n-1,n-1} \neq 0)$$

$$x_n = \frac{b_n}{a_{nn}} \quad (a_{nn} \neq 0),$$

In HW you will develop Julia code to solve triangular systems of equations, whether upper or lower triangular!

- ▶ Matrix Multiplication
- ▶ Read Chapter 4 of ROB 101 Book