

ROB 101 - Fall 2021

Matrix Multiplication

September 13, 2021



- ▶ How to partition matrices into rows and columns
- ▶ How to multiply one matrix times another
- ▶ How to swap rows of a matrix

- ▶ Multiplying a row vector by a column vector
- ▶ Recognizing the rows and columns of a matrix
- ▶ Standard definition of matrix multiplication $A \cdot B$ using the rows of A and the columns of B
- ▶ Size restrictions when multiplying matrices
- ▶ Examples that work and those that don't because the sizes are wrong
- ▶ Permutation matrices

Multiplying a Row Vector by a Column Vector

Let $a^{\text{row}} = [a_1 \ a_2 \ \cdots \ a_k]$ be a row vector with k elements

and let $b^{\text{col}} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_k \end{bmatrix}$ be a column vector with the same number
of elements as a^{row} .

Multiplying a Row Vector by a Column Vector

Definition

The *product* of a^{row} and b^{col} is defined as

$$a^{\text{row}} \cdot b^{\text{col}} := \sum_{i=1}^k a_i b_i.$$

Multiplying a Row Vector by a Column Vector

Definition

The *product* of a^{row} and b^{col} is defined as

$$a^{\text{row}} \cdot b^{\text{col}} := \sum_{i=1}^k a_i b_i.$$

For many, the following visual representation is more understandable,

$$\begin{bmatrix} a_1 & a_2 & \cdots & a_k \end{bmatrix} \cdot \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_k \end{bmatrix} := a_1 b_1 + a_2 b_2 + \cdots + a_k b_k.$$

Let $a^{\text{row}} = [2 \quad -3 \quad -1 \quad 11]$ be a row vector with $k = 4$ elements and let $b^{\text{col}} = \begin{bmatrix} 3 \\ 5 \\ -1 \\ -2 \end{bmatrix}$ be a column vector with $k = 4$ elements. Perform their multiplication if it makes sense.

Because they have the same number of elements, we can form their product and we compute

$$\begin{aligned} a^{\text{row}} \cdot b^{\text{col}} &:= \sum_{i=1}^4 a_i b_i \\ &= (2)(3) + (-3)(5) + (-1)(-1) + (11)(-2) = -30, \end{aligned}$$

Because they have the same number of elements, we can form their product and we compute

$$\begin{aligned} a^{\text{row}} \cdot b^{\text{col}} &:= \sum_{i=1}^4 a_i b_i \\ &= (2)(3) + (-3)(5) + (-1)(-1) + (11)(-2) = -30, \end{aligned}$$

or, equivalently, we write it like this

$$\begin{aligned} [2 \quad -3 \quad -1 \quad 11] \cdot \begin{bmatrix} 3 \\ 5 \\ -1 \\ -2 \end{bmatrix} \\ = (2)(3) + (-3)(5) + (-1)(-1) + (11)(-2) = -30. \end{aligned}$$

Because they have the same number of elements, we can form their product and we compute

$$\begin{aligned} a^{\text{row}} \cdot b^{\text{col}} &:= \sum_{i=1}^4 a_i b_i \\ &= (2)(3) + (-3)(5) + (-1)(-1) + (11)(-2) = -30, \end{aligned}$$

```
In [1]: a = [2. -3 -1 11];  
        b = [3., 5, -1, -2];  
        a * b
```

```
Out[1]: 1-element Array{Float64,1}:  
        -30.0
```

Examples of Row and Column Partitions

Let $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}$ be a 2×3 matrix.

Examples of Row and Column Partitions

Let $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}$ be a 2×3 matrix.

Then a *partition* of A into *rows* is

$$\begin{bmatrix} a_1^{\text{row}} \\ a_2^{\text{row}} \end{bmatrix} = \begin{bmatrix} \boxed{1 \ 2 \ 3} \\ \boxed{4 \ 5 \ 6} \end{bmatrix}, \quad \text{that is,} \quad \begin{aligned} a_1^{\text{row}} &= [1 \ 2 \ 3] \\ a_2^{\text{row}} &= [4 \ 5 \ 6]. \end{aligned}$$

Examples of Row and Column Partitions

Let $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}$ be a 2×3 matrix.

Then a *partition* of A into *rows* is

$$\begin{bmatrix} a_1^{\text{row}} \\ a_2^{\text{row}} \end{bmatrix} = \begin{bmatrix} \boxed{1 \ 2 \ 3} \\ \boxed{4 \ 5 \ 6} \end{bmatrix}, \quad \text{that is,} \quad \begin{aligned} a_1^{\text{row}} &= [1 \ 2 \ 3] \\ a_2^{\text{row}} &= [4 \ 5 \ 6]. \end{aligned}$$

Remark

We note that a_1^{row} and a_2^{row} are row vectors of size 1×3 ; they have the same number of entries as A has columns.

Examples of Row and Column Partitions

A *partition* of $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}$ into *columns* is

$$\begin{bmatrix} a_1^{\text{col}} & a_2^{\text{col}} & a_3^{\text{col}} \end{bmatrix} = \left[\begin{array}{|c|} \hline 1 \\ \hline 4 \\ \hline \end{array} \quad \begin{array}{|c|} \hline 2 \\ \hline 5 \\ \hline \end{array} \quad \begin{array}{|c|} \hline 3 \\ \hline 6 \\ \hline \end{array} \right],$$

Examples of Row and Column Partitions

A *partition* of $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}$ into *columns* is

$$\begin{bmatrix} a_1^{\text{col}} & a_2^{\text{col}} & a_3^{\text{col}} \end{bmatrix} = \left[\begin{array}{|c|} \hline 1 \\ \hline 4 \\ \hline \end{array} \quad \begin{array}{|c|} \hline 2 \\ \hline 5 \\ \hline \end{array} \quad \begin{array}{|c|} \hline 3 \\ \hline 6 \\ \hline \end{array} \right],$$

that is,

$$a_1^{\text{col}} = \begin{bmatrix} 1 \\ 4 \end{bmatrix}, \quad a_2^{\text{col}} = \begin{bmatrix} 2 \\ 5 \end{bmatrix}, \quad a_3^{\text{col}} = \begin{bmatrix} 3 \\ 6 \end{bmatrix}$$

Examples of Row and Column Partitions

A *partition* of $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}$ into *columns* is

$$\begin{bmatrix} a_1^{\text{col}} & a_2^{\text{col}} & a_3^{\text{col}} \end{bmatrix} = \left[\boxed{\begin{bmatrix} 1 \\ 4 \end{bmatrix}} \quad \boxed{\begin{bmatrix} 2 \\ 5 \end{bmatrix}} \quad \boxed{\begin{bmatrix} 3 \\ 6 \end{bmatrix}} \right],$$

that is,

$$a_1^{\text{col}} = \begin{bmatrix} 1 \\ 4 \end{bmatrix}, \quad a_2^{\text{col}} = \begin{bmatrix} 2 \\ 5 \end{bmatrix}, \quad a_3^{\text{col}} = \begin{bmatrix} 3 \\ 6 \end{bmatrix}$$

Remark

We note that a_1^{col} , a_2^{col} , and a_3^{col} are column vectors of size 2×1 ; they have the same number of entries as A has rows.

Let A be an $n \times m$ matrix. A partition of A into rows is

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1m} \\ a_{21} & a_{22} & \cdots & a_{2m} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nm} \end{bmatrix} = \begin{bmatrix} a_1^{\text{row}} \\ a_2^{\text{row}} \\ \vdots \\ a_n^{\text{row}} \end{bmatrix} = \begin{bmatrix} \boxed{a_{11} \ a_{12} \ \cdots \ a_{1m}} \\ \boxed{a_{21} \ a_{22} \ \cdots \ a_{2m}} \\ \vdots \\ \boxed{a_{n1} \ a_{n2} \ \cdots \ a_{nm}} \end{bmatrix}.$$

Let A be an $n \times m$ matrix. A partition of A into rows is

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1m} \\ a_{21} & a_{22} & \cdots & a_{2m} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nm} \end{bmatrix} = \begin{bmatrix} a_1^{\text{row}} \\ a_2^{\text{row}} \\ \vdots \\ a_n^{\text{row}} \end{bmatrix} = \begin{bmatrix} \boxed{a_{11} \ a_{12} \ \cdots \ a_{1m}} \\ \boxed{a_{21} \ a_{22} \ \cdots \ a_{2m}} \\ \vdots \\ \boxed{a_{n1} \ a_{n2} \ \cdots \ a_{nm}} \end{bmatrix}.$$

That is, the i -th row is the $1 \times m$ row vector

$$a_i^{\text{row}} = [a_{i1} \ a_{i2} \ \cdots \ a_{im}],$$

where i varies from 1 to n .

A partition of A into columns is

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1m} \\ a_{21} & a_{22} & \cdots & a_{2m} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nm} \end{bmatrix} = \begin{bmatrix} a_1^{\text{col}} & a_2^{\text{col}} & \cdots & a_m^{\text{col}} \end{bmatrix}$$

$$= \left[\begin{array}{c} a_{11} \\ a_{21} \\ \vdots \\ a_{n1} \end{array} \right] \begin{array}{c} a_{12} \\ a_{22} \\ \vdots \\ a_{n2} \end{array} \cdots \begin{array}{c} a_{1m} \\ a_{2m} \\ \vdots \\ a_{nm} \end{array} \right].$$

General Case of Partitions

A partition of A into columns is

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1m} \\ a_{21} & a_{22} & \cdots & a_{2m} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nm} \end{bmatrix} = \left[\begin{array}{c} a_{11} \\ a_{21} \\ \vdots \\ a_{n1} \end{array} \right] \left[\begin{array}{c} a_{12} \\ a_{22} \\ \vdots \\ a_{n2} \end{array} \right] \cdots \left[\begin{array}{c} a_{1m} \\ a_{2m} \\ \vdots \\ a_{nm} \end{array} \right].$$

That is, the j -th column is the $n \times 1$ column vector

$$a_j^{\text{col}} = \begin{bmatrix} a_{1j} \\ a_{2j} \\ \vdots \\ a_{nj} \end{bmatrix}, \text{ where } j \text{ varies from } 1 \text{ to } m.$$

- ▶ Let A be an $n \times k$ matrix, meaning it has n rows and k columns.

Standard Matrix Multiplication

- ▶ Let A be an $n \times k$ matrix, meaning it has n rows and k columns.
- ▶ Let B be a $k \times m$ matrix, so that it has k rows and m columns.

Standard Matrix Multiplication

- ▶ Let A be an $n \times k$ matrix, meaning it has n rows and k columns.
- ▶ Let B be a $k \times m$ matrix, so that it has k rows and m columns.
- ▶ The values of n , m , and k can be any integers greater than or equal to one.

Definition

When the number of columns of the first matrix A equals the number of rows of the second matrix B , the *matrix product* of A and B is defined and results in an $n \times m$ matrix:

$$[n \times k \text{ matrix}] \cdot [k \times m \text{ matrix}] = [n \times m \text{ matrix}].$$

Definition

When the number of columns of the first matrix A equals the number of rows of the second matrix B , the *matrix product* of A and B is defined and results in an $n \times m$ matrix:

$$[n \times k \text{ matrix}] \cdot [k \times m \text{ matrix}] = [n \times m \text{ matrix}].$$

- ▶ The standard way of doing matrix multiplication $A \cdot B$ involves multiplying the rows of A with the columns of B .

Definition

When the number of columns of the first matrix A equals the number of rows of the second matrix B , the *matrix product* of A and B is defined and results in an $n \times m$ matrix:

$$[n \times k \text{ matrix}] \cdot [k \times m \text{ matrix}] = [n \times m \text{ matrix}].$$

- ▶ The standard way of doing matrix multiplication $A \cdot B$ involves multiplying the rows of A with the columns of B .
- ▶ In general, $A \cdot B \neq B \cdot A$ even when A and B are square matrices of the same size.

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} a_1^{\text{row}} \\ a_2^{\text{row}} \end{bmatrix} = \begin{bmatrix} \boxed{1 \ 2} \\ \boxed{3 \ 4} \end{bmatrix}$$

and

$$B = \begin{bmatrix} 5 \\ 6 \end{bmatrix} = \begin{bmatrix} b_1^{\text{col}} \end{bmatrix} = \begin{bmatrix} \boxed{5} \\ \boxed{6} \end{bmatrix}.$$

The matrix product of A and B is

$$A \cdot B = \begin{bmatrix} a_1^{\text{row}} \\ a_2^{\text{row}} \end{bmatrix} \cdot \begin{bmatrix} b_1^{\text{col}} \end{bmatrix} := \begin{bmatrix} a_1^{\text{row}} \cdot b_1^{\text{col}} \\ a_2^{\text{row}} \cdot b_1^{\text{col}} \end{bmatrix} = \begin{bmatrix} 17 \\ 39 \end{bmatrix}$$

The matrix product of A and B is

$$A \cdot B = \begin{bmatrix} a_1^{\text{row}} \\ a_2^{\text{row}} \end{bmatrix} \cdot \begin{bmatrix} b_1^{\text{col}} \end{bmatrix} := \begin{bmatrix} a_1^{\text{row}} \cdot b_1^{\text{col}} \\ a_2^{\text{row}} \cdot b_1^{\text{col}} \end{bmatrix} = \begin{bmatrix} 17 \\ 39 \end{bmatrix}$$

because

$$a_1^{\text{row}} \cdot b_1^{\text{col}} = \begin{bmatrix} 1 & 2 \end{bmatrix} \cdot \begin{bmatrix} 5 \\ 6 \end{bmatrix} = 5 + 12 = 17$$

$$a_2^{\text{row}} \cdot b_1^{\text{col}} = \begin{bmatrix} 3 & 4 \end{bmatrix} \cdot \begin{bmatrix} 5 \\ 6 \end{bmatrix} = 15 + 24 = 39.$$

The matrix product of A and B is

$$A \cdot B = \begin{bmatrix} a_1^{\text{row}} \\ a_2^{\text{row}} \end{bmatrix} \cdot \begin{bmatrix} b_1^{\text{col}} \end{bmatrix} := \begin{bmatrix} a_1^{\text{row}} \cdot b_1^{\text{col}} \\ a_2^{\text{row}} \cdot b_1^{\text{col}} \end{bmatrix} = \begin{bmatrix} 17 \\ 39 \end{bmatrix}$$

```
In [2]: A = [1 2; 3 4];  
        B = [5; 6];  
        A * B
```

```
Out[2]: 2-element Array{Int64,1}:  
        17  
        39
```

We reuse A and B above and ask if we can form the matrix product in the order $B \cdot A$.

We reuse A and B above and ask if we can form the matrix product in the order $B \cdot A$.

We have that the first matrix B is 2×1 and the second matrix A is 2×2 . The number of columns of the first matrix does not match the number of rows of the second matrix, and hence the product cannot be defined in this direction.

We reuse A and B above and ask if we can form the matrix product in the order $B \cdot A$.

We have that the first matrix B is 2×1 and the second matrix A is 2×2 . The number of columns of the first matrix does not match the number of rows of the second matrix, and hence the product cannot be defined in this direction.

```
In [3]: # Don't try this at home!  
A = [1 2; 3 4];  
B = [5; 6];  
B * A
```

```
DimensionMismatch("matrix A has dimensions (2,1), matrix B has dimensions (2,2)")
```

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} a_1^{\text{row}} \\ a_2^{\text{row}} \end{bmatrix} = \begin{bmatrix} \boxed{\begin{matrix} 1 & 2 \end{matrix}} \\ \boxed{\begin{matrix} 3 & 4 \end{matrix}} \end{bmatrix}$$

and

$$B = \begin{bmatrix} 5 & -2 \\ 6 & 1 \end{bmatrix} = \begin{bmatrix} b_1^{\text{col}} & b_2^{\text{col}} \end{bmatrix} = \begin{bmatrix} \boxed{\begin{matrix} 5 \\ 6 \end{matrix}} & \boxed{\begin{matrix} -2 \\ 1 \end{matrix}} \end{bmatrix}.$$

The matrix product of A and B is

$$\begin{aligned} A \cdot B &= \left[\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 4 \\ \hline \end{array} \right] \cdot \left[\begin{array}{|c|c|} \hline 5 & -2 \\ \hline 6 & 1 \\ \hline \end{array} \right] \\ &= \left[\begin{array}{cc} (1)(5) + (2)(6) & (1)(-2) + (2)(1) \\ (3)(5) + (4)(6) & (3)(-2) + (4)(1) \end{array} \right] = \left[\begin{array}{cc} 17 & 0 \\ 39 & -2 \end{array} \right]. \end{aligned}$$

The matrix product of B and A is

$$\begin{aligned} B \cdot A &= \left[\begin{array}{|c|c|} \hline 5 & -2 \\ \hline 6 & 1 \\ \hline \end{array} \right] \cdot \left[\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 4 \\ \hline \end{array} \right] \\ &= \left[\begin{array}{cc} (5)(1) + (-2)(3) & (5)(2) + (-2)(4) \\ (6)(1) + (1)(3) & (6)(2) + (1)(4) \end{array} \right] = \left[\begin{array}{cc} -1 & 2 \\ 9 & 16 \end{array} \right]. \end{aligned}$$

Example 3

```
In [4]: A = [1 2; 3 4];  
        B = [5 -2; 6 1];  
        A * B
```

```
Out[4]: 2x2 Array{Int64,2}:  
        17  0  
        39 -2
```

```
In [5]: B * A
```

```
Out[5]: 2x2 Array{Int64,2}:  
        -1  2  
         9 16
```

General Case: What is happening inside Julia

We partition the $n \times k$ matrix A into rows and the $k \times m$ matrix B into columns, as in

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1k} \\ a_{21} & a_{22} & \cdots & a_{2k} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nk} \end{bmatrix} = \begin{bmatrix} a_1^{\text{row}} \\ a_2^{\text{row}} \\ \vdots \\ a_n^{\text{row}} \end{bmatrix}$$
$$= \begin{bmatrix} \boxed{a_{11} \quad a_{12} \quad \cdots \quad a_{1k}} \\ \boxed{a_{21} \quad a_{22} \quad \cdots \quad a_{2k}} \\ \vdots \\ \boxed{a_{n1} \quad a_{n2} \quad \cdots \quad a_{nk}} \end{bmatrix}$$

General Case: What is happening inside Julia

and

$$B = \begin{bmatrix} b_{11} & b_{12} & \cdots & b_{1m} \\ b_{21} & b_{22} & \cdots & b_{2m} \\ \vdots & \vdots & \ddots & \vdots \\ b_{k1} & b_{k2} & \cdots & b_{km} \end{bmatrix} = \begin{bmatrix} b_1^{\text{col}} & b_2^{\text{col}} & \cdots & b_m^{\text{col}} \end{bmatrix}$$
$$= \left[\begin{array}{c} b_{11} \\ b_{21} \\ \vdots \\ b_{k1} \end{array} \quad \begin{array}{c} b_{12} \\ b_{22} \\ \vdots \\ b_{k2} \end{array} \quad \cdots \quad \begin{array}{c} b_{1m} \\ b_{2m} \\ \vdots \\ b_{km} \end{array} \right],$$

General Case: What is happening inside Julia

then

$$A \cdot B := \begin{bmatrix} a_1^{\text{row}} \cdot b_1^{\text{col}} & a_1^{\text{row}} \cdot b_2^{\text{col}} & \dots & a_1^{\text{row}} \cdot b_m^{\text{col}} \\ a_2^{\text{row}} \cdot b_1^{\text{col}} & a_2^{\text{row}} \cdot b_2^{\text{col}} & \dots & a_2^{\text{row}} \cdot b_m^{\text{col}} \\ \vdots & \vdots & \ddots & \vdots \\ a_n^{\text{row}} \cdot b_1^{\text{col}} & a_n^{\text{row}} \cdot b_2^{\text{col}} & \dots & a_n^{\text{row}} \cdot b_m^{\text{col}} \end{bmatrix}.$$

General Case: What is happening inside Julia

Another way to see the pattern is like this

$$\begin{aligned}
 A \cdot B &:= \left[\begin{array}{|c|} \hline \begin{array}{|c|} \hline a_{11} & a_{12} & \cdots & a_{1k} \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline a_{21} & a_{22} & \cdots & a_{2k} \\ \hline \end{array} \\ \hline \vdots \\ \hline \begin{array}{|c|} \hline a_{n1} & a_{n2} & \cdots & a_{nk} \\ \hline \end{array} \\ \hline \end{array} \right] \cdot \left[\begin{array}{|c|} \hline \begin{array}{|c|} \hline b_{11} \\ b_{21} \\ \vdots \\ b_{k1} \\ \hline \end{array} \\ \hline \begin{array}{|c|} \hline b_{12} \\ b_{22} \\ \vdots \\ b_{k2} \\ \hline \end{array} \\ \hline \cdots \\ \hline \begin{array}{|c|} \hline b_{1m} \\ b_{2m} \\ \vdots \\ b_{km} \\ \hline \end{array} \\ \hline \end{array} \right] \\
 &= \left[\begin{array}{|c|} \hline \sum_{i=1}^k a_{1i}b_{i1} & \sum_{i=1}^k a_{1i}b_{i2} & \cdots & \sum_{i=1}^k a_{1i}b_{im} \\ \hline \sum_{i=1}^k a_{2i}b_{i1} & \sum_{i=1}^k a_{2i}b_{i2} & \cdots & \sum_{i=1}^k a_{2i}b_{im} \\ \hline \vdots & \vdots & \ddots & \vdots \\ \hline \sum_{i=1}^k a_{ni}b_{i1} & \sum_{i=1}^k a_{ni}b_{i2} & \cdots & \sum_{i=1}^k a_{ni}b_{im} \\ \hline \end{array} \right].
 \end{aligned}$$

The Matrix View of Swapping the Order of Equations

This system of equations is neither upper triangular nor lower triangular

$$3x_1 = 6$$

$$x_1 - 2x_2 + 3x_3 = 2$$

$$2x_1 - x_2 = -2,$$

The Matrix View of Swapping the Order of Equations

but if we simply re-arrange the order of the equations, we arrive at the lower triangular equations

$$3x_1 = 6$$

$$2x_1 - x_2 = -2$$

$$x_1 - 2x_2 + 3x_3 = 2.$$

The Matrix View of Swapping the Order of Equations

Let's write out the matrix equations for the “unfortunately ordered” equations and then the “nicely” re-arranged system of equations.

$$\underbrace{\begin{bmatrix} 3 & 0 & 0 \\ 1 & -2 & 3 \\ 2 & -1 & 0 \end{bmatrix}}_{A_O} \underbrace{\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}}_x = \underbrace{\begin{bmatrix} 6 \\ 2 \\ -2 \end{bmatrix}}_{b_O},$$

$$\underbrace{\begin{bmatrix} 3 & 0 & 0 \\ 2 & -1 & 0 \\ 1 & -2 & 3 \end{bmatrix}}_{A_L} \underbrace{\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}}_x = \underbrace{\begin{bmatrix} 6 \\ -2 \\ 2 \end{bmatrix}}_{b_L}.$$

- ▶ We see the second and third rows are swapped when we compare A_O to A_L and b_O to b_L .

- ▶ We see the second and third rows are swapped when we compare A_O to A_L and b_O to b_L .

Claim

The swapping of rows can be accomplished by multiplying A_O and b_O on the left by

$$P = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}.$$

Indeed, we check that

$$P \cdot A_O = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 3 & 0 & 0 \\ 1 & -2 & 3 \\ 2 & -1 & 0 \end{bmatrix} = \begin{bmatrix} 3 & 0 & 0 \\ 2 & -1 & 0 \\ 1 & -2 & 3 \end{bmatrix} = A_L$$

Indeed, we check that

$$P \cdot A_O = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 3 & 0 & 0 \\ 1 & -2 & 3 \\ 2 & -1 & 0 \end{bmatrix} = \begin{bmatrix} 3 & 0 & 0 \\ 2 & -1 & 0 \\ 1 & -2 & 3 \end{bmatrix} = A_L$$

and

$$P \cdot b_O = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 6 \\ 2 \\ -2 \end{bmatrix} = \begin{bmatrix} 6 \\ -2 \\ 2 \end{bmatrix} = b_L.$$

Indeed, we check that

$$P \cdot A_O = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 3 & 0 & 0 \\ 1 & -2 & 3 \\ 2 & -1 & 0 \end{bmatrix} = \begin{bmatrix} 3 & 0 & 0 \\ 2 & -1 & 0 \\ 1 & -2 & 3 \end{bmatrix} = A_L$$

and

$$P \cdot b_O = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 6 \\ 2 \\ -2 \end{bmatrix} = \begin{bmatrix} 6 \\ -2 \\ 2 \end{bmatrix} = b_L.$$

P is called a permutation matrix.

```
In [1]: A0 = [3. 0 0; 1 -2 3; 2 -1 0];  
b0 = [6.; 2; -2];  
P = [1 0 0; 0 0 1; 0 1 0];  
  
AL = P * A0
```

```
Out[1]: 3x3 Array{Float64,2}:  
 3.0  0.0  0.0  
 2.0 -1.0  0.0  
 1.0 -2.0  3.0
```

```
In [2]: bL = P * b0
```

```
Out[2]: 3-element Array{Float64,1}:  
 6.0  
-2.0  
 2.0
```

Remark

We note that the permutation matrix P is constructed from the 3×3 identity matrix I by swapping its second and third rows, exactly the rows we wanted to swap in A_O and b_O . This observation works in general.

Suppose we want to do the following rearrangement

$$\begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \end{bmatrix} \rightarrow \begin{bmatrix} 1 \\ 2 \\ 5 \\ 4 \\ 3 \end{bmatrix},$$

To see the resulting structure at a matrix level, we put the 5×5 identity matrix on the left and the permutation matrix P on the right

$$I = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix} \rightarrow P = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 \end{bmatrix}$$

so that it is apparent that P is just a re-ordering of the rows of I .

```
In [3]: v = [1; 2; 3; 4; 5];  
P = [1 0 0 0 0; 0 1 0 0 0; 0 0 0 0 1; 0 0 0 1 0; 0 0 1 0 0]
```

```
Out[3]: 5x5 Array{Int64,2}:  
 1  0  0  0  0  
 0  1  0  0  0  
 0  0  0  0  1  
 0  0  0  1  0  
 0  0  1  0  0
```

```
In [4]: # permute v  
P * v
```

```
Out[4]: 5-element Array{Int64,1}:  
 1  
 2  
 5  
 4  
 3
```

- ▶ LU (Lower-Upper) Factorization
- ▶ Read Chapter 5 of ROB 101 Book