

MINISTRY OF EDUCATION, SINGAPORE
in collaboration with
CAMBRIDGE ASSESSMENT INTERNATIONAL EDUCATION
General Certificate of Education Ordinary Level

CANDIDATE
NAME



CENTRE
NUMBER

S				
---	--	--	--	--

INDEX
NUMBER

--	--	--	--

ADDITIONAL MATHEMATICS

4049/01

Paper 1

October/November 2025

2 hours 15 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, index number and name in the spaces at the top of this page.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE ON ANY BARCODES.

Answer **all** the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an approved scientific calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 90.

This document consists of **16** printed pages.



Singapore Examinations and Assessment Board



Cambridge Assessment
International Education



Mathematical Formulae

1. ALGEBRA

*Quadratic Equation*For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial expansion

$$(a + b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{n(n-1)\dots(n-r+1)}{r!}$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2 \cos^2 A - 1 = 1 - 2 \sin^2 A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

Formulae for $\triangle ABC$

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2}bc \sin A$$





- 1 An item is to be sold online. At the start of the month, it is placed on sale at an initial price of \$300. At the end of each month, if the item hasn't sold, the price is automatically reduced by 5%. It remains on sale at this price for the next month. The item finally sells in the month when the item is first priced below \$120.

How many complete months have elapsed before the item is sold?

[5]

- 2 Find the set of values of the constant k for which the curve $y = kx^2 + 2x + k + 1$ is completely below the line $y = x + 1$.

[6]





3 (a) (i) Factorise $a^3 - b^3$.

[1]

(ii) Hence state a factor, other than itself and 1, of $347^3 - 286^3$.

[1]

It is given that x and n are positive integers.

(b) Use **part (a)(i)** to prove that $\frac{(x+n)^3}{2} - \frac{(x-n)^3}{2}$ is divisible by n .

[3]





4 (a) Differentiate $x^{n+1} \ln x$ with respect to x , where $n \geq 0$. [2]

(b) Hence find $\int x^n \ln x dx$. [3]





- 5 Two variables x and y are believed to be related by the equation $y = Ax^b$ where A and b are constants. When a set of points (X, Y) are plotted, where X is a function of x and Y is a function of y , the resulting graph is a straight line.

(a) Given that $\ln y$ is plotted on the Y -axis determine, with working, the function of x which must be plotted on the X -axis. [2]

(b) Given that two of the points lying on the straight line are $(1.2, 6.3)$ and $(4.1, 12.1)$ find the values of A and b . [4]



- 6 (a) Show that the x -coordinate of the stationary point on the curve $y = \frac{1}{2}x^2 - 3\sqrt{x} + 1$ is $\left(\frac{3}{2}\right)^{\frac{2}{3}}$. [4]

- (b) Determine the nature of this stationary point. [3]





- 7 (a) Find, in ascending powers of x , the first four terms in the expansion of $\left(\frac{1}{2} - ax\right)^5$, where a is a constant. Simplify the coefficients in your expansion. [4]

- (b) In the expansion of $\left(4 + \frac{5}{ax}\right)\left(\frac{1}{2} - ax\right)^5$, find the non-zero value of a for which the coefficient of x is twice the coefficient of x^2 . [3]



8 A curve has equation $\frac{(x-a)^2}{2} + (y-1)^2 = 1$, where a is a constant.

(a) Find the set of values of a for which the line $y = x + 1$ intersects this curve at two distinct points. [5]

(b) Deduce the two values of a for which the line $y = x + 1$ is a tangent to the curve. Show your working clearly. [2]





9 (a) Prove that $\frac{1 + \sin 2\theta}{\sin \theta} - \frac{1 + \cos 2\theta}{\cos \theta} = \operatorname{cosec} \theta$.

[4]

(b) Hence solve $\frac{1 + \sin 2\theta}{\sin \theta} - \frac{1 + \cos 2\theta}{\cos \theta} = 2 \sin \theta$ for $\pi < \theta < \frac{3}{2}\pi$.

[3]






- 10 (a) Find, in terms of the positive constant a , the equation of the normal to the curve $y = 2 \sin ax - 3 \cos 2x$ at the point P where $x = 0$. [7]

- (b) The normal cuts the x -axis at Q . The area of triangle OPQ , where O is the origin, is 27 units^2 . Find the value of a . [3]





11



A square $ABCD$ is part of a computer animation. The side AB is the hypotenuse of a right-angled triangle AEB with angle $AEB = 90^\circ$. The points A and E are fixed and do not change during the animation.

The length of AE is always 3 units and angle AEB is always 90° .

During the animation B moves directly away from E along the line perpendicular to AE and, at time t seconds, the distance EB is x units.

It is given that x is increasing at a constant rate of 2 units/s.

(a) Find the rate of change of the length AB , in units/s, when $x = 4$.


[5]



- (b) The rate of change of the area of the square cannot exceed $80 \text{ units}^2/\text{s}$. Find the maximum possible value for x . [3]



12



The diagram shows part of the curve $y = a \cos kx$ where a and k are positive constants and x is in radians. The curve crosses the positive x -axis at A and B .

- (a) (i) Find, in terms of k , the x -coordinates of A and of B . [2]

Point C is the minimum point on the curve between A and B .
Triangle ABC is equilateral.

- (ii) Prove that $a = \frac{\pi\sqrt{3}}{2k}$. [3]



- (b) Given that the area of triangle ABC is 1 unit^2 , show that $k = \frac{(3)^{\frac{1}{4}}\pi}{2}$. Hence find the amplitude and period of the curve. [3]

Question 13 is printed on the next page.



- 13 Points $A(-1, 7)$ and B lie on a circle.

The chord AB of this circle has equation $2y - 3x = 17$.

The centre of the circle lies on the line $y = -2$.

The diameter of the circle parallel to AB passes through the point $\left(-\frac{1}{2}, \frac{1}{4}\right)$.

- (a) Find the equation of the circle.

[7]

- (b) The point $(7, \alpha)$ lies **inside** the circle. Find the set of possible values for α .

[2]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

