

MINISTRY OF EDUCATION, SINGAPORE
in collaboration with
CAMBRIDGE ASSESSMENT INTERNATIONAL EDUCATION
General Certificate of Education Ordinary Level

CANDIDATE
NAME



CENTRE
NUMBER

S				
---	--	--	--	--

INDEX
NUMBER

--	--	--	--

ADDITIONAL MATHEMATICS

4049/02

Paper 2

October/November 2025

2 hours 15 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, index number and name in the spaces at the top of this page.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE ON ANY BARCODES.

Answer **all** the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an approved scientific calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 90.

This document consists of **17** printed pages and **3** blank pages.



Singapore Examinations and Assessment Board



Cambridge Assessment
International Education



Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial expansion

$$(a + b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{n(n-1)\dots(n-r+1)}{r!}$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2 \cos^2 A - 1 = 1 - 2 \sin^2 A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

Formulae for $\triangle ABC$


$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2}bc \sin A$$



1 Do not use a calculator in this question.



The diagram shows a trapezium $ABCD$ with BC and AD parallel. The length of AD is $(6 + \sqrt{3})$ cm and the perpendicular height of the trapezium is $(4 - \sqrt{3})$ cm. The length of BC is l cm. Given that the trapezium has area 26 cm^2 , find the exact length of BC . Give your answer in the form $a + b\sqrt{3}$ where a and b are integers. [6]





- 2 On a winter's day the temperature, T °C, h hours after measurement began, is modelled by the formula $T = 4 + 2h - \frac{1}{2}h^2$, where $0 \leq h \leq 8$.
- (a) By expressing T in the form $A - B(h + C)^2$, where A , B and C are constants, find the time at which the temperature is greatest and state the value of this temperature. [5]

- (b) On the following day, and at the same times as before, the temperatures are found to be half their previous day's values. State the time at which the temperature is greatest and the value of this temperature. [2]





- 3 The gradient of a curve, at a point (x, y) on the curve, is given by $e^x(3e^{2x} - e^{-4x}) - \cos 2x$. The curve passes through the origin. Find y in terms of x . [7]



**4 Do not use a calculator in this question.**

- (a) It is given that $f(x) = x^3 + px^2 + qx - 6$, where p and q are constants.
 $(x + 1)$ is a factor of $f(x)$.

When $f(x)$ is divided by $(x + 2)$ there is a remainder of 4.

Find the values of p and q .

[6]

- (b) It is given that $3x - 1$ is a factor of $g(x) = 9x^3 - 18x^2 - x + 2$. Express $g(x)$ as a product of three linear factors.

[3]





5 Given that $y = \frac{1 - e^{-x}}{1 + e^{-x}}$, prove that y increases for all real values of x .

[6]

DO NOT WRITE IN THIS MARGIN





6 Solve the equation $2 \operatorname{cosec}^2 \theta - \cot \theta - 3 = 0$ for $0^\circ \leq \theta \leq 360^\circ$.

[7]





- 7 A particle is travelling in a straight line. Its acceleration, $a \text{ m/s}^2$, at time t seconds is given by

$$a = (2t + 1)^{-1} + 0.3t + 0.2 \text{ for } 0 \leq t \leq 10.$$

At $t = 0$ the velocity of the particle is -3 m/s .
Find the velocity of the particle when $t = 5$.

[7]





8 (a) Solve the equation $2^p = 5(3^{2-p})$.

[4]

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN






(b) Solve the simultaneous equations

$$\begin{aligned}9(2^x) + 2(3^{y+1}) &= 38, \\15(2^{x-1}) - 21(3^y) &= 23.\end{aligned}$$

[5]





The diagram shows the line $y = kx$, where k is a constant, $k < 10$, and the curve $y = x(10-x)$. The line intersects the curve at the origin O and at point P . Point $Q(\alpha, 0)$, where α is a constant, lies on the x -axis such that PQ is parallel to the y -axis.

R_1 is the area of the region enclosed by the curve and the line OP .

(a) By finding k in terms of α , show that $R_1 = \frac{\alpha^3}{6}$.

[7]






R_2 is the area of the region enclosed by the lines OP , PQ and the x -axis.

(b) Find the numerical value of k if $R_1 = R_2$.

[3]



10



The diagram shows a quadrilateral $ABCD$ whose vertices lie on the circumference of a circle. The tangent to the circle at B meets the line AC extended at the point E . Angle $BAC = x$ and angle $FBA = y$, where F is a point on EB extended.

(a) Prove that triangle ABE is similar to triangle BCE .

[3]



(b) (i) Prove that angle $ADC = x + y$.

[3]

(ii) If DC is parallel to AB prove that $AD = BC$.

[4]





- 11 At time $t = 0$ seconds, two objects are at points A and B which are 30 cm apart. The objects are moving directly towards one another in a straight line. Their constant speeds are $6 \cos \theta$ cm/s and $8 \sin \theta$ cm/s respectively, for a certain value of θ measured in degrees. The radii of the objects may be ignored.

(a) Explain why θ must be acute.

[1]

- (b) In the case when the objects collide at the midpoint of AB , find the time taken for the objects to collide.

[3]

- (c) (i) If, instead, the objects collide when $t = 4$ seconds show that $3 \cos \theta + 4 \sin \theta = 3.75$.

[2]





- (ii) By writing $3 \cos \theta + 4 \sin \theta$ in the form $R \cos(\theta - \alpha)$, where $R > 0$ and α is acute, find the numerical values of the speeds of the two objects. [6]





BLANK PAGE

DO NOT WRITE IN THIS MARGIN





DO NOT WRITE IN THIS MARGIN





DO NOT WRITE IN THIS MARGIN

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

