

- 1 **Wildlife health perceptions and monitoring practices in globally distributed protected**
- 2 **areas**

## Abstract

Deficits in wildlife health (WH) monitoring at protected areas (PAs) can weaken the detection of infectious diseases; physical, and chemical threats; rapid response; and assessment of health management practices, threatening biodiversity conservation and global health. However, there is a lack of baseline information regarding the local perception of wildlife, human, and livestock health relevance at these sites. Current WH monitoring practices and WH data collection and management in PAs are also unknown. To address these gaps, we conducted a survey targeting globally distributed protected area data managers (PADMs). Eighty-six valid responses were considered for analysis.

PADMs considered WH as relevant to the conservation goals of PAs and >90% of them confirmed that non-healthy wildlife (injured, sick, and dead) are encountered. However, >50% and >20% of PADMs claimed that these animals were not recorded, respectively. When these animals were documented, the recording methods and information collected differed. Although domestic animal presence was common and considered a conservation concern, these animals and their health status were not always recorded. Health data were often stored in a database, but paper forms and spreadsheets were also used.

Responses suggest that valuable syndromic WH surveillance data from PAs are being lost due to non-collection or inadequate management and their value could be limited by unstandardized recording. Rangers could become a globally distributed “One Health workforce” but these flaws must be addressed first.

## Introduction

Despite the growing recognition that the health of animals, people, and their shared environment are inseparably linked, effective surveillance systems collecting, analyzing, and responding to wildlife health (WH) data remain uncommon or deficient (Machalaba et al. 2021; One Health High-Level Expert Panel et al. 2022; World Organization for Animal Health [WOAH] 2023; Delgado et al. 2023). Human encroachment and land-use change (Laurance et al. 2012; Vicente et al. 2021; Meng et al. 2023) are associated with extraction, pollution, the creation of human-wildlife-livestock interfaces, and ecosystem degradation (Plowright et al. 2021; Vicente et al. 2021; Reaser et al. 2023). These processes expose wildlife and people to physical (e.g., snaring, fire), chemical (e.g., poisoning events), and biological hazards (e.g., pathogens) with the capacity to impact biodiversity conservation and global health (De Vos et al. 2016; Vila et al. 2019; Wolf et al. 2019; Hacon et al. 2020; Machalaba et al. 2020; Becker et al. 2023; Groenenberg et al. 2023; Porco et al. 2023).

Ebola virus disease is an illustrative example. Index cases of several human outbreaks of this disease have included hunters who contacted wildlife (Judson et al. 2016). Outbreaks in wildlife have decimated populations of western gorillas and chimpanzees (Whitfield 2003; Leroy et al. 2004; Bermejo et al. 2006) and have preceded outbreaks in humans (Rouquet et al. 2005).

Protected areas (PAs), nature's last strongholds, are increasingly facing anthropogenic pressures becoming key human-wildlife interfaces for disease emergence and prevention (Mittermeier et al. 2011; Machalaba et al. 2021; IUCN & EcoHealth Alliance 2022; WOAHA 2023; Hayman et al. 2023; Hopkins et al. 2024). WH monitoring at these sites can strengthen the detection of infectious diseases, physical, and chemical threats; rapid response; and assessment of health management practices. However, there is a lack of baseline information regarding the perception of wildlife, human, and livestock health relevance for biodiversity conservation. Current WH monitoring practices and WH data collection and management at these sites are also unknown. To address these gaps, we conducted a survey targeting protected area data managers (PADMs) to assess: i) their perceptions regarding WH and pathogen transmission between wildlife, humans, and livestock, ii) the detection and documentation of dead, sick, or injured wildlife, and domestic animals in PAs, and iii) health data management in PAs.

## Methods

The "Spatial Monitoring and Reporting Tool" (SMART) is a technology platform designed to support the administration of PAs (Cronin et al. 2021) through the collection, management, assessment, and communication of PA data. SMART technology is distributed globally in more than 1,000 conservation sites by the time of the study.

We developed a web-based questionnaire aimed at these PADM SMART-users. First, respondents were asked if their job roles and responsibilities matched the definition "a person directly responsible for managing SMART data in one or more PAs or a general manager or administrator of one or more PAs that uses SMART data". Respondents who did not identify as a PADM were considered outside our target population and excluded. The survey had five sections. Section 1 assessed the perception of PADM on the importance of WH in achieving conservation goals, the role of human and livestock pathogens in affecting WH, and the role of wildlife pathogens in affecting public and livestock health. Section 2 requested PADM to rank the overall frequency of encounters with dead, sick, or injured wildlife in PAs and their documentation when found during patrols. Section 3, asked about the presence of domestic animals in the PA(s), the documentation of their health status, and the perceived threats of domestic animals to conservation goals. Likert scales were used to answer questions in Sections 1-3. Section 4 addressed health data storage practices when collected and Section 5 assessed the current state of SMART deployment in PAs.

An introductory web page explained that the survey was voluntary, anonymous, aimed at PADM, and that clicking the "Start the survey" button constituted consent. A tutorial was provided for the language-translation tool of this survey built on Google Forms (Supporting Information). The survey did not request any personal information or demographic characteristics and, consequently, it was exempt from full ethics review [placeholder for Institutional Review Board to keep authors' anonymity during peer review].

The survey was distributed globally to the SMART Community Forum users by the SMART Partnership (<https://smartconservationtools.org>) via email in October 2022 and remained open for three months. A reminder was sent to the SMART Community three weeks before the closing date.

Responses by PADM s could represent single or multiple PAs. For our analysis, we focused on what we considered to be “local” responses which included one or two PAs, and assumed they provided insights into specific local realities. “Non-local” responses represented more than two PAs, which were assumed to help understand perceptions at the decision-making level and were analyzed separately (see Supporting Information). We discarded responses that only included marine PAs based on the World Database on Protected Areas (<https://www.protectedplanet.net/en/thematic-areas/wdpa?tab=WDPA>) as marine PA management, species, and patrol logistics are markedly different. The descriptive analysis of survey responses was conducted in R v4.3.1. The responses dataset can be found here <https://figshare.com/s/36513db82ac5dfa8e71d?file=49682265>. Descriptive analysis can be found at [placeholder for github repository to keep authors’ anonymity during peer review].

## Results

We received 128 responses. Forty-two were removed because either the PA name(s) were not provided, only marine PAs were listed, or the respondents did not match the target audience (8, 7, and 27 responses, respectively). The final dataset contained 86 respondents from 23 countries. Seventy-three were local responses from 19 countries and 13 were non-local responses from 10 countries (the descriptive analysis of non-local responses is provided in the Supporting Information). The specific countries are not provided to protect the identity of the respondents, but local responses were from North, Central, and South America; West, Central, East, and Southern Africa; Southeast and South Asia; and a Balkan country in Europe, with most coming from South America. Non-local responses were from West, Central, and East Africa, Central and South America, and Southeast and South Asia.

### **Perceptions regarding wildlife health importance in conservation and potential consequences of pathogen transmission among wildlife, domestic animals, and people**

Most respondents either strongly agreed or agreed with the affirmations “Wildlife health, including infectious and non-infectious diseases, is important to achieve the conservation goals of the protected areas where I work” and “human or livestock pathogens can affect wildlife populations inhabiting the protected area(s) I work in” (92%; and 81%, respectively). Regarding the affirmation “pathogens carried by wildlife inhabiting the protected area(s) where I work in can affect livestock health”, most respondents strongly agreed or agreed (48%) although the proportion of neutral respondents was more prominent (29%). Across respondents, 63% strongly agreed or agreed that “pathogens carried by wildlife inhabiting the protected area(s) where I work in can affect human health”. Detailed response distributions are shown in Figure 1. Non-local responses followed similar trends (Figure S1).

**Overall frequency of encounters with dead, sick, or injured wildlife in protected areas and their documentation when found during patrols**

Most local PADM s (97%) reported the encounter of dead animals in the PAs (e.g., "Very rarely" or more frequently), and 76% of them reported documenting these encounters if found during patrols. Similar to dead animal encounters, 93% of local PADM s reported encounters with sick or injured animals in the PAs (e.g., "Very rarely" or more frequently), but only 35% and 48% confirmed their documentation, respectively. In general, the documentation of injured, sick, or dead animals tend to be higher as the encounter frequency increased (Figure 2). All non-local PADM s reported the encounter with dead and injured or sick wildlife ("Very rarely" up to "Very frequently"). The proportions of non-local PADM s reporting the documentation of these animals were larger compared to the proportions from local responses (92, 62, and 85%, for dead, sick, and injured wildlife respectively, Figure S2).

All 17 local PADM s who ranked encountering dead wildlife "Very rarely" or more frequently but answered that these animals were not documented, either agreed or strongly agreed with the importance of WH to achieve conservation goals. Similarly, 91 and 94% of local PADM s who ranked encountering sick or injured wildlife "Very rarely" or more frequently but answered that these animals were not documented either agreed or strongly agreed with this statement. The corresponding percentages for non-local PADM s were 0, 80 and 50% for dead, sick and injured wildlife).

The documentation method of dead, sick, or injured wildlife, varied among the 58 local PADM s that reported the recording of one or more of these groups. Most often, each animal was documented individually ("Individual observation"). The second most common method involved a complete inventory of healthy, sick, injured, or dead animals for each species ("Part of the full count"). Reporting their presence or absence was the third most common method ("Present or absent"; Table 1). For non-local responses, the predominant method was "each animal is an individual observation" across health categories (Table S1).

The data collected from each observation were not consistent. Across health categories and documentation method, photographs and the species were the main item collected. Anomalies observed in non-healthy wildlife and the condition of carcasses were not always recorded (Table 1). In non-local responses the trend was relatively similar, however, items were reported to be recorded more consistently (e.g., age, anomalies, and condition in the three health categories; Table S1).

**Presence of domestic animals in protected areas, the documentation of their health status, and the perceived threats of domestic animals to conservation goals**

Fifty-two local PADM s (71%) responded that domestic animals were found in the corresponding PAs. Among them, 67% reported that domestic animals were documented if observed during patrols, but only 26% reported recording their health status (Figure 3). Forty-two local respondents

reporting domestic animals in the PAs (81%) either agreed or strongly agreed that domestic animals are a conservation concern (Figure 3). Twenty-seven of them (64%) answered that these animals were documented. Most respondents claiming that domestic animals are not found in the corresponding PAs also either agreed or strongly agreed that they are a conservation concern.

Eight non-local PADM s (62%) responded that domestic animals were found in the PAs. Among them, seven (88%) reported that domestic animals were documented if observed during patrols, of which only two (29%) reported recording their health status (Figure S3).

### **Health data storage practices in protected areas**

Seventy-two, 54, and 65% of local PADM s reporting the documentation of either injured, sick, or dead wildlife stored all these data in a SMART database. Paper forms, reports, and spreadsheets were employed when non-healthy wildlife were documented but their data was not stored in a SMART database.

Thirty-one of the 35 local respondents (89%) recording domestic animals during patrols indicated that this information was stored in a SMART database. The health status of domestic animals, when documented, was also stored in a SMART database.

## **Discussion**

We developed a web-based questionnaire aimed at globally distributed PADM s to learn about their perceptions regarding WH; the monitoring of dead, sick, and injured wildlife and domestic animals in PAs; and health data storage practices. Regardless of any geographic biases in this study, the responses suggest that valuable syndromic WH and, consequently, One Health surveillance data are being lost due to non-collection or inadequate management. Even when WH data are collected and properly managed, the usefulness for surveillance is likely limited by the diversity of methods employed to record them.

PADM s largely considered WH as relevant to the conservation goals of PAs and most of them confirmed that dead, sick, or injured wildlife were encountered “Very rarely” or more frequently. However, the documentation of these animals was not always conducted, and it was less common in responses from local data managers. This contradiction could be explained by recent global pathogen-driven crises such as SARS-CoV-2 and H5N1 Highly Pathogenic Avian Influenza virus (Nicola et al. 2020; Leguia et al, 2023) that might have sensitized our audience by the time the survey was distributed but before health-associated monitoring objectives could be planned and rolled out. These findings could also suggest a lack of knowledge or resources to act on their understanding of the importance of WH for conservation goals. Although only 13 non-local responses were included in the final dataset, the larger proportion of non-local responses reporting the documentation of non-healthy wildlife could suggest differences between the expectations of managers in an administrative role and field realities in PAs. Protected area management agencies should take a more active local role to identify and correct weaknesses in WH data collection.

We noted a general agreement among PADMIs regarding the conservation threat that domestic animals (e.g., dogs, cats, cattle) present. Although we did not explicitly ask why domestic animals are a conservation concern, most PADMIs also agreed with the statement “human and wildlife pathogens can impact wildlife health” either when these animals were found in the PAs or not (Figure S4 and S5). This finding might imply that pathogens are part of the reason why domestic animals are considered a conservation concern. Pathogen transmission from domestic animals to wildlife can seriously harm biodiversity conservation efforts including in PAs (e.g., del Valle Ferreyra et al. [2022]) and they add to the direct and indirect pressures on wildlife from domestic animals, such as predation, competition, disturbance, and land-use change in- and out-side of PAs (du Toit 2011; Gompper 2013). While most local PADMIs reported the presence of domestic animals in PAs, their documentation was not consistent and only a minority responded that their health status was recorded. The contradiction between perceived conservation risk of domestic animals and documentation of their presence and health status could be explained by the same drivers mentioned above.

Adequate management of data and harmonization are foundational pillars for WH monitoring (WOAH 2010, 2015, 2018; Sleeman et al. 2012; Ryser-Degiorgis 2013; Stephen 2018; Lawson et al. 2021; Machalaba et al. 2021; Giacinti et al. 2022; Stephen & Berezowski 2022; Hayman et al. 2023; Heiderich et al. 2023). However, we found that paper forms and Excel sheets are used to store data from sick, injured, or dead wildlife even when SMART was available. Also, we found diverse structure and attributes in WH data that was collected. The lack of harmonization across PAs within and beyond country boundaries can limit the value of collected health data as regional, national, or across border health assessments could be unfeasible. Similarly, the longitudinal assessment of wildlife health trends in a single PA is not possible when data is recorded differently over time. These findings are aligned with historical pitfalls in WH surveillance’s data governance (e.g., Avery-Gomm et al. 2016; Lawson et al. 2021; Heiderich et al. 2023; WOA 2023).

## Recommendations

Leveraging existing PA human resources that can detect morbidity and mortality in animals offers a sound and cost-effective strategy to establish a minimal baseline of WH monitoring. Rangers can detect injured, sick, and dead animals in PAs and the few documented initiatives that explicitly report ranger participation in WH monitoring have demonstrated their potential to provide data to assess health risks and trends or trigger responses to disease outbreaks of global and conservation concern (Kuisma et al. 2019; Vila et al. 2019; Wolf et al. 2019; Montecino-Latorre et al. 2020; Orozco et al. 2020; Porco et al. 2023). Rangers also have contributed to a healthier planet through emergency medical assistance, contact tracing, and public health education (Singh et al. 2021; Stolton et al. 2023). Currently, there are approximately 280,000 rangers worldwide and it is estimated that 1.5 million will be needed by 2030 to adequately protect 30% of the planet (Appleton et al. 2022). The present and projected number of rangers reveal their unique potential as a worldwide distributed “One Health workforce” that could drastically improve the general

225 global absence of WH and One Health surveillance (Machalaba et al. 2021; Worsley-Tonks et al.  
 226 2022; Delgado et al. 2023; Hopkins et al. 2024).

227 Our recommendation is to include WH monitoring within the remit of rangers. The global  
 228 integration of this workforce into WH monitoring could be supported by SMART or the World  
 229 Commission of Protected Areas (WCPA) of the International Union for Conservation of Nature,  
 230 all of which support best practice management of PAs. Indeed, the WCPA has established a two-  
 231 year Task Force to integrate One Health in PAs and vice versa which includes WH surveillance  
 232 activities in PAs (Hopkins et al., 2024). This Task Force makes explicit the contemporaneous  
 233 relevance to develop WH surveillance policy in PAs to support the conservation of biodiversity  
 234 and global health.

235 Technology can enhance ranger integration into WH surveillance systems. The engagement of a  
 236 large community of SMART-using rangers to document “health incidents” in PAs represents a  
 237 promising opportunity to create a technology-supported worldwide network of WH sentinels  
 238 (Worsley-Tonks et al. 2022). Other platforms such as EarthRanger could also support the  
 239 integration.

240 However, there are key issues that must be addressed before. First, a definition of a health event  
 241 optimized for rangers must be established. The minimal set of variables to be recorded from each  
 242 health event and their documentation method must be harmonized across jurisdictions. Specific  
 243 variables and options can then be tailored to individual PA realities. Second, rangers need to be  
 244 trained to recognize and document health events encountered during their patrols. Third, a database  
 245 to guarantee the governance of ranger-documented health events must be available. SMART  
 246 technology is ready to support the management of harmonized syndromic health data, provided  
 247 that adequate resources for planning, training, and expert support are available. PADMs working  
 248 with fully implemented SMART should have the capacity to properly manage and query health  
 249 data and coordinate their distribution with other relevant parties, such as environmental agencies  
 250 and organizations, veterinary services, and public health managers.

251 Efforts to address the issues mentioned above and build this workforce have already started.  
 252 WildHealthNet, a Wildlife Conservation Society (WCS) initiative, focuses on creating national  
 253 surveillance networks and codifying their Standard Operating Procedures (Denstedt et al. 2021;  
 254 Porco et al. 2023; Pruvot et al. 2023). Currently, WCS is supporting the integration of rangers into  
 255 WH monitoring using the same standards in Lao, Peru, Cambodia, Guatemala, and Madagascar.  
 256 Additional efforts are underway through the Wildlife Health Intelligence Network (Noguera et al.  
 257 2024) whose objectives include building a larger community of practice that supports the scaling  
 258 of local WH surveillance globally, and the establishment of WH data collection and management  
 259 standards. We strongly recommend taking steps towards the global adoption of ranger-based WH  
 260 monitoring in PAs, utilizing a unified methodology and standards to support biodiversity  
 261 conservation and improve the health of all.



## References

- Appleton, M.R., Courtiol, A., Emerton, L., Slade, J.L., Tilker, A., Warr, L.C., Malvido, M.Á., Barborak, J.R., de Bruin, L., Chapple, R., Daltry, J.C., Hadley, N.P., Jordan, C.A., Rousset, F., Singh, R., Sterling, E.J., Wessling, E.G. & Long, B. (2022). Protected area personnel and ranger numbers are insufficient to deliver global expectations. *Nature Sustainability*, 5, 1100–1110.
- Avery-Gomm, S., Valliant, M., Schacter, C.R., Robbins, K.F., Liboiron, M., Daoust, P.-Y., Rios, L.M. & Jones, I.L. (2016). A study of wrecked Dovekies (*Alle alle*) in the western North Atlantic highlights the importance of using standardized methods to quantify plastic ingestion. *Mar. Pollut. Bull.*, 113, 75–80.
- Becker, D.J., Eby, P., Madden, W., Peel, A.J. & Plowright, R.K. (2023). Ecological conditions predict the intensity of Hendra virus excretion over space and time from bat reservoir hosts. *Ecol. Lett.*, 26, 23–36.
- Bermejo M, Rodríguez-Teijeiro JD, Illera G, Barroso A, Vilà C, Walsh PD. 2006. Ebola outbreak killed 5000 gorillas. *Science* 314,1564.
- Cronin, D.T., Dancer, A., Long, B., Lynam, A.J., Muntifering, J., Palmer, J., Bergl, R.A., Wich, S.A. & Piel, A.K. (2021). Application of SMART software for conservation area management. In: *Conservation Technology*. Oxford University Press Oxford, UK, pp. 201–224.
- Delgado, M., Ferrari, N., Fanelli, A., Muset, Thompson L., Sleeman, J. M., White, C. L., Walsh, D., Wannous, C. & Tizzani, P. (2023). Wildlife health surveillance: Gaps, needs and opportunities. *Rev. Sci. Tech.*, 42, 161-172.
- Denstedt, E., Porco, A., Hwang, J., Nga, N.T.T., Ngoc, P.T.B., Chea, S., Khammavong, K.,

285 Milavong, P., Sours, S., Osbjør, K., Tum, S., Douangngeun, B., Theppanya, W., Van Long,  
 286 N., Thanh Phuong, N., Tin Vinh Quang, L., Van Hung, V., Hoa, N.T., Le Anh, D., Fine, A.  
 287 & Pruvot, M. (2021). Detection of African swine fever virus in free-ranging wild boar in  
 288 Southeast Asia. *Transbound. Emerg. Dis.*, 68, 2669–2675.

289 De Vos, A., Cumming, G., Cumming, D., Ament, J., Baum, J., Clements, H., Grewar, J.,  
 290 Maciejewski, K. & Moore, C. (2016). Pathogens, disease, and the social-ecological  
 291 resilience of protected areas. *Ecol. Soc.*, 21.

292 del Valle Ferreyra, H., Rudd, J., Foley, J., Vanstreels, R. E. T., Martín, A. M., Donadio, E., &  
 293 Uhart, M. M. (2022). Sarcoptic mange outbreak decimates South American wild camelid  
 294 populations in San Guillermo National Park, Argentina. *Plos One*, 17, e0256616.

295 Giacinti, J.A., Jane Parmley, E., Reist, M., Bayley, D., Pearl, D.L. & Jardine, C.M. (2022).  
 296 Canadian wildlife health surveillance—patterns, challenges and opportunities identified by  
 297 a scoping review. *Facets (Ott)*, 7, 25–44.

298 Gompper, M.E. (2013). *Free-Ranging Dogs and Wildlife Conservation*. OUP Oxford.

299 Groenenberg, M., Crouthers, R., Yoganand, K., Banet-Eugene, S., Bun, S., Muth, S., Kim, M.,  
 300 Mang, T., Panha, M., Pheaktra, P., Pin, T., Sopheak, K., Sovanna, P., Vibolratanak, P.,  
 301 Wyatt, A.G. & Gray, T.N.E. (2023). Snaring devastates terrestrial ungulates whilst sparing  
 302 arboreal primates in Cambodia’s Eastern Plains Landscape. *Biol. Conserv.*, 284, 110195.

303 Hacon, S. de S., Oliveira-da-Costa, M., Gama, C. de S., Ferreira, R., Basta, P.C., Schramm, A. &  
 304 Yokota, D. (2020). Mercury Exposure through Fish Consumption in Traditional  
 305 Communities in the Brazilian Northern Amazon. *Int. J. Environ. Res. Public Health*, 17.

306 Hayman, D.T.S., Adisasmito, W.B., Almuhairi, S., Behraves, C.B., Bilivogui, P., Bukachi,  
 307 S.A., Casas, N., Becerra, N.C., Charron, D.F., Chaudhary, A., Ciacchi Zanella, J.R.,

308 Cunningham, A.A., Dar, O., Debnath, N., Dungu, B., Farag, E., Gao, G.F., Khaita, M.,  
 309 Machalaba, C., Mackenzie, J.S., Markotter, W., Mettenleiter, T.C., Morand, S., Smolenskiy,  
 310 V., Zhou, L. & Koopmans, M. (2023). Developing One Health surveillance systems. *One*  
 311 *Health*, 100617.

312 Heiderich, E., Keller, S., Pewsner, M., Origgi, F.C., Zürcher-Giovannini, S., Borel, S., Marti, I.,  
 313 Scherrer, P., Pisano, S.R.R., Friker, B., Adrian-Kalchhauser, I. & Ryser-Degiorgis, M.-P.  
 314 (2023). Analysis of a European general wildlife health surveillance program: chances,  
 315 challenges and recommendations. *bioRxiv*.

316 Hopkins, S. R., Olson, S.H., Fairban, H. T., Redford, K. H., Adams, J., Mitchell, B. A., Nova,  
 317 N., Muylaert, R. L., Morand, S., Miller, A. & Rao M. (2024). Protected areas and One  
 318 Health. *Parks*, 30, 6 – 13.

319 IUCN & EcoHealth Alliance. (2022). *Healthy people and wildlife through nature protection:*  
 320 *Guidelines for prevention, detection, response, and recovery from disease risks in and*  
 321 *around protected and conserved areas*.

322 Judson S.D., Fischer R., Judson A., Munster V.J. (2016). Ecological contexts of index cases and  
 323 spillover events of different ebolaviruses. *PLoS Pathog.*, 12, e1005780.

324 Kuisma, E., Olson, S.H., Cameron, K.N., Reed, P.E., Karesh, W.B., Ondzie, A.I., Akongo, M.-J.,  
 325 Kaba, S.D., Fischer, R.J., Seifert, S.N., Muñoz-Fontela, C., Becker-Ziaja, B., Escudero-  
 326 Pérez, B., Goma-Nkoua, C., Munster, V.J. & Mombouli, J.-V. (2019). Long-term wildlife  
 327 mortality surveillance in northern Congo: a model for the detection of Ebola virus disease  
 328 epizootics. *Philos. Trans. R. Soc. Lond. B Biol. Sci.*, 374, 20180339.

329 Laurance, W.F., Useche, D.C., Rendeiro, J., Kalka, M., Bradshaw, C.J.A., Sloan, S.P., Laurance,  
 330 S.G., Campbell, M., Abernethy, K., Alvarez, P., Arroyo-Rodriguez, V., Ashton, P., Benítez-

331 Malvido, J., Blom, A., Bobo, K.S., Cannon, C.H., Cao, M., Carroll, R., Chapman, C.,  
332 Coates, R., Cords, M., Danielsen, F., De Dijn, B., Dinerstein, E., Donnelly, M.A., Edwards,  
333 D., Edwards, F., Farwig, N., Fashing, P., Forget, P.-M., Foster, M., Gale, G., Harris, D.,  
334 Harrison, R., Hart, J., Karpanty, S., Kress, W.J., Krishnaswamy, J., Logsdon, W., Lovett, J.,  
335 Magnusson, W., Maisels, F., Marshall, A.R., McClearn, D., Mudappa, D., Nielsen, M.R.,  
336 Pearson, R., Pitman, N., van der Ploeg, J., Plumptre, A., Poulsen, J., Quesada, M., Rainey,  
337 H., Robinson, D., Roetgers, C., Rovero, F., Scatena, F., Schulze, C., Sheil, D., Struhsaker,  
338 T., Terborgh, J., Thomas, D., Timm, R., Urbina-Cardona, J.N., Vasudevan, K., Wright, S.J.,  
339 Arias-G, J.C., Arroyo, L., Ashton, M., Auzel, P., Babaasa, D., Babweteera, F., Baker, P.,  
340 Banki, O., Bass, M., Bila-Isia, I., Blake, S., Brockelman, W., Brokaw, N., Brühl, C.A.,  
341 Bunyavejchewin, S., Chao, J.-T., Chave, J., Chellam, R., Clark, C.J., Clavijo, J., Congdon,  
342 R., Corlett, R., Dattaraja, H.S., Dave, C., Davies, G., Beisiegel, B. de M., da Silva, R. de  
343 N.P., Di Fiore, A., Diesmos, A., Dirzo, R., Doran-Sheehy, D., Eaton, M., Emmons, L.,  
344 Estrada, A., Ewango, C., Fedigan, L., Feer, F., Fruth, B., Willis, J.G., Goodale, U.,  
345 Goodman, S., Guix, J.C., Guthiga, P., Haber, W., Hamer, K., Herbing, I., Hill, J., Huang,  
346 Z., Sun, I.F., Ickes, K., Itoh, A., Ivanauskas, N., Jackes, B., Janovec, J., Janzen, D.,  
347 Jiangming, M., Jin, C., Jones, T., Justiniano, H., Kalko, E., Kasangaki, A., Killeen, T.,  
348 King, H.-B., Klop, E., Knott, C., Koné, I., Kudavidanage, E., Ribeiro, J.L. da S., Lattke, J.,  
349 Laval, R., Lawton, R., Leal, M., Leighton, M., Lentino, M., Leonel, C., Lindsell, J., Ling-  
350 Ling, L., Linsenmair, K.E., Losos, E., Lugo, A., Lwanga, J., Mack, A.L., Martins, M.,  
351 McGraw, W.S., McNab, R., Montag, L., Thompson, J.M., Nabe-Nielsen, J., Nakagawa, M.,  
352 Nepal, S., Norconk, M., Novotny, V., O'Donnell, S., Opiang, M., Ouboter, P., Parker, K.,  
353 Parthasarathy, N., Pisciotta, K., Prawiradilaga, D., Pringle, C., Rajathurai, S., Reichard, U.,

354 Reinartz, G., Renton, K., Reynolds, G., Reynolds, V., Riley, E., Rödel, M.-O., Rothman, J.,  
 355 Round, P., Sakai, S., Sanaiotti, T., Savini, T., Schaab, G., Seidensticker, J., Siaka, A.,  
 356 Silman, M.R., Smith, T.B., de Almeida, S.S., Sodhi, N., Stanford, C., Stewart, K., Stokes,  
 357 E., Stoner, K.E., Sukumar, R., Surbeck, M., Tobler, M., Tschardtke, T., Turkalo, A.,  
 358 Umapathy, G., van Weerd, M., Rivera, J.V., Venkataraman, M., Venn, L., Vereza, C., de  
 359 Castilho, C.V., Waltert, M., Wang, B., Watts, D., Weber, W., West, P., Whitacre, D.,  
 360 Whitney, K., Wilkie, D., Williams, S., Wright, D.D., Wright, P., Xiankai, L., Yonzon, P. &  
 361 Zamzani, F. (2012). Averting biodiversity collapse in tropical forest protected areas. *Nature*,  
 362 489, 290–294.

363 Lawson, B., Neimanis, A., Lavazza, A., López-Olvera, J.R., Tavernier, P., Billinis, C., Duff, J.P.,  
 364 Mladenov, D.T., Rijks, J.M., Savić, S., Wibbelt, G., Ryser-Degiorgis, M.-P. & Kuiken, T.  
 365 (2021). How to Start Up a National Wildlife Health Surveillance Programme. *Animals*  
 366 (*Basel*), 11.

367 Leguia, M., Garcia-Glaessner, A., Muñoz-Saavedra, B., Juarez, D., Barrera, P., Calvo-Mac, C.,  
 368 Jara, J., Silva, W., Ploog, K., Amaro, L. & Colchao-Claux, P. (2023). Highly pathogenic  
 369 avian influenza A (H5N1) in marine mammals and seabirds in Peru. *Nat. Comm.*, 14,  
 370 p.5489.

371 Leroy EM et al. 2004. Multiple Ebola virus transmission events and rapid decline of central  
 372 African wildlife. *Science* 303:387–390.

373 Machalaba, C., Feferholtz, Y., Uhart, M. & Karesh, W.B. (2020). Wildlife conservation status  
 374 and disease trends: ten years of reports to the Worldwide Monitoring System for Wild  
 375 Animal Diseases. *Rev. Sci. Tech.*, 39, 991–1001.

376 Machalaba, C., Uhart, M., Ryser-Degiorgis, M.-P. & Karesh, W.B. (2021). Gaps in health

security related to wildlife and environment affecting pandemic prevention and preparedness, 2007-2020. *Bull. World Health Organ.*, 99, 342–350B.

Meng, Z., Dong, J., Ellis, E.C., Metternicht, G., Qin, Y., Song, X.-P., Löfqvist, S., Garrett, R.D., Jia, X. & Xiao, X. (2023). Post-2020 biodiversity framework challenged by cropland expansion in protected areas. *Nature Sustainability*, 1–11.

Mittermeier, R.A., Turner, W.R., Larsen, F.W., Brooks, T.M. & Gascon, C. (2011). Global Biodiversity Conservation: The Critical Role of Hotspots. In: *Biodiversity Hotspots: Distribution and Protection of Conservation Priority Areas* (eds. Zachos, F.E. & Habel, J.C.). Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 3–22.

Montecino-Latorre, D., Napolitano, C., Briceño, C. & Uhart, M.M. (2020). Sarcoptic mange: An emerging threat to Chilean wild mammals? *Perspect Ecol. Conserv*, 18, 267–276.

Nicola, M., Alsafi, Z., Sohrabi, C., Kerwan, A., Al-Jabir, A., Iosifidis, C., Agha, M. & Agha, R. (2020). The socio-economic implications of the coronavirus pandemic (COVID-19): A review. *International Journal of Surgery*, 78, 185-193.

Noguera Z., L. P., Kappel, C., Uhart, M. M., Diaz, F., Cayol, C., Cox-Witton, K., Death, C., Joly, D. O., Brown, K., Gardner, E. G., Suwanpakdee, S., Bett, B., Pepin, K. M., Yellowbird, K., Saulo, D., Morgan, O., Olson, S. H., & Pruvot, M. (2024). Theory of change for building stronger wildlife health surveillance systems globally. *Preprints*. <https://doi.org/10.20944/preprints202407.1055.v2>

One Health High-Level Expert Panel (OHHLEP), Adisasmito, W.B., Almuhairi, S., Behraves, C.B., Bilivogui, P., Bukachi, S.A., Casas, N., Cediell Becerra, N., Charron, D.F., Chaudhary, A., Ciacchi Zanella, J.R., Cunningham, A.A., Dar, O., Debnath, N., Dungu, B., Farag, E., Gao, G.F., Hayman, D.T.S., Khaita, M., Koopmans, M.P.G., Machalaba, C., Mackenzie,

400 J.S., Markotter, W., Mettenleiter, T.C., Morand, S., Smolenskiy, V. & Zhou, L. (2022). One  
 401 Health: A new definition for a sustainable and healthy future. *PLoS Pathog.*, 18, e1010537.  
 402 Orozco, M.M., Argibay, H.D., Minatel, L., Guillemi, E.C., Berra, Y., Schapira, A., Di Nucci, D.,  
 403 Marcos, A., Lois, F., Falzone, M. & Farber, M.D. (2020). A participatory surveillance of  
 404 marsh deer (*Blastocerus dichotomus*) morbidity and mortality in Argentina: first results.  
 405 *BMC Vet. Res.*, 16, 321.  
 406 Plowright, R.K., Reaser, J.K., Locke, H., Woodley, S.J., Patz, J.A., Becker, D.J., Oppler, G.,  
 407 Hudson, P.J. & Tabor, G.M. (2021). Land use-induced spillover: a call to action to  
 408 safeguard environmental, animal, and human health. *Lancet Planet Health*, 5, e237–e245.  
 409 Porco, A., Chea, S., Sours, S., Nou, V., Groenenberg, M., Agger, C., Tum, S., Chhuon, V., Sorn,  
 410 S., Hong, C., Davis, B., Davis, S., Ken, S., Olson, S. & Fine, A. (2023). Case Report:  
 411 Lumpy skin disease in an endangered wild banteng (*Bos javanicus*) and initiation of a  
 412 vaccination campaign in domestic livestock in Cambodia. *Frontiers in Veterinary Science*,  
 413 10.  
 414 Pruvot, M., Denstedt, E., Latinne, A., Porco, A., Montecino-Latorre, D., Khammavong, K.,  
 415 Milavong, P., Phouangsouvanh, S., Sisavanh, M., Nga, N.T.T., Ngoc, P.T.B., Thanh, V.D.,  
 416 Chea, S., Sours, S., Phommachanh, P., Theppangna, W., Phiphakhavong, S., Vanna, C.,  
 417 Masphal, K., Sothyra, T., San, S., Chamnan, H., Long, P.T., Diep, N.T., Duoc, V.T.,  
 418 Zimmer, P., Brown, K., Olson, S.H. & Fine, A.E. (2023). WildHealthNet: Supporting the  
 419 development of sustainable wildlife health surveillance networks in Southeast Asia. *Sci.*  
 420 *Total Environ.*, 863, 160748.  
 421 Reaser, J.K., Chitale, R.A., Tabor, G.M., Hudson, P.J. & Plowright, R.K. (2023). Looking Left:  
 422 Ecologically Based Biosecurity to Prevent Pandemics. *Health Secur.*

423 Rouquet P et al. 2005. Wild animal mortality monitoring and human Ebola outbreaks, Gabon and  
 424 Republic of Congo, 2001-2003. *Emerging infectious diseases* **11**:283–290. Centers for  
 425 Disease Control and Prevention (CDC).  
 426 Ryser-Degiorgis, M.-P. (2013). Wildlife health investigations: needs, challenges and  
 427 recommendations. *BMC Vet. Res.*, 9, 223.  
 428 Singh, R., Galliers, C., Moreto, W., Slade, J., Long, B., Aisha, H., Wright, A., Cartwright, F.,  
 429 Deokar, A., Wyatt, A., Deokar, D., Phoonjampa, R., Smallwood, E., Aziz, R., Koutoua  
 430 Benoit, A., Cao, R., Willmore, S., Jayantha, D. & Gosh, S. (2021). Impact of the COVID-19  
 431 pandemic on rangers and the role of rangers as a planetary health service. *Parks &*  
 432 *Recreation*, 119–134.  
 433 Sleeman, J.M., Brand, C.J. & Wright, S.D. (2012). Strategies for Wildlife Disease Surveillance.  
 434 USGS Staff -- Published Research.  
 435 Stephen, C. (2018). Evolving Urban Wildlife Health Surveillance to Intelligence for Pest  
 436 Mitigation and Monitoring. *Frontiers in Ecology and Evolution*, 6.  
 437 Stephen, C. & Berezowski, J. (2022). Wildlife Health Surveillance and Intelligence. Challenges  
 438 and Opportunities. In: *Wildlife Population Health* (ed. Stephen, C.). Springer International  
 439 Publishing, Cham, pp. 99–111.  
 440 Stolton, S., Timmins, H.L., Dudley, N., Biegus, O., Galliers, C., Jackson, W., Kettunen, M.,  
 441 Long, B., Rao, M., Rodriguez, C.M., Romanelli, C., Schneider, T., Seidl, A., Singh, R. &  
 442 Sykes, M. (2023). Essential planetary health workers: Positioning rangers within global  
 443 policy. *Conserv. Lett.*  
 444 du Toit, J.T. (2011). Ecology. Coexisting with cattle. *Science*.  
 445 Vicente, J., Vercauteren, K.C. & Gortázar, C. (2021). *Diseases at the Wildlife - Livestock*



446        *Interface: Research and Perspectives in a Changing World*. Springer Nature.

447    Vila, A.R., Briceño, C., McAloose, D., Seimon, T.A., Armien, A.G., Mauldin, E.A., Be, N.A.,

448        Thissen, J.B., Hinojosa, A., Quezada, M., Paredes, J., Avendaño, I., Silva, A. & Uhart,

449        M.M. (2019). Putative parapoxvirus-associated foot disease in the endangered huemul deer

450        (*Hippocamelus bisulcus*) in Bernardo O'Higgins National Park, Chile. *PLoS One*, 14,

451        e0213667.

452    Whitfield J. 2003. Ape populations decimated by hunting and Ebola virus. *Nature* 422, 551.

453    Wolf, T.M., Annie Wang, W., Lonsdorf, E.V., Gillespie, T.R., Pusey, A., Gilby, I.C., Travis,

454        D.A. & Singer, R.S. (2019). Optimizing syndromic health surveillance in free ranging great

455        apes: the case of Gombe National Park. *J. Appl. Ecol.*, 56, 509–518.

456    World Health Organization, Food and Agriculture Organization of the United Nations, United

457        Nations Environment Programme & World Organisation for Animal Health. (2023). *A*

458        *guide to implementing the One Health Joint Plan of Action at national level*.

459    World Organisation for Animal Health. (2010). *Training Manual on Wildlife Diseases and*

460        *Surveillance*. World Organisation for Animal Health.

461    World Organisation for Animal Health. (2015). *Training Manual on Surveillance and*

462        *International Reporting of Diseases in Wild Animals*. World Organisation for Animal

463        Health.

464    World Organisation for Animal Health. (2018). *Training Manual on Wildlife Health Information*

465        *Management - Fifth Cycle*. World Organisation for Animal Health.

466    World Organization for Animal Health. (2023). *In-country Wildlife Disease Surveillance Survey*

467        *Report*.

468    Worsley-Tonks, K.E.L., Bender, J.B., Deem, S.L., Ferguson, A.W., Fèvre, E.M., Martins, D.J.,

469 Muloi, D.M., Murray, S., Mutinda, M., Ogada, D., Omondi, G.P., Prasad, S., Wild, H.,  
470 Zimmerman, D.M. & Hassell, J.M. (2022). Strengthening global health security by  
471 improving disease surveillance in remote rural areas of low-income and middle-income  
472 countries. *Lancet Glob Health*, 10, e579–e584.

## Tables and Figures

*Table 1. Distribution of the method of documentation to register either healthy, sick, injured, or dead wildlife found during ranger patrols reported by local protected area data managers (“Individual observation”, “Part of the full count”, “Present or absent”, “Another way”) and the recording of specific data items for each wildlife health status across documentation methods.*



\* The black line shows the 50% reference.





Figure 1. Distribution of the level of agreement (grey scale) among local protected area data managers with statements: 'Pathogens carried by wildlife inhabiting the protected area(s) where I work in can affect livestock health' (brown), 'Pathogens carried by wildlife inhabiting the protected area(s) where I work in can affect human health' (red), 'Human or livestock pathogens can affect wildlife populations inhabiting the protected area(s) where I work in' (blue), and 'Wildlife health is important to achieve the conservation goals of the protected area(s) where I work' (green).



473 \* The overall encountering frequency for sick and injured wildlife was requested in a unique question, therefore, rows  
 474 one and two show the same total number of responses per encountering frequency.

*Figure 2. Distribution of local protected area data manager responses regarding the encounter of or injured (brown), sick (red), dead (blue) wildlife in the protected area(s) where they work and the recording (bright color) or non-recording (pale color) of these animals when encountered. Green bars represent the proportion of responses that reported the recording of wildlife per wildlife category and encounter frequency.*




Figure 3. Distribution of the level of agreement among local protected area data managers with the statement: ‘Introduced domestic animals (e.g., dogs, cats, cattle, pigs, cows) are a concern for the conservation goals of the protected areas where I work’ for the groups that reported the absence (red) and presence (blue) of domestic animals in the protected area(s) and their recording of the latter. Darker segments in the bars within the “Domestic animal in protected area > Yes response” category (x-axis) represent the number of responses that documented health status as part recording the presence of domestic animals.