

Wildlife health perceptions and monitoring practices in globally distributed protected areas

Diego Montecino-Latorre 1 Mathieu Pruvot 1,2
Sarah H Olson 1

1 Wildlife Conservation Society - Health Program, Bronx, NY, USA

2 University of Calgary, Faculty of Veterinary Medicine, Calgary, Alberta, Canada

Corresponding author: Diego Montecino-Latorre **email:** dmontecino@wcs.org

Keywords: conservation, protected areas, wildlife disease, wildlife health, wildlife health monitoring, wildlife health surveillance

Abstract

The status of health monitoring practices in protected areas (PAs) is largely unknown but potential gaps could undermine biodiversity conservation at these key sites. There is also a lack of baseline information regarding local perceptions of wildlife, human, and livestock health relevance that could affect health monitoring implementation in PAs. To address these deficiencies, we conducted a web-based survey of data managers from PAs worldwide. Specifically, we assessed perceptions regarding wildlife health and pathogen transmission between wildlife, humans, and livestock; the detection and documentation of unhealthy wildlife (injured, sick, and dead) and domestic animals in PAs; and health data management. Eighty-six out of 128 responses were analyzed. Respondents considered WH relevant to the conservation goals of PAs (97%), and 98% of them confirmed that unhealthy wildlife are encountered. However, >50% and >20% of respondents claimed that injured or sick and dead animals were not recorded, respectively. When these animals were documented, the recording methods and information collected differed. Although respondents considered domestic animal presence common and a conservation concern, these animals or their health status may not be recorded (30% and 74%, respectively). Health data were often stored in a database, but paper forms and spreadsheets were also used. Responses suggested that valuable syndromic wildlife health surveillance data from PAs are not collected or lost due to inadequate management and their value could be limited by a lack of standardized recording protocols.

Introduction

Protected areas (PAs) safeguard intact landscapes, conserving diverse wildlife and flora, and preserve essential ecosystem services while supporting local communities. However, they are increasingly threatened by anthropogenic pressures. Human encroachment and land-use change (Laurance et al. 2012; Vicente et al. 2021; Meng et al. 2023) drive resource extraction, pollution, creation of human-wildlife-livestock interfaces, and ecosystem degradation (Plowright et al. 2021; Vicente et al. 2021; Reaser et al. 2023). These processes expose wildlife to physical (e.g., snaring), chemical (e.g., toxins), and biological (e.g., pathogens) hazards and can negatively affect biodiversity conservation and global health (De Vos et al. 2016; Wolf et al. 2019; Machalaba et al. 2020; Hacon et al. 2020; Porco et al. 2023; Groenenberg et al. 2023).

Wildlife health (WH) monitoring, which involves tracking diseases, pathogens, and toxic agents in wild animal populations (World Organisation for Animal Health & International Union Conservation of Nature 2024) facilitates early detection of exposure and disease, enables rapid response to mitigate risks and adverse outcomes, and supports the evaluation of health management strategies (Woods et al. 2019; Machalaba et al. 2021; One Health High-Level Expert Panel [OHHLEP] et al. 2022; Porco et al. 2023; Elnaïem et al. 2023; Vora et al. 2023; Thompson et al. 2024). The One Health approach recognizes that the health of animals and humans and the functioning of their shared environment are deeply interconnected (Machalaba et al. 2021; OHHLEP et al. 2022; World Organization for Animal Health 2023). As a result, WH monitoring is essential for biodiversity conservation and for safeguarding human and animal health.

Ebola virus disease is an illustrative example. Outbreaks in wildlife have decimated populations of western gorillas and chimpanzees (Whitfield 2003; Leroy et al. 2004; Bermejo et al. 2006). Index cases of outbreaks of this disease in humans have involved hunters who extracted animals from the wild, including in PAs (Judson et al. 2016). Virus spread into urban areas of West Africa through human-to-human transmission in 2014 resulted in over 20,000 excess human deaths (Dudas et al. 2017; Jacob et al. 2020).

Despite its importance, WH monitoring systems remain uncommon or deficient (Machalaba et al. 2021; OHHLEP et al. 2022; World Organization for Animal Health 2023). To our knowledge, the status of WH monitoring practices in PAs is largely unknown. A lack of such monitoring or gaps could undermine biodiversity conservation and compromise One Health, highlighting the need for urgent attention. Additionally, baseline information regarding the perception of wildlife, human, and livestock health relevance for biodiversity conservation by PA personnel is also unknown. Their perceptions could impact affect WH monitoring practices and their implementation.

To address knowledge gaps associated with current WH monitoring practices in PAs and associated perceptions by personnel, we surveyed protected area data managers (PADMs) to assess their perceptions regarding WH and pathogen transmission between wildlife, humans,

and livestock; the detection and documentation of injured, sick, or dead wildlife and domestic animals in PAs; and health data management in PAs.

Methods

We developed a web-based questionnaire aimed at PADM users of the Spatial Monitoring and Reporting Tool (SMART). SMART is a suite of technology tools an online that support the administration of PAs (Cronin et al. 2021) and is used in more than 1,000 conservation sites worldwide. Thus, its user community offers the opportunity to engage a large number of PADM. The survey was distributed to the SMART Community Forum users by the SMART Partnership (<https://smartconservationtools.org>) via email in October 2022. The survey remained open for three months. A reminder was sent three weeks before the closing date.

Survey questions are in Appendix S1. Because the SMART Community Forum includes many conservation actors, respondents were asked first if their job roles and responsibilities matched our definition of PADM: “a person directly responsible for managing SMART data in one or more PAs or a general manager or administrator of one or more PAs that uses SMART data.” Respondents who did not identify as a PADM were excluded. The survey had four sections. Section 1 assessed the perception of PADM on the importance of WH in achieving conservation goals, the role of human and livestock pathogens in affecting WH, and the role of wildlife pathogens in affecting public and livestock health. Section 2 asked respondents to rank the overall frequency of encounters with injured, sick, or dead wildlife in PAs and to indicate whether animals in each of these classes, along with healthy wildlife, are recorded as a specific type of individual when found during patrols. Section 3, asked about the presence of domestic animals in the PAs, the documentation of their health status, and the perceived threats of domestic animals to conservation goals. In sections 1-3 Likert scales were used for responses. Section 4 addressed health data storage practices when collected.

An introductory web page explained that the survey was voluntary, anonymous, and aimed at PADM and that clicking the “start the survey” button constituted consent. A tutorial was provided for the language-translation tool of this survey built on Google Forms. No personal information was collected, so the survey was exempt from full ethics review by the Wildlife Conservation Society’s IRB (REF# 22-53).

Responses by PADM could represent either a single or multiple PAs. For our analysis, we focused on what we defined as local” responses, which included one or two PAs. “Non-local responses”, which covered more than two PAs, were analyzed separately. We assumed that PADM s responsible for more than two PAs would conduct occasional site visits and that it was unlikely they would have insights into specific PA realities (e.g., patrol findings, non-recorded data, data management, etc). However, they likely had an understanding of perceptions at central offices at the decision-making level. In contrast, we assumed “local” PADM would work on-site and understand PA realities. The two-PA cutoff was chosen to balance maximization of

the local respondent sample size with accurate classification of them. We discarded responses that pertained only to marine PAs (determined based on the World Database on Protected Areas; <https://www.protectedplanet.net/en/thematic-areas/wdpa?tab=WDPA>) because marine PA management, species, and patrol logistics are markedly different from those in terrestrial PAs. The descriptive analysis of survey responses was conducted in R 4.3.1. The responses dataset is available from <https://figshare.com/s/36513db82ac5dfa8e71d?file=49682265>. Descriptive analyses can be found at https://github.com/dmontecino/SMART_survey.

Results

We received 128 responses. Forty-two were removed because either the PA name or names were not provided, the PAs were marine, or respondents did not match the target audience (8, 7, and 27, respectively). The final dataset contained 86 respondents from 23 countries. Seventy-three were local responses, 67 surveys pertained to a single PA, and 6 pertained to 2 PAs. Local responses came from 19 countries. There were 13 non-local responses from 10 countries. The names of the countries are not provided to protect the identity of the respondents, but local responses were from North, Central, and South America (n=45); West, Central, East, and Southern Africa (n=16); Southeast and Southern Asia (n=11); and Europe (n=1). Most responses came from South America. Non-local responses were from West, Central, and East Africa (n=6), Central and South America (n=3), and Southeast and Southern Asia (n=4).

Perceptions of wildlife health importance

Most respondents either strongly agreed or agreed with the statements “Wildlife health, including infectious and non-infectious diseases, is important to achieve the conservation goals of the protected areas where I work.” and “Human or livestock pathogens can affect wildlife populations inhabiting the protected area(s) I work in.” (92% and 81%, respectively). Regarding the affirmation “Pathogens carried by wildlife inhabiting the protected area(s) where I work in can affect livestock health.”, 48% of respondents strongly agreed or agreed and 29% neither agreed nor disagreed (e.g., were neutral). Across respondents, 63% strongly agreed or agreed that “Pathogens carried by wildlife inhabiting the protected area(s) where I work in can affect human health.” Neutral responses were 19%. Detailed distributions are shown in Figure 1. Non-local responses followed similar trends; neutral responses were proportionally fewer (Appendix S2).




Figure 1: Level of agreement (grey scale) among local protected area data managers with the statements “Pathogens carried by wildlife inhabiting the protected area(s) where I work in can affect livestock health.” (brown), “Pathogens carried by wildlife inhabiting the protected area(s) where I work in can affect human health.” (red), “Human or livestock pathogens can affect wildlife populations inhabiting the protected area(s) where I work in.” (blue), and “Wildlife health is important to achieve the conservation goals of the protected area(s) where I work,” (green).

Encounters with injured, sick, or dead wildlife and documentation

Most local overall frequency-of-encounter responses concentrated between “very rarely” and “occasionally” (Figure 2). Seventy-six percent of local PADM reporting the encounter of dead animals in the PAs (e.g., responded “very rarely” or more frequently) indicated that these encounters were documented. Forty-nine percent of local PADM reporting encounters with injured or sick animals and 35% reporting encounters with sick animals in the PAs confirmed their documentation. In general, the documentation of injured, sick, or dead animals tended to be higher as the encounter frequency increased from “very rarely” to “very frequently”. For example, the percentage of local responses reporting the documentation of sick animals

as “very rarely” was <20% versus >50% for the encounter frequency “very frequently” (Figure 2). All non-local PADM reported the encounter with injured or sick wildlife and dead wildlife. The percentages of non-local PADM reporting the documentation of these animals were larger compared with local responses (85%, 62%, and 92% for injured, sick, and dead wildlife respectively; Appendix S3).




Figure 2: Number of local protected area data managers who reported that the health status of and frequency of encounters with wildlife are recorded or not recorded in the protected area they worked in. Green bars represent the proportion of respondents that reported recording of wildlife in each category. Overall frequency of encounters with sick and injured wildlife was requested in a unique question; therefore, rows one and two show the same total number of responses per encounter category.

All 17 local PADM who indicated encountering dead wildlife “very rarely” or more frequently but answered that these animals were not documented, either agreed or strongly agreed with the importance of WH to achieve conservation goals. Similarly, 94% of local PADM who indicated encountering injured wildlife “very rarely” or more frequently but answered that these animals were not documented, either agreed or strongly agreed with this statement. For sick wildlife, the percentage was 91%. The corresponding percentages for non-local PADM were 50%, 80%, and 0% for injured, sick, and dead wildlife, respectively.

Documentation methods

The documentation method of injured, sick, or dead wildlife varied among the 58 local PADMIs that reported the recording of one or more of these groups. Most often, each animal was documented individually (“individual observation”). The second most common method was a complete inventory of healthy, injured, sick, or dead animals for each species. Reporting their presence or absence was the third most common method (Figure 3). For non-local responses, the predominant method was “each animal is an individual observation” across health categories (Appendix S4).

The items recorded from each observation were not consistent across responses (Figure 3). Photographs and the species were the main items collected across documentation methods and health categories. Anomalies observed in unhealthy wildlife (e.g., sick or dead) and the condition of carcasses were not always recorded (Figure 3). In non-local responses the trend was relatively similar; however, items were reported as being recorded more consistently (e.g., age, anomalies, and condition in the three health categories; Appendix S4).




Figure 3: Distribution of the methods of documentation (second column) of healthy, sick, injured, or dead wildlife found during ranger patrols as reported by local protected area data managers and the recording of specific types of data for each wildlife health status across documentation methods (black line, 50%).

Domestic animals in protected areas

Fifty-two local PADMs (71%) responded that domestic animals were found in their PAs. Among them, 67% reported that domestic animals were documented if observed during patrols, and 26% reported recording of their health status (Figure 4). Forty-two local respondents reporting domestic animals in the PAs (81%) either agreed or strongly agreed that domestic animals are a conservation concern (Figure 4). Twenty-seven of them (64%) answered that these animals were documented. Fourteen out of 21 respondents claimed that domestic animals are not found in the corresponding PAs and either agreed or strongly agreed that they are a conservation concern (67%).

Eight non-local PADMs (62%) responded that domestic animals were found in the PAs. Among them, seven (88%) reported that domestic animals were documented if observed during patrols, of which only two (29%) reported recording their health status (Appendix S5).




Figure 4: Responses of local protected area data manager to statements that the presence and health of domestic animals in protected areas is a conservation concern (red, no recording of domestic animals; light blue, recording of domestic animals but not their health status; dark blue, recording of domestic animals and their health status). Data are from the group of protected area data managers who reported the presence of domestic animals in the protected area.

Health data storage practices

Seventy-three percent, 54%, and 65% of local PADMs reported the documentation of injured, sick, or dead wildlife being stored in a SMART database. Paper forms, reports, and spreadsheets were employed when unhealthy wildlife were documented, but their data were not stored in a SMART database.

Thirty-one of the 35 local respondents (100%) recording domestic animals during patrols indicated that this information was stored in a SMART database. The health status of domestic animals, when documented, was also stored in a SMART database.

Discussion

Our findings suggest that syndromic WH surveillance data are not collected or lost due to inadequate management. Even when WH data are collected and properly managed, the usefulness for surveillance is likely limited by the diversity of methods employed to record them. By the time of the survey, an initiative to foster the harmonized recording of WH data in PAs (Montecino-Latorre et al. 2024 [preprint]) had been implemented in a couple of sites only in a single country, and it did not affect overall results involving multiple PAs and countries.

The PADM s largely considered WH relevant to conservation goals of PAs. However, several local PADM s expressed neutral views on the potential impact of wildlife pathogens on human and livestock health. Local PADM s and field staff could be exposed to zoonotic pathogens (Adjemian et al. 2012). It is essential to provide rangers with training in biosecurity measures to mitigate these risks.

Most PADM s confirmed that unhealthy wildlife were encountered. However, the percentage of local PADM s who agreed or strongly agreed that WH is relevant for the conservation goals of PAs was higher than the percentage of those reporting the recording of unhealthy wildlife. In other words, the perceived importance of WH did not translate into reporting. A similar contradiction was also observed in the responses specifically reporting the encounter of domestic animals but not their documentation. This discrepancy could be explained by recent global pathogen-driven crises, such as SARS-CoV-2 and H5N1 highly pathogenic avian influenza virus (Nicola et al. 2020; Leguia et al, 2023). It might also be explained by other morbidity or mortality events of different etiology that our audience was sensitized to at the time of the survey but before health-associated monitoring objectives could be planned and implemented. These findings could also suggest a lack of knowledge or resources to act on their understanding of the importance of WH for conservation goals.

More non-local respondents reported the documentation of non-healthy wildlife than local respondents. This difference could be explained by non-local and local responses coming from different PAs or it could represent differences between the expectations of managers in an administrative role (non-local PADM s) and field realities in PAs (local PADM s). For example, the mandate to record non-healthy animals may exist, but it is not pursued in practice because of limitations regarding WH training, the use of a recording tool, comprehension of the methodology to record these data, or willingness to record these data (Wilfred et al. 2019; Kavhu & Mpakairi 2021). If any of these cases exist, PA management agencies should take a more active local role in identifying and correcting weaknesses in WH data collection.

We noted a general agreement among PADM s regarding the conservation threat domestic animals (e.g., dogs, cats, cattle) present. Although we did not explicitly ask why domestic animals are a conservation concern, most PADM s also agreed with the statement “Human and livestock pathogens can affect wildlife populations...” whether these animals were found in the PAs or not (Appendices S6 and S7). Pathogen transmission from domestic animals to wild animals can seriously harm biodiversity conservation, including in PAs (Porco et al. 2023), and they add to the pressures on wildlife from domestic animals in- and outside PAs (e.g., predation, competition [du Toit 2011; Gompper 2013]). The observed contradiction between perceived conservation risk of domestic animals and documentation of their presence and health status could be explained by the same drivers mentioned above.

Effective management of data and harmonization are foundational pillars for WH monitoring (World Organisation for Animal Health 2010, 2015, 2018; Sleeman et al. 2012; Ryser-Degiorgis 2013; Stephen 2018; Machalaba et al. 2021; Stephen & Berezowski 2022; Giacinti et al. 2022; Hayman et al. 2023; Heiderich et al. 2023). However, we identified challenges in WH data harmonization and governance that align with historical pitfalls in WH surveillance (Carmichael

2012; Cardoso et al. 2021; World Organization for Animal Health 2023; Heiderich et al. 2023; Suwanpakdee et al. 2024). When healthy and unhealthy wildlife have been documented, inconsistencies in data structure and variables were present. The lack of harmonization across PAs, within and beyond country borders, diminishes the value of collected health data, making transboundary health assessments challenging. Similarly, tracking WH trends over time becomes infeasible. Additionally, records of unhealthy wildlife could be stored in paper forms or Excel sheets rather than a SMART database. Data in paper forms or Excel sheets can be uploaded to a SMART database manually by filling specific fields or through a mobile application (SMART Mobile) designed to capture data as determined in the corresponding SMART database (Cronin et al. 2021). Consequently, WH data in paper forms and spreadsheets reflect that they were not considered in the PAs' SMART databases. We did not ask about the use of SMART Mobile, but challenges in its adoption have been described (Wilfred et al. 2019; Kavhu & Mpakairi 2021; Wyatt et al. 2023).

This is the first account of WH perceptions and monitoring practices in PAs. Because survey respondents were contacted through the SMART Partnership, they may not fully represent the broader population of local and non-local PADMs. Ideally, a follow-up longitudinal study and respondents selected through random sampling from a sampling frame, including local and non-local PADMs of the same PAs, could further refine and enhance our initial insights. This approach could also identify geographic differences and associations between distance patrolled per time unit, landscape type, and fauna size in the overall encounter rates with health and unhealthy wildlife and domestic animals. Potential differences in reporting as a consequence of experience with the recording tool used (SMART Mobile or other) should also be assessed.

Leveraging existing individuals who can detect morbidity and mortality in wildlife offers a cost-effective strategy to establish minimal WH monitoring. Rangers can detect non-healthy animals in PAs, and they have provided data to assess health risks and trends or trigger responses to disease outbreaks (Wolf et al. 2019; Vila et al. 2019; Kuisma et al. 2019; Orozco et al. 2020; Montecino-Latorre et al. 2020; Porco et al. 2023).

Our recommendation is to include standardized WH monitoring in the remit of rangers following a unified methodology and standards. Currently, there are approximately 280,000 rangers worldwide, and 1.5 million will be needed to protect 30% of the planet by 2030 (Appleton et al. 2022). The present and projected number of rangers needed reveal their unique potential as a worldwide One Health workforce that could drastically improve WH and One Health surveillance (Machalaba et al. 2021; Hopkins et al. 2024; Montecino-Latorre et al. 2024 [preprint]; World Organisation for Animal Health & International Union Conservation of Nature 2024). However, our findings suggest that several problems must be addressed first.

Acknowledgments

Thanks to all responders to our survey and the SMART Partnership We thank responders and the SMART Partnership, especially to M. Barcellos, and J. Palmer, Executive Director of the Wildlife Conservation Society's Conservation Technology Department.

References

- Adjemian J et al. 2012. Zoonotic infections among employees from Great Smoky Mountains and Rocky Mountain National Parks, 2008-2009. *Vector Borne and Zoonotic Diseases* 12:922–931.
- Appleton MR et al. 2022. Protected area personnel and ranger numbers are insufficient to deliver global expectations. *Nature Sustainability* 5:1100–1110.
- Bermejo M, Rodríguez-Teijeiro JD, Illera G, Barroso A, Vilà C, Walsh PD. 2006. Ebola outbreak killed 5000 gorillas. *Science* 314:1564.
- Cardoso B, García-Bocanegra I, Acevedo P, Cáceres G, Alves PC, Gortázar C. 2021. Stepping up from wildlife disease surveillance to integrated wildlife monitoring in Europe. *Research in Veterinary Science* 144:149-156.
- Carmichael C. 2012. Coordinating an effective response to wildlife diseases. *Wildlife Society Bulletin* 36:204–206.
- Cronin DT, Dancer A, Long B, Lynam AJ, Muntifering J, Palmer J, Bergl RA, Wich SA, Piel AK. 2021. Application of SMART software for conservation area management. Pages 201–224 *Conservation Technology*. Oxford University Press Oxford, UK.
- De Vos A, Cumming G, Cumming D, Ament J, Baum J, Clements H, Grewar J, Maciejewski K, Moore C. 2016. Pathogens, disease, and the social-ecological resilience of protected areas. *Ecology and Society* 21.
- Dudas G et al. 2017. Virus genomes reveal factors that spread and sustained the Ebola epidemic. *Nature* 544:309–315.
- du Toit JT. 2011. Coexisting with cattle. *Science* 333:1710–1711. Elnaïem A et al. 2023. Global and regional governance of One Health and implications for global health security. *The Lancet* 401:688–704.
- Giacinti JA, Jane Parmley E, Reist M, Bayley D, Pearl DL, Jardine CM. 2022. Canadian wildlife health surveillance—patterns, challenges and opportunities identified by a scoping review. *Facets* 7:25–44.
- Gompper ME. 2013. *Free-Ranging Dogs and Wildlife Conservation*. Oxford Univ. Press. UK.

- Groenenberg M et al. 2023. Snaring devastates terrestrial ungulates whilst sparing arboreal primates in Cambodia's Eastern Plains Landscape. *Biological Conservation* 284:110195.
- Hacon S de S, Oliveira-da-Costa M, Gama C de S, Ferreira R, Basta PC, Schramm A, Yokota D. 2020. Mercury exposure through fish consumption in traditional communities in the Brazilian northern Amazon. *International Journal of Environmental Research and Public Health* 17: 5269.
- Hayman DTS et al. 2023. Developing One Health surveillance systems. *One Health*:100617.
- Heiderich E et al. 2023, July 13. Analysis of a European general wildlife health surveillance program: chances, challenges and recommendations. *Plos One* 19: e0301438.
- Hopkins SR et al. 2024. Protected areas and One Health. *Parks* 30:6–13.
- Jacob ST et al. 2020. Ebola virus disease. *Nature Reviews Disease Primers* 6:13.
- Judson SD, Fischer R, Judson A, Munster VJ. 2016. Ecological contexts of index cases and spillover events of different Ebolaviruses. *PLoS Pathogens* 12:e1005780.
- Kavhu B, Mpakairi KS. 2021. Spatial Monitoring and Reporting Tool (SMART) in Mid-Zambezi Valley, Zimbabwe: Implementation challenges and practices. *Conservation Science and Practice* 3: e492.
- Kuisma E et al. 2019. Long-term wildlife mortality surveillance in northern Congo: a model for the detection of Ebola virus disease epizootics. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* 374:20180339.
- Laurance WF et al. 2012. Averting biodiversity collapse in tropical forest protected areas. *Nature* 489:290–294.
- Leguia M et al. 2023. Highly pathogenic avian influenza A (H5N1) in marine mammals and seabirds in Peru. *Nature Communications* 14, 5489. <https://doi.org/10.1038/s41467-023-41182-0>
- Leroy EM et al. 2004. Multiple Ebola virus transmission events and rapid decline of central African wildlife. *Science* 303:387–390.
- Machalaba C, Feferholtz Y, Uhart M, Karesh WB. 2020. Wildlife conservation status and disease trends: ten years of reports to the Worldwide Monitoring System for Wild Animal Diseases. *Revue Scientifique et Technique* 39:991–1001.
- Machalaba C, Uhart M, Ryser-Degiorgis M-P, Karesh WB. 2021. Gaps in health security related to wildlife and environment affecting pandemic prevention and preparedness, 2007-2020. *Bulletin of the World Health Organization* 99.
- Meng Z, Dong J, Ellis EC, Metternicht G, Qin Y, Song X-P, Löfqvist S, Garrett RD, Jia X, Xiao X. 2023. Post-2020 biodiversity framework challenged by cropland expansion in protected areas. *Nature Sustainability* 6: 758-768.

- Montecino-Latorre D et al. 2024. Rangers as an essential One Health workforce: Strengthening the preservation of biodiversity and global health in practice. Preprint. SSRN. <http://dx.doi.org/10.2139/ssrn.4997409>.
- Montecino-Latorre D, Napolitano C, Briceño C, Uhart MM. 2020. Sarcoptic mange: An emerging threat to Chilean wild mammals? *Perspectives in Ecology and Conservation* 18:267–276.
- Nicola M, Alsafi Z, Sohrabi C, Kerwan A, Al-Jabir A, Iosifidis C, Agha M, Agha R. 2020. The socio-economic implications of the coronavirus pandemic (COVID-19): A review. *International Journal of Surgery* 78:185–193. 10.1016/j.ijssu.2020.04.018
- One Health High-Level Expert Panel et al. 2022. One Health: A new definition for a sustainable and healthy future. *PLoS Pathogens* 18:e1010537.
- Orozco MM et al. 2020. A participatory surveillance of marsh deer (*Blastocerus dichotomus*) morbidity and mortality in Argentina: first results. *BMC Veterinary Research* 16:321.
- Plowright RK, Reaser JK, Locke H, Woodley SJ, Patz JA, Becker DJ, Oppler G, Hudson PJ, Tabor GM. 2021. Land use-induced spillover: a call to action to safeguard environmental, animal, and human health. *The Lancet Planetary Health* 5:e237–e245.
- Porco A et al. 2023. Case Report: Lumpy skin disease in an endangered wild banteng (*Bos javanicus*) and initiation of a vaccination campaign in domestic livestock in Cambodia. *Frontiers in Veterinary Science* 10.
- Reaser JK, Chitale RA, Tabor GM, Hudson PJ, Plowright RK. 2023. Looking left: ecologically based biosecurity to prevent pandemics. *Health security* 22: 74-81.
- Ryser-Degiorgis M-P. 2013. Wildlife health investigations: needs, challenges and recommendations. *BMC Veterinary Research* 9:1-17.
- Sleeman JM, Brand CJ, Wright SD. 2012. Strategies for wildlife disease surveillance. In: A.A. Aguirre, R.S. Ostfeld, P. Daszak (Eds.), *New Directions in Conservation Medicine: Applied Cases in Ecological Health*. Oxford University Press. New York. 539-551.
- Stephen C. 2018. Evolving urban wildlife health surveillance to intelligence for pest mitigation and monitoring. *Frontiers in Ecology and Evolution* 6. <https://doi.org/10.3389/fevo.2018.00127>
- Stephen C, Berezowski J. 2022. Wildlife Health Surveillance and Intelligence. Challenges and Opportunities. Pages 99–111 in Stephen C, editor. *Wildlife Population Health*. Springer.
- Suwanpakdee S et al. 2024. Wildlife health capacity enhancement in Thailand through the World Organisation for Animal Health Twinning Program. *Frontiers in Veterinary Science* 11:1462280.
- Thompson L, Cayol C, Awada L, Muset S, Shetty D, Wang J, Tizzani P. 2024. Role of the World Organisation for Animal Health in global wildlife disease surveillance. *Frontiers in Veterinary Science* 11.

- Vicente J, Vercauteren KC, Gortázar C. 2021. Diseases at the Wildlife - Livestock Interface: Research and Perspectives in a Changing World. Springer Nature.
- Vila AR et al. 2019. Putative parapoxvirus-associated foot disease in the endangered huemul deer (*Hippocamelus bisulcus*) in Bernardo O'Higgins National Park, Chile. *PloS One* 14:e0213667.
- Vora NM et al. 2023. Interventions to reduce risk for pathogen spillover and early disease spread to prevent outbreaks, epidemics, and pandemics. *Emerging Infectious Diseases* 29:1–9.
- Whitfield J. 2003. Ape populations decimated by hunting and Ebola virus. *Nature* 422:551. <https://doi.org/10.1038/422551a>
- Wilfred P, Kayeye H, Magige FJ, Kisingo A, Nahonyo CL. 2019. Challenges facing the introduction of SMART patrols in a game reserve, western Tanzania. *African Journal of Ecology* 57:523–530.
- Wolf TM, Annie Wang W, Lonsdorf EV, Gillespie TR, Pusey A, Gilby IC, Travis DA, Singer RS. 2019. Optimizing syndromic health surveillance in free ranging great apes: the case of Gombe National Park. *The Journal of Applied Ecology* 56:509–518.
- Woods R, Reiss A, Cox-Witton K, Grillo T, Peters A. 2019. The Importance of wildlife disease monitoring as part of global surveillance for zoonotic diseases: The role of Australia. *Tropical Medicine and Infectious Disease* 4. <https://doi.org/10.3390/tropicalmed4010029>
- World Organisation for Animal Health. 2010. Training manual on wildlife diseases and surveillance. Paris, World Organisation for Animal Health.
- World Organisation for Animal Health. 2015. Training manual on surveillance and international reporting of diseases in wild animals. Paris, World Organisation for Animal Health.
- World Organisation for Animal Health. 2018. Training manual on wildlife health information management. Paris, World Organisation for Animal Health.
- World Organisation for Animal Health, International Union Conservation of Nature (IUCN). 2024. General guidelines for surveillance of diseases, pathogens and toxic agents in free-ranging wildlife: An overview for wildlife authorities and others working with wildlife. Paris, World Organisation for Animal Health and Gland, Switzerland, IUCN.
- World Organization for Animal Health. 2023. In-country wildlife disease surveillance survey report. Paris, World Organisation for Animal Health.
- Wyatt A, Singh R, Read C. 2023. Assessing the adaptative management process in SMART sites: Lessons learned. *Parks* 29.

Supporting Information ‘Wildlife health perceptions and monitoring practices in globally distributed protected areas’

Methods

Appendix S1. Survey

Use of SMART to record wildlife health information in protected areas around the world

* Indicates required question

1. **1. Please enter all the name(s) of the protected area(s) that apply.** *

If there is more than one protected area(s) name(s), please separate them by hitting enter in the answer space below

Please read the six statements and mark how you feel about them (from “**strongly disagree**” to “**strongly agree**”).

2. **2. Wildlife health, including infectious and non-infectious diseases, is important to achieve the conservation goals of the protected areas where I work** *

Mark only one oval.

- ☐ Strongly disagree
- ☐ Disagree
- ☐ Somewhat Disagree
- ☐ Neutral
- ☐ Agree
- ☐ Strongly agree

3. **3. Pathogens carried by wildlife inhabiting the protected area I work in can affect livestock health** *

Mark only one oval.

- ☐ Strongly disagree
- ☐ Disagree
- ☐ Somewhat Disagree
- ☐ Neutral
- ☐ Agree
- ☐ Strongly agree

4. **4. Pathogens carried by wildlife inhabiting the protected area I work in can affect public health** *

Mark only one oval.

- ☐ Strongly disagree
- ☐ Disagree
- ☐ Somewhat Disagree
- ☐ Neutral
- ☐ Agree
- ☐ Strongly agree

5. **5. Human or livestock pathogens can affect wildlife populations inhabiting the protected area I work in** *

Mark only one oval.

- ☐ Strongly disagree
- ☐ Disagree
- ☐ Somewhat Disagree
- ☐ Neutral
- ☐ Agree
- ☐ Strongly agree

6. **6. Introduced domestic animals (e.g., dogs, cats, cattle, pigs, cats) are a concern for the conservation goals of the protected areas where I work** *

Mark only one oval.

- ☐ Strongly disagree
- ☐ Disagree
- ☐ Somewhat Disagree
- ☐ Neutral
- ☐ Agree
- ☐ Strongly agree

Please mark one of the alternatives provided to answer the following questions based on your experience

7. **7. Are dead wildlife encountered in the protected area?** *

Mark only one oval.

- ☐ Always
- ☐ Very frequently
- ☐ Occasionally
- ☐ Sometimes
- ☐ Rarely
- ☐ Very rarely
- ☐ Never

8. **8. Are sick or injured wildlife encountered in the protected area? ***

Mark only one oval.

- ☐ Always
- ☐ Very frequently
- ☐ Occasionally
- ☐ Sometimes
- ☐ Rarely
- ☐ Very rarely
- ☐ Never

9. **9. Are livestock encountered in the protected area? ***

Mark only one oval.

- ☐ Always
- ☐ Very frequently
- ☐ Occasionally
- ☐ Sometimes
- ☐ Rarely
- ☐ Very rarely
- ☐ Never

10. **10. How long, on average, are the patrols in the protected area? ***

Mark only one oval.

- ☐ Less than a day
- ☐ Between one to two days
- ☐ Between two and five days
- ☐ Between four days and a week

11. **11. On average, how many patrols are completed in the protected area in one month?** *
-

Please respond to the following questions related to data recorded in SMART (e.g., data collected by ranger patrols, community members, or other users and managed in SMART)

12. **12. Are healthy wildlife found during patrols (rangers or others) recorded as a specific category of individuals?** *

Mark only one oval.

☐ Yes *Skip to question 13*

☐ No *Skip to question 17*

13. **12.a Please select one of the choices below to help us understand how healthy wildlife are recorded during a patrol:** *

Mark only one oval.

☐ Healthy wildlife are recorded as present/absent (the number of healthy individuals is not specified)

☐ Healthy wildlife are counted and reported as part of the full count of individuals belonging to an observed species (e.g. "3 total healthy animals of species X and 2 total healthy animals of species Y")

☐ Each healthy animal is recorded as an individual observation

☐ Healthy wildlife is recorded in another way

14. 12.b Please indicate what type of data are recorded (check all that apply) *

Check all that apply.

- ☐ Photographs
- ☐ Species
- ☐ Age
- ☐ Sex
- ☐ Body condition
- ☐ Other

15. 12.c Are these data entered and stored in SMART Desktop?

Mark only one oval.

- ☐ All of these items are entered and stored in the corresponding SMART Conservation Area *Skip to question 17*
- ☐ Some of these items are entered and stored in the corresponding SMART Conservation Area *Skip to question 17*
- ☐ None of these items are recorded in the corresponding SMART Conservation Area *Skip to question 16*

16. 12.c.i If none of the items are recorded in the corresponding SMART Conservation Area where are they recorded? *

17. 13. Are dead wildlife found during patrols recorded as a specific category of individuals? *

Mark only one oval.

- ☐ Yes *Skip to question 18*
- ☐ No *Skip to question 22*

18. **13.a Please select one of the choices below to help us understand how dead wildlife are recorded during a patrol:** *

Mark only one oval.

- ☐ Dead wildlife are recorded as present/absent (the number of dead individuals is not specified)
- ☐ Dead wildlife are counted and reported as part of the full count of individuals belonging to an observed species (e.g. "3 total dead animals of species X and 2 total dead animals of species Y").
- ☐ Each dead animal is recorded as an individual observation
- ☐ Dead wildlife is recorded in another way

19. **13.b Please indicate what type of data are recorded (check all that apply)** *

Check all that apply.

- ☐ Photographs
- ☐ Species
- ☐ Age
- ☐ Sex
- ☐ Carcass condition
- ☐ Anomalies in carcass (if any)
- ☐ Suspect cause of death
- ☐ Other

20. **13.c Are these data entered and stored in SMART Desktop?** *

Mark only one oval.

- ☐ All of these items are entered and stored in the corresponding SMART Conservation Area *Skip to question 22*
- ☐ Some of these items are entered and stored in the corresponding SMART Conservation Area *Skip to question 22*
- ☐ None of these items are recorded in the corresponding SMART Conservation Area *Skip to question 21*

21. **13.c.i If none of the items are recorded in the corresponding SMART Conservation Area where are they recorded?** *
-

22. **14. Are injured wildlife found during patrols recorded as a specific category of individuals?** *

Mark only one oval.

- ☐ Yes
- ☐ No *Skip to question 27*

23. **14.a Please select one of the choices below to help us understand how injured wildlife are recorded during a patrol:** *

Mark only one oval.

- ☐ Injured wildlife are recorded as present/absent (the number of injured individuals is not specified)
- ☐ Injured wildlife are counted and reported as part of the full count of individuals belonging to an observed species (e.g. "3 total injured animals of species X and 2 total injured animals of species Y").
- ☐ Each injured animal is recorded as an individual observation
- ☐ Injured wildlife is recorded in another way

24. **14.b Please indicate what type of data are recorded (check all that apply)** *

Check all that apply.

- ☐ Photographs
- ☐ Species
- ☐ Age
- ☐ Sex
- ☐ Anomalies/signs if any
- ☐ Body condition
- ☐ Suspect cause of injury
- ☐ Other

25. **14.c Are these data entered and stored in SMART Desktop? ***

Mark only one oval.

- ☐ All of these items are entered and stored in the corresponding SMART Conservation Area *Skip to question 27*
- ☐ Some of these items are entered and stored in the corresponding SMART Conservation Area *Skip to question 27*
- ☐ None of these items are recorded in the corresponding SMART Conservation Area *Skip to question 26*

26. **14.c.i If none of the items are recorded in the corresponding SMART Conservation Area where are they recorded? ***

27. **15. Are sick wildlife found during patrols recorded as a specific category of individuals? ***

Mark only one oval.

- ☐ Yes *Skip to question 28*
- ☐ No *Skip to question 32*

28. **15.a Please select one of the choices below to help us understand how sick wildlife are recorded during a patrol: ***

Mark only one oval.

- ☐ Sick wildlife are recorded as present/absent (the number of sick individuals is not specified)
- ☐ Sick wildlife are counted and reported as part of the full count of individuals belonging to an observed species (e.g. "3 total sick animals of species X and 2 total sick animals of species Y").
- ☐ Each sick animal is recorded as an individual observation
- ☐ Sick wildlife is recorded in another way

29. **15.b Please indicate what type of data are recorded (check all that apply) ***

Check all that apply.

- ☐ Photographs
- ☐ Species
- ☐ Age
- ☐ Sex
- ☐ Anomalies/signs if any
- ☐ Body condition
- ☐ Suspect cause of disease
- ☐ Other

30. **15.c Are these data entered and stored in SMART Desktop? ***

Mark only one oval.

- ☐ All of these items are entered and stored in the corresponding SMART Conservation Area *Skip to question 32*
- ☐ Some of these items are entered and stored in the corresponding SMART Conservation Area *Skip to question 32*
- ☐ None of these items are recorded in the corresponding SMART Conservation Area *Skip to question 31*

31. **15.c.i If none of the items are recorded in the corresponding SMART Conservation Area where are they recorded? ***

Please mark "Yes/No" to answer the following questions

32. **16. Are domestic animals found in the protected area (free-ranging, captive, on a farm)? ***

Mark only one oval.

☐ Yes *Skip to question 33*

☐ No *Skip to question 37*

33. **17. If observed on a patrol, is the presence of domestic animals recorded? ***

Mark only one oval.

☐ Yes *Skip to question 34*

☐ No *Skip to question 37*

34. **17.a Are these data entered in SMART Desktop? ***

Mark only one oval.

☐ Yes

☐ No

35. **18. Is the health status of the observed domestic animals recorded (healthy, sick or injured, dead) recorded? ***

Mark only one oval.

☐ Yes *Skip to question 36*

☐ No *Skip to question 37*

36. **18.a Are these data entered in SMART Desktop? ***

Mark only one oval.

☐ Yes

☐ No

37. **19. Is any other data relevant to wildlife health collected during the patrols? ***

38. **20. If no data are collected on dead, sick, or injured wildlife, please select the best explanation(s) for why this information is not collected (check all that apply) ***

Check all that apply.

- ☐ Dead, sick or injured wildlife are seldom found
- ☐ It would add too much work or time to the patrol
- ☐ Wildlife disease is not relevant in the protected area
- ☐ Lack of expertise on how to properly record this information
- ☐ Never thought about it
- ☐ We use the default SMART data model which does not include sick, dead or injured wildlife

39. **21. What version of SMART Desktop is currently used in the protected area? ***

40. **22. Is SMART Connect available to manage and transfer information between SMART Desktop and SMART Mobile? ***

Mark only one oval.

- ☐ Yes *Skip to question 42*
- ☐ No *Skip to question 41*

41. **22.a Are there plans to set up SMART Connect instance and when? ***

42. **23. Is SMART fully rolled-out in the protected area(s) you work in or is it being piloted?** *

Mark only one oval.

- ☐ Fully rolled-out
- ☐ Under pilot
- ☐ Partially rolled-out

43. **24. Would you be interested in adding a set of categories and attributes to your data model in order to facilitate the collection of wildlife health data (morbidity/mortality findings and events)?** *

Mark only one oval.

- ☐ Yes
- ☐ No

44. **25. Please select the option that best describes your position** *

Mark only one oval.

- ☐ I am directly responsible for managing SMART data in one or more protected areas
- ☐ I am a general manager or administrator of one or more protected areas that uses SMART data
- ☐ Other


This content is neither created nor endorsed by Google.

Google Forms

Results


Perceptions of wildlife health importance

Appendix S2. Level of agreement (grey scale) among non-local protected area data managers with the statements “Pathogens carried by wildlife inhabiting the protected area(s) where I work in can affect livestock health.” (brown), “Pathogens carried by wildlife inhabiting the protected area(s) where I work in can affect human health.” (red), “Human or livestock pathogens can affect wildlife populations inhabiting the protected area(s) where I work in.” (blue), and “Wildlife health is important to achieve the conservation goals of the protected area(s) where I work,” (green).




Encounters with injured, sick, or dead wildlife and documentation

Appendix S3. Number of non-local protected area data managers who reported that the health status of and frequency of encounters with wildlife are recorded or not recorded in the protected area they worked in. Green bars represent the proportion of respondents that reported recording of wildlife in each category. Overall frequency of encounters with sick and injured wildlife was requested in a unique question; therefore, rows one and two show the same total number of responses per encounter category.




Documentation methods

Appendix S4. Distribution of the methods of documentation (second column) of healthy, sick, injured, or dead wildlife found during ranger patrols as reported by non-local protected area data managers and the recording of specific types of data for each wildlife health status across documentation methods (black line, 50%).




Domestic animals in protected areas

Appendix S5. Responses of non-local protected area data manager to statements that the presence and health of domestic animals in protected areas is a conservation concern (red, no recording of domestic animals; light blue, recording of domestic animals but not their health status; dark blue, recording of domestic animals and their health status). Data are from the group of protected area data managers who reported the presence of domestic animals in the protected area.



Discussion

Appendix S6. Distribution of protected area data managers responses (local and non-local) across their overall agreement with the statements “Human or livestock pathogens can affect wildlife populations inhabiting the protected area I work in” and “Introduced domestic animals are a concern for the conservation goals of the protected area” for those protected area managers (local and non-local) that reported the absence of domestic animals in the protected area.



Appendix S7. Distribution of protected area data managers responses (local and non-local) across their overall agreement with “Human or livestock pathogens can affect wildlife populations inhabiting the protected area I work in” and “Introduced domestic animals are a concern for the conservation goals of the protected area” for those protected area data managers that reported the presence of domestic animals in the protected area.

