

01 Math Review

Engineering Statics, STCS 200

Updated on: August 13, 2025

- ▶ Statics is all math! All but the most trivial statics problems require algebra and/or trigonometry and/or geometry to solve.
- ▶ The good news is that the math is not very difficult. You won't need anything more advanced than high-school math.
- ▶ We will do a quick review here that should cover all the math you'll need for STCS 200.

Triangles are a strong, stable shape and often used in engineering.


Triangles help avoid issues like this:



Triangles mean we need trigonometry.


Right Triangle

A **right triangle** is a triangle having one 90° angle.



Right Triangle


A **right triangle** is a triangle having one 90° angle.



Label the three sides a , b and c . The side a , opposite the right angle, is called the **hypotenuse**.

Right Triangle

A **right triangle** is a triangle having one 90° angle.




Label the three sides a , b and c . The side a , opposite the right angle, is called the **hypotenuse**.

If we know the lengths of any two sides, we can calculate the length of the third side using the **Pythagorean Theorem**:

Pythagorean Theorem

$$a^2 = b^2 + c^2$$

Right Triangle Exercises (1)



1. Use the Pythagorean Theorem to determine the lengths of CE and CB

- ▶ Significant **digits** and significant **figures** are the same thing.

- ▶ Significant **digits** and significant **figures** are the same thing.
- ▶ Significant digits are a measure of the **accuracy** of a number.

- ▶ Significant **digits** and significant **figures** are the same thing.
- ▶ Significant digits are a measure of the **accuracy** of a number.

It is **extremely important** to recognize that we can get no more accuracy out of a calculation than we put in. If the inputs to a problem have three significant digits, we cannot expect any higher accuracy than three significant digits in our result — even if the calculator does give ten digits.

Non-zero digits

Non-zero digits **are** significant:

- ▶ 1234 has 4 significant digits.
- ▶ 12.34 has 4 significant digits.

Non-zero digits

Non-zero digits **are** significant:

- ▶ 1234 has 4 significant digits.
- ▶ 12.34 has 4 significant digits.

Zeros between non-zero digits are significant

- ▶ 12034 has 5 significant digits.
- ▶ 12.0034 has 6 significant digits.

Leading zeros are **not** significant

- ▶ 0.1234 has 4 significant digits.
- ▶ 0.0001234 has 4 significant digits.

Leading zeros are **not** significant

- ▶ 0.1234 has 4 significant digits.
- ▶ 0.0001234 has 4 significant digits.

Trailing zeros (after a decimal point) **are** significant

- ▶ 1234.0 has 5 significant digits.
- ▶ 1.23400 has 6 significant digits.

Trailing zeros (on whole numbers, i.e. integers) are more **complicated**

- ▶ 12300 can have 3, 4 or 5 significant digits!

Trailing zeros (on whole numbers, i.e. integers) are more **complicated**

- ▶ 12300 can have 3, 4 or 5 significant digits!
 - ▶ Consider 12.3 m. This value has 3 significant digits. It is equal to 12300 mm, so in this case the 12300 also has 3 significant digits.

Trailing zeros (on whole numbers, i.e. integers) are more complicated

- ▶ 12300 can have 3, 4 or 5 significant digits!
 - ▶ Consider 12.3 m. This value has 3 significant digits. It is equal to 12300 mm, so in this case the 12300 also has 3 significant digits.
 - ▶ Now consider 12.30 m. This value has 4 significant digits. But it is still equal to 12300 mm, so in this case the 12300 has 4 significant digits.

Trailing zeros (on whole numbers, i.e. integers) are more complicated

- ▶ 12300 can have 3, 4 or 5 significant digits!
 - ▶ Consider 12.3 m. This value has 3 significant digits. It is equal to 12300 mm, so in this case the 12300 also has 3 significant digits.
 - ▶ Now consider 12.30 m. This value has 4 significant digits. But it is still equal to 12300 mm, so in this case the 12300 has 4 significant digits.
 - ▶ What if 12300 mm refers to 12.300 m? Then it has 5 significant digits.

Trailing zeros (on whole numbers, i.e. integers) are more complicated

- ▶ 12300 can have 3, 4 or 5 significant digits!
 - ▶ Consider 12.3 m. This value has 3 significant digits. It is equal to 12300 mm, so in this case the 12300 also has 3 significant digits.
 - ▶ Now consider 12.30 m. This value has 4 significant digits. But it is still equal to 12300 mm, so in this case the 12300 has 4 significant digits.
 - ▶ What if 12300 mm refers to 12.300 m? Then it has 5 significant digits.
- ▶ Usually, the trailing zeros are placeholders for the magnitude of a value and we don't need to worry unduly.

Trailing zeros (on whole numbers, i.e. integers) are more complicated

- ▶ 12300 can have 3, 4 or 5 significant digits!
 - ▶ Consider 12.3 m. This value has 3 significant digits. It is equal to 12300 mm, so in this case the 12300 also has 3 significant digits.
 - ▶ Now consider 12.30 m. This value has 4 significant digits. But it is still equal to 12300 mm, so in this case the 12300 has 4 significant digits.
 - ▶ What if 12300 mm refers to 12.300 m? Then it has 5 significant digits.
- ▶ Usually, the trailing zeros are placeholders for the magnitude of a value and we don't need to worry unduly.
- ▶ If we want to emphasize that 12300 has 4 significant digits, we can write $1.230 \times (10^3)$.

Calculations for Exercises

- ▶ In practice, it is often difficult to measure objects more accurately than to three significant digits so **input values for exercises are generally given to 3 significant digits.** (Or sometimes 4 significant digits when the leading significant digit is a 1)

Calculations for Exercises

- ▶ In practice, it is often difficult to measure objects more accurately than to three significant digits so **input values for exercises are generally given to 3 significant digits.** (Or sometimes 4 significant digits when the leading significant digit is a 1)
- ▶ We cannot expect to get more accuracy in our result at the end of a calculation than from our given input values at the beginning of the calculation so **solutions should be correct to 3 significant digits, not more than the accuracy of the calculation inputs!**

Calculations for Exercises

- ▶ In practice, it is often difficult to measure objects more accurately than to three significant digits so **input values for exercises are generally given to 3 significant digits.** (Or sometimes 4 significant digits when the leading significant digit is a 1)
- ▶ We cannot expect to get more accuracy in our result at the end of a calculation than from our given input values at the beginning of the calculation so **solutions should be correct to 3 significant digits, not more than the accuracy of the calculation inputs!**
- ▶ Intermediate calculations will accumulate rounding errors quickly if we use only three significant digits and these can affect the final result. **For intermediate calculations, use 5 or more significant digits.**

(When I write solutions down, I use 5 significant digits for intermediate calculations. You may use more if it is more convenient for you, e.g., if you are storing intermediate results in your calculator.)

Significant Digits and Rounding

- ▶ When using five significant digits for intermediate calculations, it is necessary to convert to three (or four) significant digits when providing the final answer to an exercise. Then, **rounding** is often involved:

Significant Digits and Rounding

- ▶ When using five significant digits for intermediate calculations, it is necessary to convert to three (or four) significant digits when providing the final answer to an exercise. Then, **rounding** is often involved:
 - ▶ 2.3456 becomes 2.35 because the first non-significant digit (the 5 in this case) is ≥ 5 and so the 4 rounds up.

Significant Digits and Rounding

- ▶ When using five significant digits for intermediate calculations, it is necessary to convert to three (or four) significant digits when providing the final answer to an exercise. Then, **rounding** is often involved:
 - ▶ 2.3456 becomes 2.35 because the first non-significant digit (the 5 in this case) is ≥ 5 and so the 4 rounds up.
 - ▶ 2.3446 becomes 2.34 because the first non-significant digit (the 4 in this case) is < 5 and the last significant digit, 4, remains unchanged.

Significant Digits and Rounding


- ▶ When using five significant digits for intermediate calculations, it is necessary to convert to three (or four) significant digits when providing the final answer to an exercise. Then, **rounding** is often involved:
 - ▶ 2.3456 becomes 2.35 because the first non-significant digit (the 5 in this case) is ≥ 5 and so the 4 rounds up.
 - ▶ 2.3446 becomes 2.34 because the first non-significant digit (the 4 in this case) is < 5 and the last significant digit, 4, remains unchanged.
- ▶ When the first discarded digit is a 5 (or higher), round up the digit before the 5 (or higher)

Significant Digits and Rounding

- ▶ When using five significant digits for intermediate calculations, it is necessary to convert to three (or four) significant digits when providing the final answer to an exercise. Then, **rounding** is often involved:
 - ▶ 2.3456 becomes 2.35 because the first non-significant digit (the 5 in this case) is ≥ 5 and so the 4 rounds up.
 - ▶ 2.3446 becomes 2.34 because the first non-significant digit (the 4 in this case) is < 5 and the last significant digit, 4, remains unchanged.
- ▶ When the first discarded digit is a 5 (or higher), round up the digit before the 5 (or higher)
- ▶ There are various rules (such as the odd-even rule) which take a more complicated approach to rounding 5 but, for our purposes, **5 rounds up!**


More About Right Triangle

The sine, cosine and tangent trigonometrical functions relate an acute angle (θ , in this example) in a right triangle to two of the sides of the triangle.



More About Right Triangle

The sine, cosine and tangent trigonometrical functions relate an acute angle (θ , in this example) in a right triangle to two of the sides of the triangle.




Right Triangle Trigonometry Formulæ

$$\sin \theta = \frac{\text{opposite}}{\text{hypotenuse}}, \quad \cos \theta = \frac{\text{adjacent}}{\text{hypotenuse}}, \quad \tan \theta = \frac{\text{opposite}}{\text{adjacent}}$$

Remember: **SOHCAHTOA**


Right Triangle Exercises (2)



2. Use the **tangent** function to calculate $\angle CEF$.
3. From $\angle CEF$ just found (use the **intermediate, 5 or more significant digit, form!**) and the **sine** rule to verify the length of CE found earlier.
4. Use the **cosine** function and the length of CB found earlier to calculate the angle between BC and the horizontal.
5. Use the **tangent** function to verify the previous result.

Triangles - Sine Rule

Not all triangles contain a right angle. To solve for these triangles (finding the lengths of the side and the triangle angles), we have to employ some different tools: the **sine rule** and (later) the **cosine rule**



Sine Rule

$$\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}$$


or

Sine Rule

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

Triangles - Sine Rule Exercises

6. Using the sine rule, find $\angle ACB$.
7. Using the sine rule, find $\angle ABC$.
8. Sum the interior angles of the triangle.



A couple of trig identities that will come in useful:

Identities

$$\sin(180^\circ - \theta) = \sin \theta$$

$$\cos(-\theta) = \cos \theta$$

Note:

$$\sin(140^\circ) = \sin(40^\circ) = 0.64279$$


$$\cos(42^\circ) = \cos(-42^\circ) = 0.74314$$

Thus, we have to be careful when using inverse trigonometric functions:

$$\sin^{-1}(0.64279) = 40^\circ \text{ or } 140^\circ$$

$$\cos^{-1} 0.74314 = -42^\circ \text{ or } 42^\circ$$

Triangles - Cosine Rule



Cosine Rule

$$a^2 = b^2 + c^2 - 2bc \cos A$$


$$b^2 = a^2 + c^2 - 2ac \cos B$$

$$c^2 = a^2 + b^2 - 2ab \cos C$$


The cosine rule is useful when you have all the sides of a triangle and want to find the angles.

Triangles - Cosine Rule Exercises

9. Determine $\angle ABC$, using the value for AB found earlier
10. Compare the value for $\angle ABC$ with the value calculated earlier. Is it the same? It should be!




Similar Triangles



If triangles ABC and XYZ have the same angles, they are said to be **similar triangles**.

Similar Triangles



If triangles ABC and XYZ have the same angles, they are said to be **similar triangles**.

The ratios of the lengths of corresponding sides of similar triangles are equal:

$$\frac{AB}{XY} = \frac{BC}{XZ} = \frac{AC}{YZ}$$


Similar Triangles - Exercises

$ABCD$ is a rigid (i.e., it does not deform) plate, pinned at C .

When horizontal force P is applied at A , $ABCD$ rotates about C and A deflects 2.45 mm horizontally rightwards.

Assume that BF remains horizontal and that DE remains vertical.

11. Determine δ_{BF} , the change in length of BF .
12. Determine δ_{DE} , the change in length of DE .



Similar Triangles - Exercises

$ABCD$ is a rigid (i.e., it does not deform) plate, pinned at C .

When horizontal force P is applied at A , $ABCD$ rotates about C and A deflects 2.45 mm horizontally rightwards.

Assume that BF remains horizontal and that DE remains vertical.

13. Determine δ_{BF} , the change in length of BF .
14. Determine δ_{DE} , the change in length of DE .

