

Computing with cut systems

Daniel Murfet

December 14, 2015

Abstract

The bicategory \mathcal{LG} of Landau-Ginzburg models has polynomials as objects and matrix factorisations as 1-morphisms. The composition of 1-morphisms naturally produces infinite rank matrix factorisations, which is a nuisance. In this paper we define a bicategory \mathcal{C} which is equivalent to \mathcal{LG} but for which composition of 1-morphisms produces finite rank matrix factorisations equipped with the action of a Clifford algebra.

Contents

1	Introduction	2
1.1	A_∞ -algebras and minimal models	3
1.2	A geometric enrichment of \mathcal{LG}	4
1.3	Semantics of linear logic	4
2	Background	5
2.1	Supercategories	5
2.2	The superbicategory of Landau-Ginzburg models	8
2.3	Partial derivatives as homotopies	10
2.4	Clifford algebras	11
2.5	The Clifford thickening	13
3	The superbicategory \mathcal{C}	16
4	The equivalence of \mathcal{C} and \mathcal{LG}	23
4.1	Homological perturbation and fixed points	24
4.2	Clifford actions and Koszul complexes	27
4.3	The finite model	34
4.4	The equivalence of \mathcal{C} and \mathcal{LG}	39
4.5	Example: the Hom complex	41
A	Tensor products in supercategories	43

1 Introduction

The bicategory \mathcal{LG} of Landau-Ginzburg models has a rich structure, as can be seen by its important role in topological field theory [26, 7, 11] homological knot invariants [23] and 2-representations of quantum groups [?], but the definition is still relatively elementary. This makes \mathcal{LG} an attractive algebraic object of study, but for one major wrinkle: the composition operation for 1-morphisms is infinitary.

The objects of \mathcal{LG} are potentials $W(x)$, while 1-morphisms $W(x) \longrightarrow V(y)$ are matrix factorisations of the difference $V(y) - W(x)$ over $\mathbb{C}[x, y]$. Such a matrix factorisation is a \mathbb{Z}_2 -graded free $\mathbb{C}[x, y]$ -module $X = X^0 \oplus X^1$ together with an odd operator d_X satisfying $d_X^2 = (V(y) - W(x)) \cdot 1_X$. For a third potential $U(z)$, the composition of 1-morphisms is a functor

$$\begin{aligned} \mathcal{LG}(V, U) \times \mathcal{LG}(W, V) &\longrightarrow \mathcal{LG}(W, U), \\ (Y, X) &\mapsto Y \circ X \end{aligned}$$

defined for matrix factorisations Y of $V - U$ and X of $W - V$ by

$$(1.1) \quad Y \circ X = (Y \otimes_{\mathbb{C}[y]} X, d_Y \otimes 1 + 1 \otimes d_X).$$

This is the analogue of convolution of Fourier-Mukai kernels in the matrix factorisation setting and suffers from the same defect, namely, the resulting matrix factorisation $Y \circ X$ is a free module of *infinite rank* over $\mathbb{C}[x, z]$ which is only homotopy equivalent (that is, isomorphic in the homotopy category) to a finite rank matrix factorisation. While this finite model can be proven abstractly to exist, there is no obvious way to describe it as a function of the input data Y, X viewed as matrices of polynomials. This need to deal throughout with infinite rank matrix factorisations introduces a certain minimal level of complexity into the theory, which for many applications is undesirable.

Together with Dyckerhoff we initiated in [15] the study of these finite models in terms of homological perturbation and Atiyah classes, and in the present paper this program reaches its logical conclusion. We introduce a bicategory \mathcal{C} in which the objects are again potentials and where the 1-morphisms are finite rank matrix factorisations equipped with the structure of a representation of a Clifford algebra. We will prove that this bicategory is equivalent to \mathcal{LG} , but is much simpler - in particular, it has a composition operation which is described by a finite number of polynomial functions of the input data.

More precisely: the composition of 1-morphisms in \mathcal{C} is a functor

$$(1.2) \quad \mathcal{C}(V, U) \times \mathcal{C}(W, V) \longrightarrow \mathcal{C}(W, U),$$

$$(1.3) \quad (Y, X) \mapsto Y \mid X$$

which we call the *cut operation*. The definition of the cut is very simple. With $J_V = \mathbb{C}[y]/(\partial V)$ the Jacobi algebra of \mathbb{C} (which is a finite-dimensional \mathbb{C} -vector space by the assumption that $V(y)$ is a potential) the cut of Y and X is a finite rank matrix factorisation

$$(1.4) \quad Y \mid X = Y \otimes_{\mathbb{C}[y]} J_V \otimes_{\mathbb{C}[y]} X = (Y \otimes_{\mathbb{C}[y]} X)/(\partial V)(Y \otimes_{\mathbb{C}[y]} X)$$

over $\mathbb{C}[x, z]$

equipped with odd operators $\gamma_i, \gamma_i^\dagger$ satisfying Clifford relations

$$\begin{aligned} \gamma_i e &= \gamma_i = e \gamma_i, \\ \gamma_i^\dagger e &= \gamma_i^\dagger = e \gamma_i^\dagger. \end{aligned}$$

up to homotopy (see Theorem 3.9). The differential on $Y \mid X$ is induced from the differential on $Y \otimes X$ and can be written explicitly as a matrix over $\mathbb{C}[x, z]$ in terms of the polynomial entries of d_Y, d_X . The Clifford action and the matrices which define the Clifford actions on Y, X . In particular, infinite rank matrix factorisations are never involved. In the remainder of the introduction we sketch how the cut operation arises from homological perturbation, why Clifford algebras are involved, where the equivalence $\mathcal{C} \cong \mathcal{LG}$ comes from, and our motivations for pursuing this “finite model” \mathcal{C} of \mathcal{LG} .

The main theorem (Theorem 4.34) says

$$(1.5) \quad Y \mid X \begin{array}{c} \xleftarrow{\Phi} \\ \xrightarrow{\Phi^{-1}} \end{array} S_m \otimes_k (Y \otimes X)$$

The paper is structured as follows: in Section 2 we recall the definition of superbicategories and Clifford algebras, and introduce the Clifford thickening of a supercategory. In Section 3 we define \mathcal{C} and in Section 4 we prove it is equivalent to \mathcal{LG} as a supercategory without units. We do not consider unit 1-morphisms here because \mathcal{C} does not make working with units any simpler.

1.1 A_∞ -algebras and minimal models

The relationship between \mathcal{LG} and \mathcal{C} is similar to the relationship between a DG-algebra and its A_∞ -minimal model. Given a DG algebra A over a field k , which is infinite-dimensional but has finite-dimensional cohomology, the A_∞ -minimal model is an A_∞ -structure on the cohomology $H^*(A)$ which is A_∞ -quasi-isomorphic to A . The tradeoff being made is that A is infinite-dimensional but with a simple structure (i.e. one multiplication map m_2) while $H^*(A)$ is finite-dimensional but with possibly an infinite family of higher multiplications. In \mathcal{LG} the composition operation is not described by a finite amount of data, and the price that one pays by going to \mathcal{C} where this is the case, is that the composition has additional structure (a Clifford action). There is a more precise link to A_∞ -minimal models. A special case of the equivalence between \mathcal{C} and \mathcal{LG} gives, for

a matrix factorisation X of the potential $W(x_1, \dots, x_m)$ viewed, a strong deformation retract

$$(1.6) \quad \Phi : S_m \otimes_k \text{Hom}_R(X, X) \longrightarrow X^\vee | X.$$

where $S_m = \bigwedge(k\theta_1 \oplus \dots \oplus k\theta_m)$ is an exterior algebra, $\text{Hom}_R(X, X)$ is the Hom complex. This deformation retract can be used as the input data for the usual construction of an A_∞ -minimal on the left hand side, viewed as a DG-algebra.

1.2 A geometric enrichment of \mathcal{LG}

A finite rank matrix factorisation of $V(y) - W(x)$ is a matrix of polynomials

$$D \in M_{2r}(\mathbb{C}[x, y]) \quad \text{satisfying} \quad D^2 = (V(y) - W(x)) \cdot I_{2r}$$

where we require that, dividing D into $r \times r$ blocks, the upper left and bottom right blocks are zero matrices. The set of finite rank matrix factorisations is denoted $\mathcal{LG}(W, V)_{fin}$. For technical reasons the category of 1-morphisms $\mathcal{LG}(W, V)$ contains, in addition to the finite rank matrix factorisations, those infinite rank matrix factorisations which are homotopy equivalent to finite rank ones; this means in particular that the inclusion

$$\mathcal{LG}(W, V)_{fin} \longrightarrow \mathcal{LG}(W, V)$$

is an equivalence of categories. If we fix the size r of the blocks and bound the degrees of the monomials appearing in D by some integer k , we can parametrise the possible matrix factorisations by the closed points of an affine scheme $\mathcal{M}_{r,k}(W, V)$ whose coordinates are the coefficients of the various monomials in each row and column of D . Hence as sets we have

$$(1.7) \quad \mathcal{LG}(W, V)_{fin} = \bigsqcup_{r \geq 1} \bigcup_{k \geq 0} \mathcal{M}_{r,k}(W, V)_{\mathbb{C}}.$$

does indeed lift to a morphism of the parametrising schemes - fulfilling the original goal. The schemes parametrising 1-morphisms in \mathcal{C} together with the morphisms of schemes lifting composition, and vector bundles encoding the 2-morphisms, is what we refer to as a *cut system* in the sequel [27]. However the present paper is concerned only with defining the supercategory \mathcal{C} and proving that it is equivalent to \mathcal{LG} . TODO: A cut system is a bicategory enriched in a certain sense in the category of (indexed families of) affine schemes.

1.3 Semantics of linear logic

In forthcoming work we will use the results of this paper to define a novel semantics of linear logic, in which formulas are interpreted by tuples $(\mathcal{X}, \mathbb{C}[x], W)$ where \mathcal{X} is

a scheme and $W \in \mathbb{C}[x]$ is a potential. Proofs in the logic are interpreted by certain correspondences between these pairs. To explain briefly why we are forced to use \mathcal{C} rather than \mathcal{LG} , we will use the denotation of the Church numeral $\underline{2}$ (see [28]).

Let x be a variable of the logic and suppose it has denotation $\llbracket x \rrbracket = (\text{Spec}(\mathbb{C}), \mathbb{C}[x], W)$ in the semantics. Let $\mathcal{M} = \mathcal{M}(W, W)$ be the (indexed) affine scheme parametrising 1-morphisms $X : W \rightarrow W$ in \mathcal{C} , as discussed above. Then the semantics works as follows:

$$\begin{aligned} \llbracket x \multimap x \rrbracket &= (\text{Spec}(\mathbb{C}), \mathbb{C}[x, x'], W(x) - W(x')) , \\ \llbracket !(x \multimap x) \rrbracket &= (\mathcal{M}, \mathbb{C}, 0) . \end{aligned}$$

The proof $\underline{2}$ of the sequent $!(x \multimap x) \vdash x \multimap x$ has for its denotation a correspondence between $\llbracket !(x \multimap x) \rrbracket$ and $\llbracket x \multimap x \rrbracket$, which concretely means that $\llbracket \underline{2} \rrbracket$ is a bundle over the scheme \mathcal{M} whose fiber over a loop X , viewed as a point $[X] \in \mathcal{M}$, is the square $X \mid X$. More precisely, $\llbracket \underline{2} \rrbracket$ is the bundle over $\mathcal{M} \times \text{Spec}(\mathbb{C}[x, x'])$ whose fiber over $([X], a)$ is

$$(1.8) \quad \llbracket \underline{2} \rrbracket_{[X], a} = (X \mid X) \otimes_{\mathbb{C}[x, x']} \mathbb{C}(a) .$$

For these definitions to work, the coefficients of the monomials in the matrix factorisation $X \mid X$ need to be given explicitly as polynomial functions of the coefficients in X . This means we have to use \mathcal{C} rather than \mathcal{LG} .

Acknowledgements. Thanks to Nils Carqueville for helpful comments on the draft, and to Ingo Runkel and Rafal Suszek for kindly allowing the use of Figure ?? from [29].

2 Background

Throughout k is a commutative \mathbb{Q} -algebra. By default categories and functors are k -linear.

2.1 Supercategories

A supercategory has for every 1-morphism F an associated 1-morphism ΨF with $\Psi^2 \cong 1$, and in this sense the categories of 1-morphisms are \mathbb{Z}_2 -graded. The bicategory of Landau-Ginzburg models is an example of a supercategory. Since Clifford algebras (which are \mathbb{Z}_2 -graded) play a fundamental role in the definition of cut systems, supercategories are the natural language. Our main references for the foundations are [16, 19, 20], but we give the full definitions below as these references work only with strict bicategories.

Definition 2.1. A *supercategory* is a category \mathcal{C} together with a functor $\Psi : \mathcal{C} \rightarrow \mathcal{C}$ and a natural isomorphism $\xi : \Psi^2 \rightarrow 1_{\mathcal{C}}$ satisfying the condition

$$\xi * 1_{\Psi} = 1_{\Psi} * \xi$$

as natural transformations $\Psi^3 \rightarrow \Psi$. A *superfunctor* (F, γ) from a supercategory $(\mathcal{C}, \Psi_{\mathcal{C}})$ to a supercategory $(\mathcal{D}, \Psi_{\mathcal{D}})$ is a functor $F : \mathcal{C} \rightarrow \mathcal{D}$ together with a natural isomorphism $\gamma : F\Psi_{\mathcal{C}} \rightarrow \Psi_{\mathcal{D}}F$ satisfying

$$1_F * \xi = (\xi * 1_F)(1_{\Psi} * \gamma)(\gamma * 1_{\Psi}) .$$

A *supernatural transformation* $\varphi : F \longrightarrow G$ between superfunctors $(F, \gamma_F), (G, \gamma_G)$ is a natural transformation making the following diagram commute:

$$\begin{array}{ccc} F\Psi & \xrightarrow{\varphi\Psi} & G\Psi \\ \gamma_F \downarrow & & \downarrow \gamma_G \\ \Psi F & \xrightarrow{\Psi\varphi} & \Psi G \end{array} .$$

Example 2.2. If A is a \mathbb{Z}_2 -graded k -algebra there is a supercategory \mathcal{A} with set of objects $|\mathcal{A}| = \mathbb{Z}_2$ and morphisms $\mathcal{A}(0, 0) = \mathcal{A}(1, 1) = A_0, \mathcal{A}(0, 1) = \mathcal{A}(1, 0) = A_1$ with the obvious composition and addition rules. The functor Ψ is defined on objects by $\Psi(i) = i + 1$ and on morphisms by the identity, while $\xi = 1$.

Let \mathcal{C} be a supercategory. For objects $X, Y \in \mathcal{C}$ we define the \mathbb{Z}_2 -graded k -module

$$\mathcal{C}^*(X, Y) = \prod_{i \in \mathbb{Z}_2} \mathcal{C}(X, \Psi^i Y) .$$

There is for any triple X, Y, Z of objects a morphism of graded modules

$$\mathcal{C}^*(Y, Z) \otimes_k \mathcal{C}^*(X, Y) \longrightarrow \mathcal{C}^*(X, Z)$$

given for instance by $\varphi \otimes \psi \mapsto \xi \Psi(\varphi) \psi$ when φ, ψ are both of degree one. This makes \mathcal{C} into a category enriched over the monoidal category of \mathbb{Z}_2 -graded k -modules.

There are isomorphisms of graded modules $\mathcal{C}^*(X, \Psi Y) \cong \Psi \mathcal{C}^*(X, Y) \cong \mathcal{C}^*(\Psi X, Y)$ where $\Psi V = V[1]$ denotes the grading shift on a \mathbb{Z}_2 -graded k -module. For \mathbb{Z}_2 -graded k -modules V, W we write $\text{Hom}_k^0(V, W)$ for the module of degree zero maps and $\text{Hom}_k^*(V, W)$ for the \mathbb{Z}_2 -graded module of homogeneous maps. If A is a \mathbb{Z}_2 -graded k -algebra then A^{op} denotes the same underlying graded module with multiplication $f * g = (-1)^{|f||g|} gf$.

Definition 2.3. Let A be a \mathbb{Z}_2 -graded k -algebra and \mathcal{C} a supercategory. A *left A -module* in \mathcal{C} is an object $X \in \mathcal{C}$ together with a morphism of \mathbb{Z}_2 -graded algebras $A \longrightarrow \mathcal{C}^*(X, X)$. A *morphism* of left A -modules is a morphism in \mathcal{C} which commutes with the A -action in the obvious sense. The category of left A -modules in \mathcal{C} is denoted \mathcal{C}_A . Right modules are defined similarly, using A^{op} .

Remark 2.4. Viewing A as a supercategory \mathcal{A} as in Example 2.2, there is an equivalence between the category of A -modules in \mathcal{C} and the category of superfunctors $\mathcal{A} \longrightarrow \mathcal{C}$ and supernatural transformations.

Remark 2.5. The *idempotent completion* \mathcal{C}^ω of a category \mathcal{C} has as objects pairs (X, e) consisting of an object $X \in \mathcal{C}$ and an idempotent endomorphism $e : X \longrightarrow X$. In \mathcal{C}^ω a morphism $\phi : (X, e) \longrightarrow (Y, e)$ is a morphism $\phi : X \longrightarrow Y$ satisfying $\phi e = \phi$ and $e \phi = \phi$. The identity on (X, e) is e . There is a fully faithful functor $\iota : \mathcal{C} \longrightarrow \mathcal{C}^\omega, \iota(X) = (X, 1_X)$ which is an equivalence if \mathcal{C} is idempotent complete (i.e. all idempotents split in \mathcal{C}).

If \mathcal{C} is a supercategory then there is a functor $\Psi^\omega : \mathcal{C}^\omega \longrightarrow \mathcal{C}^\omega$ defined on objects by $\Psi(X, e) = (\Psi X, \Psi e)$ and a natural isomorphism $\xi : \Psi^\omega \circ \Psi^\omega \longrightarrow 1_{\mathcal{C}^\omega}$ defined by $\xi_{(X, e)} = e \xi_X$. The tuple $(\mathcal{C}^\omega, \Psi^\omega, \xi^\omega)$ is a supercategory.

For background on bicategories see [3, 2, 18, 22, 24]. We follow the notation of [7], so that lower case letters a, b, \dots denote objects of a bicategory, while upper case letters X, Y, \dots and greek letters α, β, \dots respectively denote 1-morphisms and 2-morphisms. Units are denoted Δ , associators are α , and unitors are λ, ρ . The composition of Y, X is denoted YX or $Y \circ X$.

Definition 2.6. A *superbicategory* is a bicategory \mathcal{B} together with the data:

- For each object a a 1-morphism $\Psi_a : a \longrightarrow a$ and a 2-isomorphism $\xi_a : \Psi_a^2 \longrightarrow \Delta_a$.
- For each 1-morphism $X : a \longrightarrow b$ a natural 2-isomorphism $\gamma_X : X\Psi_a \longrightarrow \Psi_b X$.

This data is required to satisfy the following axioms:

- For each composable pair X, Y of 1-morphisms the diagram

$$\begin{array}{ccc}
 (YX)\Psi & \xrightarrow{\gamma_{YX}} & \Psi(YX) \\
 \alpha \downarrow & & \uparrow \alpha \\
 Y(X\Psi) & \xrightarrow{1_Y * \gamma_X} Y(\Psi X) \xrightarrow{\alpha^{-1}} (Y\Psi)X \xrightarrow{\gamma_Y * 1_X} & (\Psi Y)X
 \end{array}$$

commutes.

- For every object a , $\xi_a * 1_\Psi = 1_\Psi * \xi_a$.
- For every 1-morphism $X : a \longrightarrow b$, $1_X * \xi_a = (\xi_b * 1_X)(1_\Psi * \gamma_X)(\gamma_X * 1_\Psi)$.

Example 2.7. Supercategories, superfunctors and supernatural transformations form a superbicategory $\text{Cat}_k^{\text{sup}}$.

A superbicategory can be constructed out of a bicategory \mathcal{B} in which the categories $\mathcal{B}(a, b)$ are all equipped with the structure of supercategories; see Appendix B.

Example 2.8. There is a bicategory of \mathbb{Z}_2 -graded k -algebras where 1-morphisms are \mathbb{Z}_2 -graded bimodules and 2-morphisms are degree zero bimodule maps. Given a B - A -module M the shift $M[1]$ has the grading $M[1]_i = M_{i+1}$ and the left and right action given by

$$b \cdot m = (-1)^{|b|}bm, \quad m \cdot a = (-1)^{|a|}ma.$$

This is a functor $\Psi = (-)[1]$ on the category of B - A -bimodules and with $\xi = 1$ this defines the structure of a supercategory on this category of bimodules. The usual isomorphisms of bimodules τ

$$(2.1) \quad N[1] \otimes M \longrightarrow (N \otimes M)[1], \quad n \otimes m \mapsto n \otimes m$$

$$(2.2) \quad N \otimes M[1] \longrightarrow (N \otimes M)[1], \quad n \otimes m \mapsto (-1)^{|n|}n \otimes m$$

satisfy the conditions in Appendix B and therefore give the bicategory of \mathbb{Z}_2 -graded algebras and bimodules the structure of a superbicategory.

Definition 2.9. Given two supercategories \mathcal{B}, \mathcal{C} a *lax superfunctor* $\Phi : \mathcal{B} \longrightarrow \mathcal{C}$ is a lax functor together with, for each object a in \mathcal{B} , a 2-morphism

$$\kappa_a : \Psi_{\Phi a} \longrightarrow \Phi(\Psi_a)$$

satisfying two coherence conditions:

- (a) for all a , commutativity of

$$\begin{array}{ccc} \Psi_{\Phi a} \Psi_{\Phi a} & \xrightarrow{\kappa * \kappa} & \Phi(\Psi_a) \Phi(\Psi_a) \\ \downarrow \xi & & \downarrow \\ & & \Phi(\Psi_a \Psi_a) \\ & & \downarrow \Phi(\xi) \\ \Delta_{\Phi a} & \longrightarrow & \Phi(\Delta_a) \end{array}$$

- (b) for each 1-morphism $X : a \longrightarrow b$ in \mathcal{B} , commutativity of

$$\begin{array}{ccccc} \Phi(X) \Psi_{\Phi a} & \xrightarrow{1 * \kappa} & \Phi(X) \Phi(\Psi_a) & \longrightarrow & \Phi(X \Psi_a) \\ \downarrow \gamma & & & & \downarrow \Phi(\gamma) \\ \Psi_{\Phi b} \Phi(X) & \xrightarrow{\kappa * 1} & \Phi(\Psi_b) \Phi(X) & \longrightarrow & \Phi(\Psi_b X) \end{array}$$

A *strong superfunctor* is a lax superfunctor with κ_a an isomorphism for every object a .

2.2 The supercategory of Landau-Ginzburg models

A polynomial $W \in k[x_1, \dots, x_n]$ is a *potential* if it satisfies the three conditions set out in [7, Section 2.2], the two most important being that the partial derivatives $\partial_{x_1} W, \dots, \partial_{x_n} W$ form a quasi-regular sequence in $k[x]$ and that the quotient $J_W = k[x]/(\partial_{x_1} W, \dots, \partial_{x_n} W)$ is a finitely generated free k -module. Typical examples are the ADE singularities [?, I §2.4] of which the simplest are the A_N -singularities $W_{A_N} = x_1^{N+1} + x_2^2 + \dots + x_n^2$.

A *matrix factorisation* of W over $k[x]$ is a \mathbb{Z}_2 -graded free $k[x]$ -module $X = X^0 \oplus X^1$ together with an odd operator (the differential) $d_X : X \longrightarrow X$ satisfying $d_X^2 = W \cdot 1_X$. A matrix factorisation (X, d_X) is *finite rank* if the underlying free module is finitely generated. In this case we may, after choosing a homogeneous basis, write

$$d_X = \begin{pmatrix} 0 & d_X^1 \\ d_X^0 & 0 \end{pmatrix}$$

for matrices of polynomials d_X^0, d_X^1 . Morphisms of matrix factorisations are degree zero $k[x]$ -linear maps which commute with the differentials. There is a homotopy relation on morphisms, and the *homotopy category* $\text{HMF}(k[x], W)$ of matrix factorisations has as

objects matrix factorisations and as morphisms the equivalence classes of morphisms of matrix factorisations up to homotopy. By $\text{hmf}(k[x], W)$ we denote the full subcategory of finite rank matrix factorisations. For background we refer to [31].

By $\text{hmf}(k[x], W)^\oplus$ we denote the full subcategory of $\text{HMF}(k[x], W)$ consisting of matrix factorisations which are direct summands (in the homotopy category) of finite rank matrix factorisations. Since $\text{HMF}(k[x], W)$ is idempotent complete, there is an equivalence (see Remark 2.5 for the notation)

$$(2.3) \quad \text{hmf}(k[x], W)^\omega \longrightarrow \text{hmf}(k[x], W)^\oplus$$

sending a pair (X, e) to an infinite rank matrix factorisation splitting the idempotent e .

If (X, d_X) is a matrix factorisation then so is $(X[1], -d_X)$, and this defines a functor $\Psi = (-)[1]$ on the homotopy category of matrix factorisations. Together with the identity $\xi : \Psi^2 \longrightarrow 1$ this makes $\text{HMF}(k[x], W)$ and $\text{hmf}(k[x], W)$ into supercategories.

The bicategory of Landau-Ginzburg models \mathcal{LG} has for its objects pairs (x, W) consisting of an ordered set of variables $x = (x_1, \dots, x_n)$ and a potential $W \in k[x] = k[x_1, \dots, x_n]$. Given potentials $W \in k[x]$ and $V \in k[y]$ the supercategory $\mathcal{LG}(W, V)$ is

$$\mathcal{LG}(W, V) = \text{hmf}(k[x, y], V - W)^\oplus.$$

Given a third potential $U \in k[z]$ the composition law is a functor

$$\mathcal{LG}(W, V) \otimes_k \mathcal{LG}(V, U) \longrightarrow \mathcal{LG}(W, U)$$

which sends a pair of 1-morphisms $X : W \longrightarrow V$ and $Y : V \longrightarrow U$ to the tensor product

$$(2.4) \quad Y \otimes X = (Y \otimes_{k[y]} X, d_{Y \otimes X} = d_Y \otimes 1 + 1 \otimes d_X).$$

Thus $Y \circ X := Y \otimes X$. This statement requires some care: $k[x, y, z]$ is an infinite rank free module over $k[x, z]$ and so $Y \otimes X$ is an infinite rank matrix factorisation of $U - W$ over $k[x, z]$. However one can prove that $Y \otimes X$ is a direct summand of a finite rank matrix factorisation [15] and therefore a valid object in $\mathcal{LG}(W, U)$.

Let $W \in k[x_1, \dots, x_n]$ be a potential and $W(x')$ the same polynomial but in a second set of variables x'_1, \dots, x'_n . Using formal symbols Θ_i of odd degree, we define the $k[x, x']$ -module

$$\Delta_W = \bigwedge \left(\bigoplus_{i=1}^n k[x, x'] \cdot \Theta_i \right).$$

Equipped with a certain differential (see [7]) this is a matrix factorisation of $W(x) - W(x')$ and defines the unit in $\mathcal{LG}(W, W)$ for composition of 1-morphisms. For the associator and unitor maps $\rho : X \otimes \Delta_W \longrightarrow X$ and $\lambda : \Delta_V \otimes X \longrightarrow X$ we also refer to *ibid*.

Lemma 2.10. *\mathcal{LG} is a superbicategory.*

Proof. The categories $\mathcal{LG}(W, V)$ are all naturally supercategories, and the same isomorphisms as in (2.1), (2.2) define the necessary natural isomorphisms τ to equip \mathcal{LG} with the structure of a superbicategory with $\Psi_W = \Delta_W[1]$, using Appendix B. \square

The superbicategory structure on \mathcal{LG} is used implicitly in [7, Section 7].

2.3 Partial derivatives as homotopies

Let $R = k[x_1, \dots, x_n]$ be a polynomial ring and X a finite rank matrix factorisation of W over R . Let us fix a homogeneous basis for X so that we may identify d_X with a matrix, and differentiate this matrix entry-wise to obtain the matrix $\lambda_i = \partial_{x_i}(d_X)$. It is easy to check that

Lemma 2.11. *The odd R -linear operator λ_i on X is independent, up to homotopy, of the choice of basis.*

Lemma 2.12. *As operators on X we have $[d_X, \lambda_i] = \partial_{x_i} W \cdot 1_X$.*

Proof. Applying the Leibniz rule to the equation $d_X d_X = W$ we obtain

$$(2.5) \quad d_X \partial_{x_i}(d_X) + \partial_{x_i}(d_X) d_X = \partial_{x_i} W$$

as claimed. □

It follows that any element in the Jacobi ideal $(\partial_{x_1} W, \dots, \partial_{x_n} W)$ acts null-homotopically on any morphism in the homotopy category $\text{hmf}(R, W)$. Moreover these homotopies λ_i are central, in the following sense:

Lemma 2.13. *Let $\phi : X \rightarrow X'$ be a morphism of matrix factorisations and write $\lambda_i = \partial_{x_i}(d_X)$ and $\lambda'_i = \partial_{x_i}(d_{X'})$. There is a homotopy $\phi \lambda_i \simeq \lambda'_i \phi$.*

Proof. From $d_{X'} \phi = \phi d_X$ we deduce

$$\partial_{x_i}(d_{X'}) \phi + d_{X'} \partial_{x_i}(\phi) = \partial_{x_i}(\phi) d_X + \phi \partial_{x_i}(d_X)$$

and hence

$$\partial_{x_i}(d_{X'}) \phi - \phi \partial_{x_i}(d_X) = \partial_{x_i}(\phi) d_X - d_{X'} \partial_{x_i}(\phi)$$

as claimed. □

Lemma 2.14. *There is a homotopy $[\lambda_i, \lambda_j] \simeq \partial_{x_i x_j} W \cdot 1_X$.*

Proof. Differentiating (2.5) with respect to j leaves

$$\partial_{x_j}(d_X) \partial_{x_i}(d_X) + d_X \partial_{x_j x_i}(d_X) + \partial_{x_j x_i}(d_X) d_X + \partial_{x_i}(d_X) \partial_{x_j}(d_X) = \partial_{x_i x_j} W$$

from which we read off

$$\lambda_i \lambda_j + \lambda_j \lambda_i = \partial_{x_i x_j} W + [d, -\partial_{x_j x_i}(d_X)].$$

□

2.4 Clifford algebras

The cut operation in \mathcal{C} produces objects which are representations of Clifford algebras. We briefly recall the basic theory of these algebras and their modules; for a full discussion see [17]. For $n \geq 0$ the Clifford algebra C_n is the associative \mathbb{Z}_2 -graded k -algebra generated by odd elements $\gamma_1, \dots, \gamma_n$ and $\gamma_1^\dagger, \dots, \gamma_n^\dagger$ subject to the following relations

$$(2.6) \quad [\gamma_i, \gamma_j] = 0, \quad [\gamma_i^\dagger, \gamma_j^\dagger] = 0, \quad [\gamma_i, \gamma_j^\dagger] = \delta_{ij}.$$

for all $1 \leq i, j \leq n$. In this paper all our commutators are graded, $[a, b] = ab - (-1)^{|a||b|}ba$. We set $C_0 = k$. These Clifford algebras have essentially only one nontrivial representation, sometimes called the *spinor representation* [17, p.14], defined as follows:

Definition 2.15. We use the \mathbb{Z}_2 -graded k -module

$$(2.7) \quad F_n = k\theta_1 \oplus \dots \oplus k\theta_n$$

where the θ_i are formal variables of odd degree, and set

$$(2.8) \quad S_n = \bigwedge F_n = \bigwedge (k\theta_1 \oplus \dots \oplus k\theta_n).$$

Definition 2.16. Left multiplication in the exterior algebra defines an odd operator

$$\begin{aligned} \theta_i \wedge (-) : S_n &\longrightarrow S_n, \\ \theta_{j_1} \dots \theta_{j_r} &\mapsto \theta_i \theta_{j_1} \dots \theta_{j_r}. \end{aligned}$$

Contraction from the left defines an odd operator

$$\begin{aligned} \theta_i^* \lrcorner (-) : S_n &\longrightarrow S_n, \\ \theta_{j_1} \dots \theta_{j_r} &\mapsto \sum_{l=1}^r (-1)^{(l-1)} \delta_{i,j_l} \theta_{j_1} \dots \widehat{\theta_{j_l}} \dots \theta_{j_r}. \end{aligned}$$

Usually we write simply θ_i for the operator $\theta_i \wedge (-)$ and θ_i^* for the operator $\theta_i^* \lrcorner (-)$. The spinor representation is defined by the next lemma, which also shows that the algebras C_n are Morita trivial in the sense that they are isomorphic to the endomorphism algebras of a finite rank free \mathbb{Z}_2 -graded k -module.

Lemma 2.17. *The map $C_n \longrightarrow \text{End}_k(S_n)$ defined by*

$$\gamma_i \mapsto \theta_i \wedge (-), \quad \gamma_i^\dagger \mapsto \theta_i^* \lrcorner (-)$$

is an isomorphism of \mathbb{Z}_2 -graded k -algebras.

In particular this means that every \mathbb{Z}_2 -graded C_n -module is isomorphic to $S_n \otimes_k V$ for some \mathbb{Z}_2 -graded k -module V . Using the isomorphism of k -modules

$$F_m \oplus F_n \cong F_{m+n}$$

which maps the ordered basis $\theta_1, \dots, \theta_m$ of F_m to the first m basis elements of F_{m+n} in their given order, and the ordered basis of F_n to the last n basis elements, we define

$$S_m \otimes_k S_n = \bigwedge F_m \otimes_k \bigwedge F_n \cong \bigwedge F_{m+n} = S_{m+n}.$$

From this we deduce an isomorphism of algebras

$$(2.9) \quad C_m \otimes_k C_n \cong C_{m+n}.$$

Given $m, n \geq 0$ we introduce notation for the C_m - C_n -bimodule

$$(2.10) \quad S_{m,n} = S_m \otimes_k S_n^* \cong \text{Hom}_k(S_n, S_m)$$

where $S_n^* = \text{Hom}_k(S_n, k)$. There is an isomorphism of C_l - C_n -bimodules

$$(2.11) \quad S_{l,m} \otimes_{C_m} S_{m,n} = S_l \otimes_k S_m^* \otimes_{C_m} S_m \otimes_k S_n^* \cong S_{l,n}.$$

Definition 2.18. For $n \geq 0$ let $e_n \in C_n$ denote the element

$$e_n = \gamma_1^\dagger \cdots \gamma_n^\dagger \gamma_n \cdots \gamma_1.$$

This is the idempotent corresponding to the summand $k \cdot 1$ of S_n .

Definition 2.19. More generally, given $m, n \geq 0$ consider the k -linear map

$$S_n \longrightarrow k \cdot 1 \xrightarrow{\text{inc}} S_m$$

given by the projection to $k \cdot 1$ followed by the inclusion into S_m . This defines an element

$$\iota_{m,n} \in \text{Hom}_k(S_n, S_m) = S_m \otimes_k S_n^*.$$

Then obviously $\iota_{n,n} = e_n$.

Remark 2.20. Recall that given a finite rank free k -module V and a symmetric bilinear form $B : V \otimes_k V \longrightarrow k$ the associated Clifford algebra $\text{Cl}(V, B)$ is the associative k -algebra generated by the elements of V subject to the relations

$$vw + wv = B(v, w) \cdot 1.$$

This is a \mathbb{Z}_2 -graded k -algebra when we assign $|v| = 1$ for all $v \in V$ and $|1| = 0$. If we take $V = F \oplus F^*$ with the bilinear form B under which $B(F, F) = 0, B(F^*, F^*) = 0$ and for $x \in F$ and $\nu \in F^*$, $B(x, \nu) = \frac{1}{2}\nu(x)$, then C_n is the Clifford algebra of the pair (V, B) .

2.5 The Clifford thickening

Let \mathcal{T} be a small idempotent complete supercategory. We construct a new supercategory \mathcal{T}^\bullet called the *Clifford thickening* in which the objects are pairs (X, n) of an integer $n \geq 0$ and a left C_n -module X in \mathcal{T} . Recall that if A is a \mathbb{Z}_2 -graded k -algebra then \mathcal{T}_A denotes the supercategory of A -modules in \mathcal{T} with A -linear maps.

Similarly to Example 2.8 there is a supercategory \mathcal{M} of Morita trivial \mathbb{Z}_2 -graded k -algebras: 1-morphisms are \mathbb{Z}_2 -graded bimodules which are finitely generated and projective over k and 2-morphisms are degree zero bimodule maps. We will need the operation of tensoring objects of \mathcal{T} with k -modules, which is recalled in Appendix A.

Proposition 2.21. *The assignment of the supercategory \mathcal{T}_A to an algebra A and of the superfunctor $\Phi_V = V \otimes_A (-)$ to a B - A -bimodule V determines a strong superfunctor*

$$\Phi^{\mathcal{T}} : \mathcal{M} \longrightarrow \text{Cat}_k^{\text{sup}}$$

to the supercategory of small supercategories and superfunctors.

Proof. The isomorphism $\Phi_W \circ \Phi_V \cong \Phi_{W \otimes V}$ is given by Lemma A.10, and apart from that the only checking that needs to be done are straightforward coherence diagrams. \square

Next we construct a strong functor into \mathcal{M} which picks out the Clifford algebras.

Definition 2.22. Let \mathbb{N} denote the category of integers $n \geq 0$ with a unique morphism $\phi_{m,n} : n \longrightarrow m$ for each pair m, n .

We view \mathbb{N} as a bicategory with only identity 2-morphisms.

Lemma 2.23. *There is a strong functor $\mathbb{N} \longrightarrow \mathcal{M}$ defined by*

$$\begin{aligned} n &\mapsto C_n = \text{End}_k(S_n), \\ \phi_{m,n} &\mapsto S_{m,n} = S_m \otimes_k S_n^*. \end{aligned}$$

The composite of these strong functors is a strong functor

$$(2.12) \quad \mathbb{N} \longrightarrow \mathcal{M} \xrightarrow{\Phi^{\mathcal{T}}} \text{Cat}_k^{\text{sup}}$$

sending n to the category of left C_n -modules in \mathcal{T} and $\phi_{m,n}$ to the functor $S_{m,n} \otimes_{C_n} -$. There is a close connection between strong functors into the bicategory of small categories, and what are called cofibered categories, given by the Grothendieck construction [30].

Definition 2.24. The *Clifford thickening* \mathcal{T}^\bullet of the supercategory \mathcal{T} is the category which results from the Grothendieck construction applied to the strong functor (2.12).

More concretely \mathcal{T}^\bullet is the category with objects all pairs (X, n) of an integer $n \geq 0$ and an object $X \in \mathcal{T}$ together with a left C_n -action. A morphism $\alpha : (X, n) \longrightarrow (Y, m)$ is a morphism of C_m -modules $\alpha : S_{m,n} \otimes_{C_n} X \longrightarrow Y$. Composition of a pair of morphisms

$$(2.13) \quad (X, n) \xrightarrow{\alpha} (Y, m) \xrightarrow{\beta} (Z, l)$$

is given by the morphism of C_l -modules

$$(2.14) \quad \begin{array}{c} S_{l,n} \otimes_{C_n} X \xrightarrow{\cong} (S_{l,m} \otimes_{C_m} S_{m,n}) \otimes_{C_n} X \\ \downarrow \cong \\ S_{l,m} \otimes_{C_m} (S_{m,n} \otimes_{C_n} X) \xrightarrow{1 \otimes \alpha} S_{l,m} \otimes_{C_m} Y \xrightarrow{\beta} Z. \end{array}$$

There is a canonical functor $p : \mathcal{T}^\bullet \longrightarrow \mathbb{N}$ defined by $p(X, n) = n$, which defines a cofibered category over \mathbb{N} by the definition of the Grothendieck construction. This definition has the benefit of being derived from a general construction, but there is an alternative description of morphisms which is less cumbersome:

Definition 2.25. Let \mathcal{T}^\star be the category with the same objects as \mathcal{T}^\bullet and

$$\mathcal{T}^\star((X, n), (Y, m)) = \left\{ \alpha \in \mathcal{T}(X, Y) \mid \alpha \gamma_i = 0 \text{ for } 1 \leq i \leq n \text{ and } \gamma_i^\dagger \alpha = 0 \text{ for } 1 \leq i \leq m \right\}.$$

Composition is as in \mathcal{T} , but the identity on (X, n) in \mathcal{T}^\star is the morphism

$$1_{(X,n)}^\star = e_n$$

where e_n is the action on X of the canonical idempotent from Definition 2.18.

Lemma 2.26. \mathcal{T}^\star is a category and there is an equivalence of categories

$$\Theta : \mathcal{T}^\bullet \longrightarrow \mathcal{T}^\star$$

Proof. To prove that \mathcal{T}^\star is a category it only needs to be checked that $1_{(X,n)}^\star$ really acts as an identity. Let $\alpha : X \longrightarrow Y$ be a morphism in \mathcal{T}^\star and consider

$$\begin{aligned} \alpha e_n &= \alpha \gamma_1^\dagger \cdots \gamma_n^\dagger \gamma_n \cdots \gamma_1 \\ &= \alpha \left(\gamma_1^\dagger \cdots \gamma_{n-1}^\dagger \gamma_{n-1} \cdots \gamma_1 - \gamma_1^\dagger \cdots \gamma_n \gamma_n^\dagger \cdots \gamma_1 \right) \\ &= \alpha \gamma_1^\dagger \cdots \gamma_{n-1}^\dagger \gamma_{n-1} \cdots \gamma_1 \end{aligned}$$

since in the second summand within the brackets, there is nothing to stop γ_n from commuting to the left and annihilating with α . Proceeding inductively, we see that $\alpha e_n = \alpha$. The proof that $e_n \alpha = \alpha$ is similar.

The functor Θ is defined to be the identity on objects, and given $\beta : (X, n) \longrightarrow (Y, m)$ in \mathcal{T}^\bullet which is a morphism $\beta : S_{m,n} \otimes_{C_n} X \longrightarrow Y$ in \mathcal{T} we define

$$\begin{aligned}\Theta(\beta) : X &\longrightarrow Y, \\ \Theta(\beta)(x) &= \beta(\iota_{m,n} \otimes x)\end{aligned}$$

where $\iota_{m,n}$ is as in Definition 2.19. We relegate the proof that this functor gives a bijection on morphism sets to Lemma A.13. Functoriality is easily checked, and $\Theta(1_X) = e_n$ since by definition $\iota_{n,n} = e_n$. \square

Every C_n -module X in \mathcal{T} is of the form $S_n \otimes_k V$ for some object V . It will be important later to know exactly how to extract the object V from X by splitting an idempotent.

Lemma 2.27. *Let X be a C_n -module in \mathcal{T} . The idempotent $e_n : X \longrightarrow X$ splits, say*

$$X \begin{array}{c} \xrightarrow{f_0} \\ \xleftarrow{g_0} \end{array} V$$

with $f_0 g_0 = 1_V$ and $g_0 f_0 = e_n$ and in \mathcal{T}^\bullet there are mutually inverse isomorphisms

$$(X, n) \begin{array}{c} \xrightarrow{f} \\ \xleftarrow{g} \end{array} (V, 0).$$

induced by f_0, g_0 . In particular

$$(2.15) \quad S_n^* \otimes_{C_n} X \cong V, \quad S_n \otimes_k V \cong X.$$

Proof. The map f_0 is a morphism $(X, n) \longrightarrow (V, 0)$ in \mathcal{T}^\bullet since

$$f_0 \gamma_i = f_0 e_n \gamma_i = f_0 \gamma_1^\dagger \cdots \gamma_n^\dagger \gamma_n \cdots \gamma_1 \gamma_i = 0.$$

Similarly g_0 defines a morphism $(V, 0) \longrightarrow (X, n)$. Let f, g be the morphisms in \mathcal{T}^\bullet corresponding to f_0, g_0 under the equivalence of Lemma 2.26. To prove that $fg = 1_{(X,n)}$ and $gf = 1_{(V,0)}$ in \mathcal{T}^\bullet it suffices to prove that $f_0 g_0 = 1_{(X,n)}^* = e_n$ and $g_0 f_0 = 1_{(V,0)}^*$ in \mathcal{T}^\bullet . But this is true by definition. \square

Lemma 2.28. *\mathcal{T}^\bullet is a supercategory and the functor $\iota : \mathcal{T} \longrightarrow \mathcal{T}^\bullet$ defined by $\iota(X) = (X, 0)$ is an equivalence of supercategories.*

Proof. The supercategory structure is given by the functor $\Psi : \mathcal{T}^\bullet \longrightarrow \mathcal{T}^\bullet$ where $\Psi(X, n) = (\Psi X, n)$ and ΨX has the C_n -action given by Definition A.8. The functor ι is fully faithful and it follows from Lemma 2.27 that it is essentially surjective. \square

3 The superbicategory \mathcal{C}

In this section we define a superbicategory without units \mathcal{C} . Later we will prove that this bicategory is equivalent to \mathcal{LG} , although this will not be apparent at first. The objects of \mathcal{C} are the same as \mathcal{LG} , namely potentials (x, W) .

Definition 3.1. Given potentials (x, W) and (y, V) we define

$$(3.1) \quad \mathcal{C}(W, V) = \left(\text{hmf}(k[x, y], V - W)^\omega \right)^\bullet,$$

where $(-)^\omega$ denotes the idempotent completion (see Remark 2.5) and $(-)^\bullet$ is the Clifford thickening (see Section 2.5).

Thus a 1-morphism $W \rightarrow V$ in \mathcal{C} is a finite rank matrix factorisation X of $V(y) - W(x)$ together with an idempotent endomorphism e and a family of odd operators $\gamma_i, \gamma_i^\dagger$ satisfying Clifford relations and the equations (all up to homotopy)

$$\begin{aligned} \gamma_i e &= \gamma_i = e \gamma_i, \\ \gamma_i^\dagger e &= \gamma_i^\dagger = e \gamma_i^\dagger. \end{aligned}$$

Most of the time we only care about 1-morphisms where $e = 1$, and let us only describe 2-morphisms in this case. For matrix factorisations X, Y of $V(y) - W(x)$ with respective Clifford actions $\{\gamma_i, \gamma_i^\dagger\}_{i=1}^a$ and $\{\rho_j, \rho_j^\dagger\}_{j=1}^b$ the 2-morphisms $\phi : (X, a) \rightarrow (Y, b)$ in \mathcal{C} are by Lemma 2.26 in bijection with morphisms of matrix factorisations $\phi : X \rightarrow Y$ satisfying

$$\phi \gamma_i = 0, \quad \rho_j^\dagger \phi = 0, \quad 1 \leq i \leq a, \quad 1 \leq j \leq b.$$

Observe that the homotopy category $\mathcal{LG}(W, V)_{fin}$ of finite rank matrix factorisations sits fully faithfully inside $\mathcal{C}(W, V)$ as the subcategory where all idempotents are identities and there are no Clifford actions, and by Lemma 2.28 the inclusion of this subcategory into $\mathcal{C}(W, V)$ is an idempotent completion.

The first aim of this section is to define, for any object (z, U) of \mathcal{LG} , a functor

$$\begin{aligned} \mathcal{C}(V, U) \otimes_k \mathcal{C}(W, V) &\rightarrow \mathcal{C}(W, U) \\ (Y, X) &\mapsto Y \mid X \end{aligned}$$

which we call the *cut functor*. The cut operation is defined on matrix factorisations X of $V - W$ and Y of $U - V$ as follows, assuming $y = (y_1, \dots, y_m)$ and writing

$$(3.2) \quad J_V = k[y]/(\partial_{y_1} V, \dots, \partial_{y_m} V).$$

Lemma 3.2. *The \mathbb{Z}_2 -graded $k[x, z]$ -module*

$$Y \mid X = Y \otimes_{k[y]} J_V \otimes_{k[y]} X$$

with the differential $d_Y \otimes 1 + 1 \otimes d_X$ is a finite rank matrix factorisation of $U - W$.

Proof. Since V is a potential J_V is a finite rank free k -module, and it follows that $Y | X$ is a finite rank free $k[x, z]$ -module. \square

Remark 3.3. If we choose a k -basis for J_V then multiplication by y_i , as a k -linear operator $J_V \rightarrow J_V$, gives a matrix $[y_i]$ over k . The cut $Y | X$ has a differential which, as a matrix, can be described by taking the matrix of $d_Y \otimes 1 + 1 \otimes d_X$ over $k[x, y, z]$ and replacing every occurrence of y_i by the scalar matrix $[y_i]$. This “inflation” produces a matrix of polynomials over $k[x, z]$.

Definition 3.4. Given morphisms of matrix factorisations $\phi : X \rightarrow X'$ and $\psi : Y \rightarrow Y'$ we define the morphism of matrix factorisations

$$\psi | \phi : Y | X \rightarrow Y' | X'$$

to be induced on the quotient by $\psi \otimes \phi : Y \otimes X \rightarrow Y' \otimes X'$. This is clearly functorial in the sense that for morphisms $\phi' : X' \rightarrow X''$ and $\psi' : Y' \rightarrow Y''$ we have

$$(3.3) \quad (\psi' | \phi') \circ (\psi | \phi) = (\psi' \psi) | (\phi' \phi),$$

and $1_Y | 1_X = 1_{Y | X}$.

The next step is to equip $Y | X$ with a Clifford action, which will be defined using the partial derivatives $\partial_{y_i}(d_X)$ and Atiyah classes. The Atiyah classes are defined by first taking the technical step of extending scalars

$$(3.4) \quad Y \check{\otimes} X = Y \otimes_{k[y]} k[[y]] \otimes_{k[y]} X$$

and putting a connection on this module. We write $t = (t_1, \dots, t_m)$ for the quasi-regular sequence of partial derivatives $t_i = \partial_{y_i} V$ and recall that there is by [15, Appendix B] a k -linear flat connection (which is “standard” in the sense of [15, Definition 8.6])

$$(3.5) \quad \nabla : k[[y]] \rightarrow k[[y]] \otimes_{k[t]} \Omega_{k[t]/k}^1$$

the components of which are k -linear operators $\partial_{t_i} : k[[y]] \rightarrow k[[y]]$. A simple example of such a connection is given in Section 4.5. Using a chosen homogeneous basis $\{e_a \otimes f_b\}_{a,b}$ for $Y \otimes X$ we may extend the operators ∂_{t_i} extend to $k[x, z]$ -linear operators

$$\begin{aligned} \partial_{t_i} : Y \check{\otimes} X &\rightarrow Y \check{\otimes} X \\ e_a \otimes h(y) \otimes f_b &\mapsto e_a \otimes \partial_{t_i}(h(y)) \otimes f_b. \end{aligned}$$

Lemma 3.5. For $1 \leq i \leq m$ the operator

$$(3.6) \quad [d_{Y \otimes X}, \partial_{t_i}] = d_{Y \otimes X} \partial_{t_i} - \partial_{t_i} d_{Y \otimes X} : Y \check{\otimes} X \rightarrow Y \check{\otimes} X$$

is $k[t]$ -linear and passes to a $k[x, z]$ -linear operator on $Y | X \cong (Y \check{\otimes} X) \otimes_{k[t]} k[t]/(t)$.

Proof. The fact that ∇ is a connection over $k[t]$ means that the operators ∂_{t_i} obey the Leibniz rule, that is

$$\partial_{t_i}(t_j h) = \partial_{t_i}(t_j)h + t_j \partial_{t_i} h = \delta_{ij}h + t_j \partial_{t_i} h$$

so $[\partial_{t_i}, t_j] = \delta_{ij}$. We calculate

$$[[d_{Y \otimes X}, \partial_{t_i}], t_j] = -[[t_j, d_{Y \otimes X}], \partial_{t_i}] - [[\partial_{t_i}, t_j], d_{Y \otimes X}] = -[\delta_{ij}, d_{Y \otimes X}] = 0$$

which shows that $[d_{Y \otimes X}, \partial_{t_i}]$ is $k[t]$ -linear. \square

Definition 3.6. The i th Atiyah class is this odd $k[x, z]$ -linear operator

$$\text{At}_i^{Y, X} = [d_{Y \otimes X}, \partial_{t_i}] : Y | X \longrightarrow Y | X .$$

Usually we just write At_i where it will not cause confusion.

This operator is the i th Atiyah class of $Y \check{\otimes} X$ relative to the ring morphism $k[x, y] \longrightarrow k[x, y, t]$ as defined in [15, Section 9]. A reference for Atiyah classes is [4], but we will prove the necessary properties here for the reader's convenience.

Lemma 3.7. *The operator At_i on $Y | X$ is closed, and if $\phi : X \longrightarrow X'$ and $\psi : Y \longrightarrow Y'$ are morphisms of matrix factorisations then there are $k[x, z]$ -linear homotopies*

$$(3.7) \quad \text{At}_i^{Y', X'} \circ (\psi | \phi) \simeq (\psi | \phi) \circ \text{At}_i^{Y, X} .$$

That is, the Atiyah classes are natural.

Proof. The Atiyah class is closed since

$$[d, \text{At}_i] = [d, [d, \partial_{t_i}]] = 0 .$$

If $\kappa : Y \otimes X \longrightarrow Y' \otimes X'$ is any $k[x, y, z]$ -linear even closed operator,

$$[\kappa, [d, \partial_{t_i}]] + [\partial_{t_i}, [\kappa, d]] + [d, [\partial_{t_i}, \kappa]] = 0$$

from which we deduce that $[\partial_{t_i}, \kappa]$ is a homotopy establishing (3.7). \square

The other ingredient in defining the Clifford action are the partial derivatives $\partial_{y_i}(d_X)$, as discussed in Section 2.3. On $Y \otimes X$ we have the odd operator $\partial_{y_i}(d_X) = 1 \otimes \partial_{y_i}(d_X)$ which is a homotopy for the action of $\partial_{y_i}(V - W) = \partial_{y_i}(V)$. Hence when we pass to the quotient, $\partial_{y_i}(d_X)$ is a closed odd operator on $Y | X$.

Definition 3.8. On $Y | X$ we define odd $k[x, z]$ -linear closed operators

$$(3.8) \quad \gamma_i^\dagger = \text{At}_i, \quad \gamma_i = -\partial_{y_i}(d_X) - \frac{1}{2} \sum_q \partial_{y_q y_i}(V) \text{At}_q .$$

For a concrete example, see Section 4.5.

Theorem 3.9. *Given matrix factorisations Y, X as above, the data*

$$(3.9) \quad Y | X = \left(Y \otimes_{k[y]} J_V \otimes_{k[y]} X, \{\gamma_i, \gamma_i^\dagger\}_{i=1}^m \right)$$

defines a representation of C_m in the homotopy category of matrix factorisations, that is, the $\gamma_i, \gamma_i^\dagger$ satisfy the Clifford relations (2.6) up to homotopy.

Proof. This will follow from the considerations of Section 4.3 but we feel it is worthwhile to give a direct proof, both to demonstrate how straightforward it is and in order to have written down explicit homotopies for the various relations.

By the graded Jacobi identity

$$\begin{aligned} [\text{At}_i, \text{At}_j] &= [\text{At}_i, [d, \partial_{t_j}]] \\ &= -[\partial_{t_j}, [\text{At}_i, d]] - [d, [\partial_{t_j}, \text{At}_i]] \\ &= [d, -[\partial_{t_j}, \text{At}_i]]. \end{aligned}$$

Hence $[\gamma_i^\dagger, \gamma_j^\dagger] = [d, h_{ij}]$ where $h_{ij} = -[\partial_{t_j}, \text{At}_i]$. Next we compute

$$\begin{aligned} [\partial_{y_i}(d_X), \text{At}_j] &= [\partial_{y_i}(d_X), [d, \partial_{t_j}]] \\ &= -[\partial_{t_j}, [\partial_{y_i}(d_X), d]] + [d, [\partial_{t_j}, \partial_{y_i}(d_X)]] \\ &= -[\partial_{t_j}, t_i] + [d, [\partial_{t_j}, \partial_{y_i}(d_X)]] \\ &= -\delta_{ij} + [d, [\partial_{t_j}, \partial_{y_i}(d_X)]]. \end{aligned}$$

Again we set $g_{ij} = [\partial_{t_j}, \partial_{y_i}(d_X)]$ for later use. We compute that

$$\begin{aligned} [\gamma_i, \gamma_j^\dagger] &= \left[-\partial_{y_i}(d_X) - \frac{1}{2} \sum_q \partial_{y_q y_i}(V) \text{At}_q, \text{At}_j \right] \\ &= \delta_{ij} - [d, g_{ij}] - \frac{1}{2} \sum_q \partial_{y_q y_i}(V) [\text{At}_q, \text{At}_j] \\ &= \delta_{ij} - [d, g_{ij}] - \frac{1}{2} \sum_q \partial_{y_q y_i}(V) [d, h_{qj}] \\ &= \delta_{ij} + [d, -g_{ij} - \frac{1}{2} \sum_q \partial_{y_q y_i}(V) h_{qj}]. \end{aligned}$$

Finally, using Lemma 2.14

$$\begin{aligned}
[\gamma_i, \gamma_j] &= \left[\partial_{y_i}(d_X) + \frac{1}{2} \sum_q \partial_{y_q y_i}(V) \text{At}_q, \partial_{y_j}(d_X) + \frac{1}{2} \sum_q \partial_{y_q y_j}(V) \text{At}_q \right] \\
&= [\partial_{y_i}(d_X), \partial_{y_j}(d_X)] + \frac{1}{2} \sum_q \partial_{y_q y_j}(V) [\partial_{y_i}(d_X), \text{At}_q] \\
&\quad + \frac{1}{2} \sum_q \partial_{y_q y_i}(V) [\text{At}_q, \partial_{y_j}(d_X)] \\
&\quad + \frac{1}{4} \sum_{p,q} \partial_{y_p y_i}(V) \partial_{y_q y_j}(V) [\text{At}_p, \text{At}_q] \\
&= \partial_{y_i y_j}(V) + [d, -\partial_{y_j y_i}(d_X)] \\
&\quad + \frac{1}{2} \sum_q \partial_{y_q y_j}(V) (-\delta_{iq} + [d, [\partial_{t_q}, \partial_{y_i}(d_X)]]) \\
&\quad + \frac{1}{2} \sum_q \partial_{y_q y_i}(V) (-\delta_{jq} + [d, [\partial_{t_q}, \partial_{y_j}(d_X)]]) \\
&\quad + \frac{1}{4} \sum_{p,q} \partial_{y_p y_i}(V) \partial_{y_q y_j}(V) [d, -[\partial_{t_q}, \text{At}_p]] \\
&= [d, c_{ij}]
\end{aligned}$$

where

$$\begin{aligned}
c_{ij} &= -\partial_{y_j y_i}(d_X) + \frac{1}{2} \sum_q \partial_{y_q y_j}(V) [\partial_{t_q}, \partial_{y_i}(d_X)] \\
&\quad + \frac{1}{2} \sum_q \partial_{y_q y_i}(V) [\partial_{t_q}, \partial_{y_j}(d_X)] \\
&\quad - \frac{1}{4} \sum_{p,q} \partial_{y_p y_i}(V) \partial_{y_q y_j}(V) [\partial_{t_q}, \text{At}_p].
\end{aligned}$$

□

Lemma 3.10. *Given morphisms $\phi : X \longrightarrow X', \psi : Y \longrightarrow Y'$, the morphism of matrix factorisations $\psi| \phi$ is a morphism of C_m -representations in the homotopy category.*

Proof. This follows easily from the fact that Atiyah classes are natural (Lemma 3.7) and the homotopies $\partial_{y_i}(d_X)$ are central (Lemma 2.13). □

Proposition 3.11. *The cut operation defines a functor*

$$(3.10) \quad \mathcal{C}(V, U) \otimes_k \mathcal{C}(W, V) \longrightarrow \mathcal{C}(W, U)$$

Proof. What we have done so far in this section defines a functor

$$\mathrm{hmf}(k[y, z], U - V) \otimes \mathrm{hmf}(k[x, y], V - W) \longrightarrow \mathrm{hmf}(k[x, z], U - W)^{\omega^\bullet}$$

and since the latter category is idempotent complete, this lifts to a functor

$$\mathrm{hmf}(k[y, z], U - V)^\omega \otimes \mathrm{hmf}(k[x, y], V - W)^\omega \longrightarrow \mathrm{hmf}(k[x, z], U - W)^{\omega^\bullet}.$$

Explicitly, this functor is given for $(Y, e_Y) \in \mathcal{C}(V, U)$ and $(X, e_X) \in \mathcal{C}(W, V)$ by

$$(Y, e_Y) \mid (X, e_X) = (Y \mid X, e_Y \mid e_X).$$

Now suppose we have representations $(Y, \{\rho_j, \rho_j^\dagger\}_{j=1}^r)$ of C_r in $\mathcal{C}(V, U)$ and $(X, \{\tau_l, \tau_l^\dagger\}_{l=1}^s)$ of C_s in $\mathcal{C}(W, V)$. Here Y, X may come equipped with idempotents but these are easily handled so we ignore them in what follows. Since the cut operation is functorial, we have representations of C_r and C_s on $Y \mid X$ given by

$$\{\rho_j \mid 1_X, \rho_j^\dagger \mid 1_X\}_{j=1}^r, \quad \{1_Y \mid \tau_l, 1_Y \mid \tau_l^\dagger\}_{l=1}^s.$$

The operators from the action of C_r anticommute with the operators from the action of C_s because they act on different components of a tensor product. Moreover, both actions anticommute with the new C_m action $\{\gamma_i, \gamma_i^\dagger\}_{i=1}^m$ coming from the cut, by Lemma 3.10. Thus $Y \mid X$ is equipped with the structure of a $C_r \otimes C_m \otimes C_s \cong C_{r+m+s}$ -representation. \square

Definition 3.12. A *(super)bicategory without units* has all the data and satisfies all the conditions of a (super)bicategory, with the exception of the existence of unit 1-morphisms, unitors, and their coherence diagram.

Definition 3.13. The superbicategory without units \mathcal{C} has

- **Objects:** same as \mathcal{LG} , that is, potentials (x, W) .
- **1- and 2-morphisms:** defined by (3.1).
- **Composition:** defined by the cut operation (3.10).
- **Associators:** given a triple of composable 1-morphisms

$$(3.11) \quad (q, Q) \xleftarrow{Z} (y, U) \xleftarrow{Y} (z, V) \xleftarrow{X} (x, W) .$$

with $V \in k[z_1, \dots, z_m]$ and $U \in k[y_1, \dots, y_p]$ the isomorphism

$$\begin{aligned} \alpha : Z \mid (Y \mid X) &\longrightarrow (Z \mid Y) \mid X, \\ z \otimes (y \otimes x) &\mapsto (z \otimes y) \otimes x \end{aligned}$$

is $C_m \otimes C_p$ -linear (see Lemma 3.14 below) and is the associator for \mathcal{C} .

Lemma 3.14. *The morphism $\alpha : (Z | Y) | X \longrightarrow Z | (Y | X)$ is $C_m \otimes C_p$ -linear.*

Proof. We prove the C_p -linearity of α , the proof for C_m is similar. The algebra C_p acts on the cut between Z and Y involving the variables y_1, \dots, y_p . The action on $Z | (Y | X) = Z \otimes J_U \otimes (Y \otimes J_V \otimes X)$ is via $1 \otimes (\partial_{y_i}(d_Y) \otimes 1)$ and the Atiyah classes of

$$M = Z \otimes_{k[y]} k[[y]] \otimes_{k[y]} (Y \otimes_{k[z]} J_V \otimes_{k[z]} X)$$

relative to the ring morphism $k[q, x] \longrightarrow k[q, x][\partial_{y_1}U, \dots, \partial_{y_p}U]$. Specifically, the generator γ_i^\dagger of C_p is the commutator $[d_M, \partial_s]$ where $s = \partial_{y_i}U$. But this operator can be identified with an Atiyah class of the linear factorisation

$$(3.12) \quad Z \otimes_{k[y]} k[[y]] \otimes_{k[y]} (Y \otimes_{k[z]} k[[z]] \otimes_{k[z]} X)$$

relative to the ring morphism

$$(3.13) \quad k[q, x] \longrightarrow k[q, x][\partial_{y_1}U, \dots, \partial_{y_p}U, \partial_{z_1}V, \dots, \partial_{z_m}V].$$

By the same argument the action of C_p on $(Z | Y) | X$ is via $(1 \otimes \partial_{y_i}(d_Y)) \otimes 1$ and the Atiyah classes of $(Z \otimes_{k[y]} k[[y]] \otimes_{k[y]} Y) \otimes_{k[z]} k[[z]] \otimes_{k[z]} X$ relative to the same ring morphism (3.13). It is therefore clear that the associator α identifies these two actions. \square

Proposition 3.15. *Thus defined \mathcal{C} is a superbicategory without units.*

Proof. The only thing to check is the coherence of the associator, which is trivial. \square

Remark 3.16. The cut system presented here refines earlier work in [15]. To see the connection, recall the idempotent $e_m \in C_m$ of Definition 2.18 which corresponds to projection onto lowest degree in S_m . The projector onto top degree $k \cdot \theta_1 \cdots \theta_m \subset S_m$ is

$$e'_m = \gamma_1 \cdots \gamma_m \gamma_m^\dagger \cdots \gamma_1^\dagger.$$

Observe that each of the γ_i 's acts as $-\lambda_i$ because all of the γ_j^\dagger 's to the right. Hence

$$\begin{aligned} e'_m &= (-1)^{\binom{m}{2}} \gamma_1 \cdots \gamma_m \gamma_1^\dagger \cdots \gamma_m^\dagger \\ &= (-1)^{\binom{m+1}{2}} \lambda_1 \cdots \lambda_m \text{At}_1 \cdots \text{At}_m. \end{aligned}$$

which is precisely the idempotent which appears in [15, Corollary 10.4].

In *ibid.* we constructed finite models of convolutions $Y \circ X$ one pair (Y, X) at time. To make this assignment of finite models coherently for all 1-morphisms in \mathcal{LG} simultaneously it is natural to identify the Clifford actions which underlie the idempotents, as we do here.

Remark 3.17. There is a question of whether or not the C_m -action on $Y | X$ is canonical, since it depends on the choice of connection ∇ and on a choice of homogeneous basis for X and Y . But Atiyah classes are independent, up to homotopy, of the choice of connection.

4 The equivalence of \mathcal{C} and \mathcal{LG}

We prove that the cut functor defined in Section 3 models composition in \mathcal{LG} by providing an equivalence of this superbicategory with \mathcal{C} . This is done by presenting the cut $Y \mid X$ with its Clifford action as the solution of the problem of finding a finite model for the matrix factorisation $Y \otimes X$ over $k[x, z]$.

As was mentioned in Section 2.2, $Y \otimes X$ is a matrix factorisation of infinite rank over $k[x, z]$. The obvious notion of a “finite model” would be a finite rank matrix factorisation to which $Y \otimes X$ is homotopy equivalent, but providing such a finite model generically seems impossible. We propose a different notion of finite model based on Clifford representations, and solve the problem in this setting. The solution we give here builds on earlier progress on this problem with Dyckerhoff [15].

Setup 4.1. *The situation is as follows:*

- $(x, W), (y, V), (z, U)$ are objects of \mathcal{LG} with $k[y] = k[y_1, \dots, y_m]$.
- Y is a finite rank matrix factorisation of $U - V$ over $k[y, z]$.
- X is a finite rank matrix factorisation of $V - W$ over $k[x, y]$.

The strategy is to consider the direct sum of (shifted) copies

$$(4.1) \quad S_m \otimes_k (Y \otimes X) = (Y \otimes X) \oplus (Y \otimes X)[1] \oplus (Y \otimes X) \oplus \dots$$

where S_m is the exterior algebra viewed as a \mathbb{Z}_2 -graded k -module

$$S_m = \bigwedge (k[1]^{\oplus m}).$$

We find a finite model of this larger object, and in addition record a method of extracting the subobject $Y \otimes X$. Some necessary background is presented in Section 4.1 and Section 4.2, and finally the full construction is presented in Section 4.3. The overall strategy is:

Strategy 4.2. We show there is a differential d_K (K stands for Koszul) such that

- 1) The complex $(S_m \otimes_k (Y \otimes X), d_K)$ has a finite model: there is a homotopy equivalence

$$(4.2) \quad (Y \mid X, 0) \cong (S_m \otimes_k (Y \otimes X), d_K)$$

over $k[x, z]$ where $Y \mid X$ is the graded module from Section 3.

- 2) The perturbation lemma (Section 4.1) promotes this to a homotopy equivalence

$$(4.3) \quad (Y \mid X, d_{Y \otimes X}) \cong (S_m \otimes_k (Y \otimes X), d_K + d_{Y \otimes X})$$

by mixing in the differential $d_{Y \otimes X}$ on both sides.

3) Using Section 4.2 we show there is a homotopy equivalence

$$(4.4) \quad (S_m \otimes_k (Y \otimes X), d_K + d_{Y \otimes X}) \cong (S_m \otimes_k (Y \otimes X), d_{Y \otimes X}).$$

4) Combining (4.3) and (4.4) we have the desired finite model

$$(4.5) \quad (Y \mid X, d_{Y \otimes X}) \cong (S_m \otimes_k (Y \otimes X), d_{Y \otimes X}).$$

There is a canonical action of the Clifford algebra C_m on S_m and thus on the right hand side, which “remembers” how to extract $Y \otimes X$ from this larger object. Transferring this action yields the operators $\{\gamma_i, \gamma_i^\dagger\}_{i=1}^m$ on $Y \mid X$ from Section 3.

In conclusion, there is an isomorphism of Clifford representations (4.5) in the homotopy category of finite rank matrix factorisations of $U - W$ over $k[x, z]$, and this is our solution to the problem of finding a finite model of $Y \otimes X$. Once we have established all of this, proving that \mathcal{C} and \mathcal{LG} are equivalent bicategories will be straightforward.

4.1 Homological perturbation and fixed points

A fundamental role in this paper is played by the homological perturbation lemma. While this result is usually stated for complexes the standard results generalise immediately to linear factorisations, and in this section we collect these standard results. But we begin by recalling the motivating problem for which the perturbation lemma is the solution.

Let Q be a complex of vector spaces (possibly infinite dimensional) with finite dimensional cohomology, so that Q contains in some sense only a “finite” amount of information. One way of making this finiteness manifest is to simply list the cohomology groups of Q , but a more flexible and categorical solution is to find a finite representative in the homotopy equivalence class of Q .

A *deformation retract* of complexes is a pair of morphisms

$$(4.6) \quad C \begin{array}{c} \xleftarrow{\pi} \\ \xrightarrow{\sigma} \end{array} Q, \quad \phi$$

and a degree -1 operator $\phi : Q \rightarrow Q$ satisfying $\pi\sigma = 1_C, \sigma\pi = 1_Q - [d_Q, \phi]$ so that π, σ are mutually inverse homotopy equivalences. A deformation retract is *strong* if it satisfies some additional conditions, listed below. The point is that if C is finitely generated, then (4.6) gives the desired finite representative in the homotopy equivalence class of Q .

Example 4.3. Consider the following diagram of complexes of k -vector spaces

$$\begin{array}{ccc} 0 & \begin{array}{c} \xleftarrow{\quad} \\ \xrightarrow{\quad} \end{array} & k[x] \\ \downarrow & & \downarrow x \\ k & \begin{array}{c} \xleftarrow{\pi} \\ \xrightarrow{\sigma} \end{array} & k[x] \end{array} \quad \begin{array}{c} \swarrow \phi \\ \searrow \phi \end{array}$$

in which the first column is the complex C which has k concentrated in degree zero, and the second column Q is the Koszul complex on x . The map $\pi : k[x] \longrightarrow k$ sends every polynomial to its constant term, while $\sigma : k \longrightarrow k[x]$ is the inclusion of the constants. The operator ϕ is defined by $\phi(x^n) = x^{n-1}$ and $\phi(1) = 0$. Together these maps are a deformation retract giving k as the “finite model” of Q .

Suppose now that a deformation retract computing a finite representative for Q cannot be found immediately, but that we know how to decompose the differential as $d_Q = d + \tau$ in such a way that there is a finite representative for the complex (Q, d) . The perturbation lemma allows us, with some conditions, to “mix in” the operator τ to obtain a finite model for the original complex Q .

Let R be a commutative ring and $W \in R$. Everything we say applies, in the case where $W = 0$, to both \mathbb{Z}_2 -graded and \mathbb{Z} -graded complexes. For our purposes the reformulation of the perturbation lemma in terms of fixed points by Barnes and Lambe is more natural, so we emphasise the “splitting homotopies” of [1].

Definition 4.4. A *splitting homotopy* on a linear factorisation (A, d) of W is a degree -1 operator ϕ on A satisfying

- (i) $\phi^2 = 0$,
- (ii) $\phi d \phi = \phi$.

A *morphism* $(A, d, \phi) \longrightarrow (A', d', \phi')$ of splitting homotopies is a morphism $\alpha : A \longrightarrow A'$ of linear factorisations satisfying $\phi' \alpha = \alpha \phi$.

The condition (ii) says that ϕ is a *fixed point* of the operator $F(x) = xdx$.

Definition 4.5. A *strong deformation retract* of linear factorisations of W is a pair of morphisms of linear factorisations

$$(4.7) \quad (M, d) \begin{array}{c} \xleftarrow{\pi} \\ \xrightarrow{\sigma} \end{array} (A, d), \quad \phi$$

together with a degree -1 operator $\phi : A \longrightarrow A$ satisfying

- (i) $\pi \sigma = 1_M$,
- (ii) $\sigma \pi = 1_A - [d, \phi]$,
- (iii) $\phi^2 = 0$,
- (iv) $\phi \sigma = 0$,
- (v) $\pi \phi = 0$.

A *morphism* of strong deformation retracts is a commutative diagram

$$\begin{array}{ccc} (M, d) & \begin{array}{c} \xleftarrow{\pi} \\ \xrightarrow{\sigma} \end{array} & (A, d) \\ \alpha \downarrow & & \downarrow \beta \\ (M', d') & \begin{array}{c} \xleftarrow{\pi'} \\ \xrightarrow{\sigma'} \end{array} & (A', d') \end{array}$$

that is, a pair of morphisms α, β such that $\beta\sigma = \sigma'\alpha$, $\alpha\pi = \pi'\beta$ and $\phi'\beta = \beta\phi$.

We follow the conventions of [1] in defining strong deformation retracts; up to a sign this the same as the special deformation retracts of [12] and [15]. Namely, a strong deformation retract in the above sense is the same as a special deformation retract in the sense of *ibid.* with $h = -\phi$.

Lemma 4.6. *There is an equivalence between the category of splitting homotopies \mathcal{C}_1 and the category of strong deformation retracts \mathcal{C}_2 .*

Proof. We briefly recall the construction from [1, p.883], which is not stated in terms of categories but is obviously functorial. Given a strong deformation retract (4.7) the data (A, ϕ) is a splitting homotopy. In the reverse direction, if (A, ϕ) is a splitting homotopy then $e = 1_A - [d, \phi]$ is idempotent and we define the linear factorisation $M = \text{im}(e)$ with the associated projection π and inclusion σ . Then this is a strong deformation retract, together with the original ϕ . \square

Here is the problem that the formalism is designed to solve: suppose (A, d) is a linear factorisation of W and that τ is a degree +1 operator on A such that $(A, d + \tau)$ is a linear factorisation of V (possibly different to W). This τ is called the *perturbation*.

Definition 4.7. Given a splitting homotopy ϕ on (A, d) the *transference problem* is to find a splitting homotopy ϕ' on $(A, d + \tau)$ such that $\text{Im}(e) \cong \text{Im}(e')$ as graded R -modules, where $e = 1_A - [d, \phi]$ and $e' = 1_A - [d + \tau, \phi']$.

That is, with $\xi = d + \tau$ the problem is to find a fixed point of the operator

$$F(x) = x\xi x$$

among operators with $x^2 = 0$ satisfying the boundary condition $\text{Im}(1_A - [d, x]) \cong \text{Im}(e)$. If $\phi\tau$ has finite order, i.e. $(\phi\tau)^m = 0$ for some m , then

$$\phi_\infty = \sum_{m \geq 0} (-1)^m (\phi\tau)^m \phi = \phi - \phi\tau\phi + \phi\tau\phi\tau\phi - \dots$$

is a solution of this fixed point problem:

Theorem 4.8 (Perturbation lemma). *Suppose $\phi\tau$ has finite order. Then ϕ_∞ is a splitting homotopy on the linear factorisation (A, ξ) and satisfies the isomorphism condition in the transference problem.*

Proof. The proof for complexes [1, p.886] goes through unchanged for linear factorisations, with some minor modifications that are elaborated in [7, §2.5]. \square

The more common statement of the basic perturbation lemma involves the deformation retract corresponding to the splitting homotopy ϕ_∞ under the equivalence of Lemma 4.6. The statement is that given a splitting homotopy ϕ as above with associated deformation retract (4.7) there is a strong deformation retract

$$(4.8) \quad (M, d_\infty) \xrightleftharpoons[\sigma_\infty]{\pi_\infty} (A, \xi), \quad \phi_\infty$$

where $A = \tau(1 + \phi\tau)^{-1} = \sum_{m \geq 0} (-1)^m \tau(\phi\tau)^m$ and

$$\begin{aligned} \sigma_\infty &= \sigma - \phi A \sigma, \\ \pi_\infty &= \pi - \pi A \phi, \\ d_\infty &= d + \pi A \sigma. \end{aligned}$$

As explained in [1], to derive these formulas one has only to notice that the sub-complex in the deformation retract associated to ϕ_∞ can, by the isomorphism

$$\text{Im}(1 - [\xi, \phi_\infty]) \cong \text{Im}(e) = M$$

be identified with M , and the induced maps are the $\sigma_\infty, \pi_\infty, d_\infty$ given above. It is also possible to give a direct proof as in [12].

4.2 Clifford actions and Koszul complexes

Let R be a k -algebra and (X, d) a complex of R -modules on which $t_1, \dots, t_m \in R$ act null-homotopically. Tensoring X with the Koszul complex $K(t_i)$ yields the same complex as taking the cone of $t_i \cdot 1_X = 0 : X \rightarrow X$, and therefore we have a homotopy equivalence

$$(4.9) \quad K(t_i) \otimes_R X \cong \text{cone}(t_i \cdot 1_X) \cong X \oplus X[1] \cong (k \oplus k[1]) \otimes_k X.$$

Iterating this operation we obtain a homotopy equivalence with $K = K(t_1, \dots, t_m)$,

$$(4.10) \quad K \otimes_R X \cong \bigwedge (k[1]^{\oplus m}) \otimes_k X$$

where the left hand side is a complex with differential $d_K + d$ and the right hand side has only the differential d . On the right hand side we have an obvious action of

$$(4.11) \quad C_m = \text{End}_k \left(\bigwedge (k[1]^{\oplus m}) \right)$$

and the purpose of this section is to compute the corresponding action of C_m via closed odd endomorphisms of the complex $K \otimes_R X$. In fact we will do this more generally:

Setup 4.9. Let R be a k -algebra and write \otimes for \otimes_R . Let (X, d) be a linear factorisation of $W \in R$ and t_1, \dots, t_m a sequence of elements of R acting null-homotopically on X . We choose odd R -linear operators λ_i on X with $[\lambda_i, d] = t_i \cdot 1_X$ for $1 \leq i \leq m$.

Remark 4.10. All equalities in this section mean equalities of linear maps. If we mean homotopy, we will explicitly write \simeq .

We introduce formal variables $\theta_1, \dots, \theta_m$ of odd degree and set

$$S_m = \bigwedge (k\theta_1 \oplus \dots \oplus k\theta_m) .$$

On this \mathbb{Z} -graded k -module there are canonical odd k -linear operators θ_i, θ_i^* defined as in Definition 2.16. The Koszul complex K of the sequence t_1, \dots, t_m is defined by

$$(4.12) \quad K = S_m \otimes_k R, \quad d_K = \sum_{i=1}^m t_i \theta_i^* .$$

The tensor products $K \otimes X = (K \otimes X, d + d_K)$ and $S_m \otimes_k X = (S_m \otimes_k X, d)$ are both linear factorisations of W with isomorphic underlying graded modules. On this underlying graded module $K \otimes X \cong S_m \otimes_k X$ we have the odd R -linear operators $\theta_k^* = \theta_k^* \otimes 1$ and $\theta_k = \theta_k \otimes 1$ and we introduce the following even R -linear operators

$$\begin{aligned} \delta &= \sum_{i=1}^m \lambda_i \theta_i^*, \\ \exp(-\delta) &= \sum_{n \geq 0} (-1)^n \frac{1}{n!} \delta^n . \end{aligned}$$

Observe that δ is nilpotent so the exponential makes sense.

Proposition 4.11. *There are mutually inverse isomorphisms of linear factorisations*

$$(4.13) \quad K \otimes X \begin{array}{c} \xrightarrow{\exp(\delta)} \\ \xleftarrow{\exp(-\delta)} \end{array} S_m \otimes_k X$$

where the left hand side has differential $d + d_K$ and the right hand side has d .

We break the proof into a series of lemmas:

Lemma 4.12. $[d, \delta^n] = n\delta^{n-1}d_K$ for $n \geq 1$.

Proof. The case of $n = 1$ is clear

$$[d, \delta] = \sum_i [d, \lambda_i] \theta_i^* = d_K .$$

For $n > 1$ we use the case $n = 1$ to show that

$$[d, \delta^n] = \sum_{i=0}^{n-1} \delta^i [d, \delta] \delta^{n-i-1} = \sum_{i=0}^{n-1} \delta^i d_K \delta^{n-i-1} = \sum_{i=0}^{n-1} \delta^i \delta^{n-i-1} d_K = n\delta^{n-1} d_K .$$

□

Lemma 4.13. $[d, \exp(-\delta)] = -\exp(-\delta)d_K$ and $[d, \exp(\delta)] = \exp(\delta)d_K$.

Proof. We prove the first identity using Lemma 4.12, the second is the same:

$$\begin{aligned}
[d, \exp(-\delta)] &= \sum_{n \geq 1} \frac{(-1)^n}{n!} [d, \delta^n] \\
&= \sum_{n \geq 1} \frac{(-1)^n}{n!} n \delta^{n-1} d_K \\
&= - \sum_{n \geq 0} \frac{(-1)^n}{n!} \delta^n d_K \\
&= -\exp(-\delta)d_K
\end{aligned}$$

□

Proof of Proposition 4.11. It suffices to show $\exp(-\delta)$ is a morphism of linear factorisations. But using Lemma 4.13

$$(d + d_K) \exp(-\delta) - \exp(-\delta)d = [d, \exp(-\delta)] + d_K \exp(-\delta) = 0$$

and similarly for the other equation. □

Definition 4.14. Let γ be a homogeneous R -linear operator on $S_m \otimes_k X$. We define the *transfer* $\mathcal{T}(\gamma)$ to be the homogeneous R -linear operator on $K \otimes X$ induced by γ using the equivalence (4.13), that is,

$$(4.14) \quad \mathcal{T}(\gamma) = \exp(-\delta)\gamma \exp(\delta).$$

Remark 4.15. We do not necessarily require that γ be closed, but since

$$(4.15) \quad [\mathcal{T}(\gamma), d_K + d] = \mathcal{T}([\gamma, d])$$

if γ is a closed operator then so is $\mathcal{T}(\gamma)$.

Of particular interest are the transfers of the generators θ_i, θ_i^* of the Clifford action. The transfer of the contraction operator θ_i^* is easy:

Lemma 4.16. $\mathcal{T}(\theta_i^*) = \theta_i^*$.

Proof. Since θ_i^* commutes with δ . □

However the calculation of $\mathcal{T}(\theta_i)$ is much more involved. To compute $\mathcal{T}(\gamma)$ the strategy is to try and commute γ past $\exp(-\delta)$ and see what happens. To this end we must first compute $[\gamma, \delta^m]$. We begin the calculations with some general statements about graded commutators.

To this end let A be a \mathbb{Z} -graded \mathbb{Q} -algebra and a, b, c homogeneous elements.

Lemma 4.17. *We have*

$$(4.16) \quad [ab, c] = a[b, c] + (-1)^{|b||c|}[a, c]b,$$

$$(4.17) \quad [a, bc] = [a, b]c + (-1)^{|a||b|}b[a, c].$$

We adopt the following notation for iterated commutators

Definition 4.18. The iterated commutator $[a, b]^{(n)}$ is defined by $[a, b]^{(0)} = b$ and

$$[a, b]^{(n+1)} = [a, [a, b]^{(n)}].$$

Observe that $[a, b]^{(n)}$ is a linear function of b , but not a .

Lemma 4.19. *For $n \geq 1$ we have*

$$[a, b^n] = \sum_{i=0}^{n-1} (-1)^{i|a||b|} b^i [a, b] b^{n-i-1}.$$

Proof. Easy proof by induction using (4.17). □

Lemma 4.20. *For $n \geq 1$ we have*

$$(4.18) \quad a^n [a, b] = \sum_{\substack{0 \leq i, j \leq n \\ i+j=n}} \binom{n}{i} [a, b]^{(i+1)} a^j.$$

Proof. By induction, with the case $n = 1$ being clear. Using the inductive hypothesis

$$a^{n+1} [a, b] = \sum_{i+j=n} \binom{n}{i} a [a, b]^{(i+1)} a^j = \sum_{i+j=n} \binom{n}{i} \left[[a, b]^{(i+2)} a^j + [a, b]^{(i+1)} a^{j+1} \right].$$

The powers of a range from 0 to $n + 1$ and the coefficient of a^j for $1 \leq j \leq n$ is

$$\binom{n}{n-j} [a, b]^{(n-j+2)} + \binom{n}{n-j+1} [a, b]^{(n-j+2)} = \binom{n+1}{n+1-j} [a, b]^{(n-j+2)}.$$

The coefficient of a^0 is $[a, b]^{(n+2)}$ and the coefficient of a^{n+1} is $[a, b]^{(1)}$, so we are done. □

Lemma 4.21. *Let odd elements $a_i, b_i \in A$ be given for $1 \leq i \leq m$ such that $[a_i, b_j] = 0$ for all $1 \leq i, j \leq m$ and define*

$$D = \sum_{i=1}^m a_i b_i.$$

Suppose $[b_i, c] = 0$ for $1 \leq i \leq m$. Then for $n \geq 0$

$$(4.19) \quad [D, c]^{(n)} = (-1)^{n|c|} \sum_{1 \leq q_1, \dots, q_n \leq m} [a_{q_n}, [a_{q_{n-1}}, \dots, [a_{q_1}, c] \dots]] b_{q_1} \dots b_{q_n}.$$

Proof. For convenience we introduce, for any tuple of indices $\mathbf{q} = (q_1, \dots, q_n)$ of integers $1 \leq q_i \leq m$, the notation

$$(4.20) \quad \tau_{\mathbf{q}} = [a_{q_n}, [a_{q_{n-1}}, \dots [a_{q_1}, c] \dots]]$$

$$(4.21) \quad b_{\mathbf{q}} = b_{q_1} \dots b_{q_n}.$$

For $n = 0$ the right hand side of (4.19) is c by convention, so the statement is trivial. We compute using the inductive hypothesis that

$$\begin{aligned} [D, c]^{(n+1)} &= [D, [D, c]^{(n)}] \\ &= [D, (-1)^{n|c|} \sum_{\mathbf{q}} \tau_{\mathbf{q}} b_{\mathbf{q}}] \\ &= (-1)^{n|c|} \sum_{\mathbf{q}} \sum_i [a_i b_i, \tau_{\mathbf{q}} b_{\mathbf{q}}] \\ &= (-1)^{n|c|} \sum_{\mathbf{q}} \sum_i (a_i b_i \tau_{\mathbf{q}} b_{\mathbf{q}} - \tau_{\mathbf{q}} b_{\mathbf{q}} a_i b_i) \\ &= (-1)^{n|c|} \sum_{\mathbf{q}} \sum_i ((-1)^{|c|} a_i \tau_{\mathbf{q}} b_{\mathbf{q}} b_i - (-1)^n \tau_{\mathbf{q}} a_i b_{\mathbf{q}} b_i) \\ &= (-1)^{(n+1)|c|} \sum_{\mathbf{q}, i} [a_i, \tau_{\mathbf{q}}] b_{\mathbf{q}} b_i \end{aligned}$$

which completes the proof. \square

Eventually we are going to substitute $a_i = \lambda_i, b_i = \theta_i^*$ so that $D = \delta$, and we will take further $c = \gamma$. In this notation, the aim of all of this is to compute the commutators $[c, D^n]$ and then $[c, \exp(-D)]$. Since most of the calculation can be done in A , so we stick to the general setting as long as possible, and maintain the notation a_i, b_i, c, D and (4.20), (4.21) of the previous lemma.

Lemma 4.22. *For $n \geq 1$*

$$[c, D^n] = \sum_{i=0}^{n-1} (-1)^{(i+1)|c|+1} \sum_{i \leq p \leq n-1} \binom{p}{i} \sum_{\mathbf{q}=(q_1, \dots, q_{i+1})} \tau_{\mathbf{q}} b_{\mathbf{q}} D^{n-i-1}.$$

Proof. By Lemma 4.19 we have

$$(4.22) \quad [c, D^n] = \sum_{p=0}^{n-1} D^p [c, D] D^{n-p-1} = - \sum_{p=0}^{n-1} D^p [c, D] D^{n-p-1}.$$

Then using Lemma 4.20 this becomes

$$(4.23) \quad - \sum_{p=0}^{n-1} \sum_{i+j=p} \binom{p}{i} [D, c]^{(i+1)} D^{n-p-1+j}$$

which by Lemma 4.21 equals

$$(4.24) \quad \sum_{p=0}^{n-1} \sum_{i+j=p} \binom{p}{i} (-1)^{(i+1)|c|+1} \sum_{\mathbf{q}=(q_1, \dots, q_{i+1})} \tau_{\mathbf{q}} b_{\mathbf{q}} D^{n-i-1}$$

which is what we needed to show. \square

Putting this all together:

Proposition 4.23. *Let A be a \mathbb{Z}_2 -graded \mathbb{Q} -algebra and*

- *Let $a_i, b_i \in A$ be odd elements with $[a_i, b_j] = 0$ for $1 \leq i, j \leq m$.*
- *Assume that $D = \sum_{i=1}^m a_i b_i$ has finite order, that is, $D^r = 0$ for some $r > 0$.*
- *Let $c \in A$ be an odd element, with $[b_i, c] = 0$ for $1 \leq i \leq m$.*

Then

$$(4.25) \quad [c, \exp(-D)] = - \sum_{n \geq 1} \frac{1}{n!} \sum_{q_1, \dots, q_n} [a_{q_n}, [a_{q_{n-1}}, \dots [a_{q_1}, c] \dots] b_{q_1} \dots b_{q_n} \exp(-D).$$

Proof. Using Lemma 4.22

$$\begin{aligned} [c, \exp(-D)] &= \sum_{n \geq 0} \frac{(-1)^n}{n!} [c, D^n] \\ &= \sum_{n \geq 0} \frac{(-1)^{n+i}}{n!} \sum_{i=0}^{n-1} \sum_{i \leq p \leq n-1} \binom{p}{i} \sum_{\mathbf{q}=(q_1, \dots, q_{i+1})} \tau_{\mathbf{q}} b_{\mathbf{q}} D^{n-i-1}. \end{aligned}$$

Rewriting this sum in terms of the variable $t = n - i - 1$ yields

$$- \sum_{t \geq 0} \sum_{i \geq 0} \frac{(-1)^t}{(t+i+1)!} \sum_{i \leq p \leq t+i} \binom{p}{i} \sum_{\mathbf{q}=(q_1, \dots, q_{i+1})} \tau_{\mathbf{q}} b_{\mathbf{q}} D^t.$$

Using Lemma 4.24 below this may be rewritten as

$$- \sum_{t \geq 0} \sum_{i \geq 0} \frac{(-1)^t}{t!} \frac{1}{(i+1)!} \sum_{\mathbf{q}=(q_1, \dots, q_{i+1})} \tau_{\mathbf{q}} b_{\mathbf{q}} D^t$$

which clearly agrees with (4.25). \square

Lemma 4.24. *For integers $a, b \geq 0$*

$$(4.26) \quad \frac{b!}{(a+b+1)!} \sum_{a \leq n \leq a+b} \binom{n}{a} = \frac{1}{(a+1)!}$$

Proof. Denoting the left hand side of (4.26) by $\Omega_{a,b}$ the proof is an induction on b , i.e. by computing $\Omega_{a,b+1}$ in terms of $\Omega_{a,b}$. \square

Now let us return to the original problem, of calculating the transfer $\mathcal{T}(\gamma)$ from Definition 4.14. We will only need this in the following special case:

Corollary 4.25. *Let γ be an odd R -linear operator on X . Then*

$$(4.27) \quad \mathcal{T}(\gamma) = \gamma + \sum_{n \geq 1} \sum_{q_1, \dots, q_n} \frac{1}{n!} [\lambda_{q_n}, [\lambda_{q_{n-1}}, \dots [\lambda_{q_1}, \gamma] \dots]] \theta_{q_1}^* \dots \theta_{q_n}^*$$

is the transferred operator on $K \otimes X$, where the q_j 's range over $1, \dots, m$.

Proof. This is immediate from Proposition 4.23, substituting $a_i = \lambda_i, b_i = \theta_i^*, c = \gamma$, since

$$\exp(-D)c \exp(D) = c - [c, \exp(-D)] \exp(D).$$

\square

Remark 4.26. Observe that since the θ_i^* square to zero, only sequences of distinct indices q_1, \dots, q_n contribute in the sum (4.27). The low order terms are

$$(4.28) \quad \mathcal{T}(\gamma) = \gamma + \sum_q [\lambda_q, \gamma] \theta_q^* + \frac{1}{2} \sum_{p,q} [\lambda_p, [\lambda_q, \gamma]] \theta_q^* \theta_p^* + \dots$$

Finally, we compute the transfer of the operator θ_i on $S_m \otimes_k X$:

Theorem 4.27. *We have*

$$(4.29) \quad \mathcal{T}(\theta_i) = \theta_i - \sum_{n \geq 0} \sum_{q_1, \dots, q_n} \frac{1}{(n+1)!} [\lambda_{q_n}, [\lambda_{q_{n-1}}, [\dots [\lambda_{q_1}, \lambda_i] \dots]] \theta_{q_1}^* \dots \theta_{q_n}^*.$$

Proof. Observe that Corollary 4.25 and the preceding developments do not apply directly to θ_i , because $[\theta_j^*, \theta_i] = \delta_{ij}$ whereas we have required the commutator $[b_j, c] = 0$ to vanish in the above. However, we do have

$$[\delta, \theta_i] = \sum_j [\lambda_j \theta_j^*, \theta_i] = \sum_j \lambda_j \delta_{ij} = \lambda_i.$$

And we can adjust the calculations above as follows. Firstly, using Lemma 4.20 we obtain

$$\delta^p [\delta, \theta_i] = \sum_{i+j=p} \binom{p}{i} [\delta, \theta_i]^{(i+1)} \delta^j = \sum_{i+j=p} \binom{p}{i} [\delta, \lambda_i]^{(i)} \delta^j$$

Tracking this through the rest of the calculation, as above, yields (4.29). \square

Remark 4.28. The low order terms of (4.29) are

$$(4.30) \quad \mathcal{T}(\theta_i) = \theta_i - \lambda_i - \frac{1}{2} \sum_q [\lambda_q, \lambda_i] \theta_q^* + \cdots$$

It is immediate that

Theorem 4.29. *The closed odd operators $\mathcal{T}(\theta_i)$ and $\mathcal{T}(\theta_i^*) = \theta_i^*$ on the linear factorisation $(K \otimes X, d_K + d)$ form a representation of the Clifford algebra C_m .*

By construction, Proposition 4.11 is now promoted to an isomorphism of representations of C_m , with $\mathcal{T}(\theta_i), \mathcal{T}(\theta_i^*)$ acting on $K \otimes X$ and the θ_i, θ_i^* acting on $S_m \otimes_k X$.

Remark 4.30. The construction of the Clifford operators ψ_i, ψ_i^\dagger depends on the choice of homotopies $\lambda_1, \dots, \lambda_m$, but it is clear from what we have just said that different choices give rise to isomorphic representations of C_m . An explicit isomorphism can be written down using the techniques developed in this Section.

Remark 4.31. Suppose for each pair $1 \leq i, j \leq m$ that up to homotopy

$$(4.31) \quad [\lambda_i, \lambda_j] \simeq f_{ij} \cdot 1_X$$

for elements $f_{ij} \in R$. Then the formula for $\mathcal{T}(\theta_i)$ simplifies substantially: for any indices i, j, k we have $[\lambda_i, [\lambda_j, \lambda_k]] \simeq [\lambda_i, f_{jk}] = 0$ so that (4.29) becomes

$$(4.32) \quad \mathcal{T}(\theta_i) \simeq \theta_i - \lambda_i - \frac{1}{2} \sum_q f_{qi} \theta_q^*.$$

4.3 The finite model

We proceed to rediscover $Y \mid X$ with its Clifford action as a finite model of $Y \otimes X$. Using this we prove in the next section that \mathcal{C} and \mathcal{LG} are equivalent bicategories. The notation is as in Setup 4.1. We introduce formal variables θ_i of odd degree and set

$$S_m = \bigwedge (k\theta_1 \oplus \cdots \oplus k\theta_m).$$

The Koszul complex K of the sequence t_1, \dots, t_m (where $t_i = \partial_{y_i} V$) is defined by

$$(4.33) \quad K = S_m \otimes_k k[y], \quad d_K = \sum_{i=1}^m t_i \theta_i^*.$$

According to Strategy 4.2 the first step is to find a finite model of

$$(K \otimes_{k[y]} (Y \otimes X), d_K) = (S_m \otimes_k (Y \otimes X), d_K).$$

First step. A finite model for (K, d_K) is obtained using connections. The connection ∇ of (3.5) extends canonically to an operator on $k[[y]] \otimes_{k[t]} \Omega_{k[t]/k}^*$ where $\Omega^* = \bigwedge \Omega^1$ is the exterior algebra. By identifying dt_i with θ_i we may make the identification

$$\check{K} := K \otimes_{k[y]} k[[y]] \cong \Omega_{k[t]/k}^* \otimes_{k[t]} k[[y]]$$

so that ∇ becomes identified with a degree -1 k -linear operator on \check{K} . It is straightforward to check (see [15, Lemma 8.7]) that $[d_K, \nabla]$ is invertible on the graded submodule $\check{K}_{\leq -1}$ of negative degree terms (in the usual \mathbb{Z} -grading on the Koszul complex) and we define

$$H = [d_K, \nabla]^{-1} \nabla.$$

Proposition 4.32. *H is a k -linear splitting homotopy on the complex (\check{K}, d_K) .*

Proof. See [15, Section 8.1]. □

The associated strong deformation retract of \mathbb{Z} -graded k -complexes is

$$(4.34) \quad (J_V, 0) \xrightleftharpoons[\sigma]{\pi} (\check{K}, d_K), \quad H$$

where J_V is the Jacobi algebra from (3.2), $\pi : \check{K} \rightarrow J_V$ is the canonical quasi-isomorphism which vanishes on $\check{K}_{\leq -1}$ and is in degree zero the quotient map $k[[y]] \rightarrow J_V$, and σ is a k -linear embedding of J_V into $k[[y]]$ uniquely determined by ∇ . The splitting homotopy H , or equivalently the deformation retract (4.34), is the desired finite model of the Koszul complex \check{K} .

Recall that the tensor product $Y \check{\otimes} X$ of (3.4) is a finite rank matrix factorisation of $U - W$ over $R = k[x, z][[y]]$. The connection ∇ extends to a standard $k[x, z]$ -linear flat connection on R . A choice of homogeneous basis for X and Y allows us to write $Y \otimes X \cong k[x, y, z] \otimes_k Q$ for some \mathbb{Z}_2 -graded free k -module Q , so

$$(4.35) \quad \check{K} \otimes_{k[[y]]} (Y \check{\otimes} X) \cong \check{K} \otimes_{k[[y]]} (R \otimes_k Q) \cong \check{K} \otimes_k k[x, z] \otimes_k Q$$

In this way the k -linear splitting homotopy H on \check{K} induces an $k[x, z]$ -linear operator $H \otimes 1$ on $\check{K} \otimes_{k[[y]]} (Y \check{\otimes} X)$, which we again denote by H . It is immediate that

Lemma 4.33. *H is a $k[x, z]$ -linear splitting homotopy on $(\check{K} \otimes_{k[[y]]} (Y \check{\otimes} X), d_K)$.*

The associated strong deformation retract is

$$(Y \mid X, 0) \xrightleftharpoons[\sigma]{\pi} (\check{K} \otimes_{k[[y]]} Y \check{\otimes} X, d_K), \quad H$$

The map π is R -linear induced by the map $\pi : \check{K} \rightarrow J_V$, and $\sigma = 1 \otimes \sigma$ is defined using map σ of (4.34) and the isomorphism (4.35). This is the desired finite model of $K \otimes Y \otimes X$ equipped with only the differential d_K .

Second step. We now view $d_{Y \otimes X}$ as a perturbation. The transference problem asks how to define a splitting homotopy ϕ_∞ on the matrix factorisation $(\check{K} \otimes_{k[[y]]} Y \check{\otimes} X, d_K + d_{Y \otimes X})$ in such a way that the underlying graded module is $Y | X$. Since $Hd_{Y \otimes X}$ has finite order, it follows from the perturbation lemma (Theorem 4.8) that

$$\phi_\infty = \sum_{m \geq 0} (-1)^m (Hd_{Y \otimes X})^m H$$

is $k[x, z]$ -linear splitting homotopy achieving the desired aim.

Specifically, by [15, Proposition 7.1] ϕ_∞ is an $k[x, z]$ -linear splitting homotopy on $\check{K} \otimes_{k[[y]]} Y \check{\otimes} X$ and the associated $k[x, z]$ -linear strong deformation retract of linear factorisations of $U - W$ is of the form

$$(4.36) \quad (Y | X, d_{Y \otimes X}) \xrightleftharpoons[\sigma_\infty]{\pi} (\check{K} \otimes_{k[[y]]} Y \check{\otimes} X, d_K + d_{Y \otimes X}), \quad \phi_\infty.$$

Third step. Since each t_i acts null-homotopically on $Y \otimes X$ there is by Section 4.2 an isomorphism of linear factorisations

$$(4.37) \quad (\check{K} \otimes_{k[[y]]} Y \otimes X, d_K + d_{Y \otimes X}) \xrightleftharpoons[\exp(\delta)]{\exp(-\delta)} (S_m \otimes_k Y \check{\otimes} X, d_{Y \otimes X})$$

where $\delta = \sum_i \lambda_i \theta_i^*$ for a choice λ_i of null-homotopy for the action of t_i . From the identity $d_X^2 = V - W$ we deduce that $\partial_{y_i}(d_X)d_X + d_X\partial_{y_i}(d_X) = \partial_{y_i}V = t_i$ so that we may, and do, choose $\lambda_i = \partial_{y_i}(d_X)$, see Section 2.3.

Fourth step. Combining (4.36) and (4.37) we have a homotopy equivalence of matrix factorisations over $k[x, z]$

$$(4.38) \quad Y | X \xrightleftharpoons[\sigma_\infty]{\pi} \check{K} \otimes_{k[[y]]} Y \check{\otimes} X \xrightleftharpoons[\exp(\delta)]{\exp(-\delta)} S_m \otimes_k (Y \check{\otimes} X).$$

By [15, Remark 7.7] the canonical map $\epsilon : Y \otimes X \longrightarrow Y \check{\otimes} X$ is a homotopy equivalence over $k[x, z]$ so finally we have

Theorem 4.34. *There is a homotopy equivalence of matrix factorisations over $k[x, z]$*

$$(4.39) \quad Y | X \xrightleftharpoons[\Phi^{-1}]{\Phi} S_m \otimes_k (Y \otimes X)$$

where $\Phi = \pi \exp(-\delta)\epsilon$ and $\Phi^{-1} = \epsilon^{-1} \exp(\delta)\sigma_\infty$.

The canonical action of C_m on the spinor representation S_m by wedge product and contraction $\{\theta_i, \theta_i^*\}_{i=1}^m$ (see Lemma 2.17) induces an action of C_m on $S_m \otimes_k (Y \otimes X)$. In Section 2.2 we defined an action of C_m on the cut $Y | X$ by operators $\{\gamma_i, \gamma_i^\dagger\}_{i=1}^m$, and it remains to show that these two actions are compatible.

Proposition 4.35. *The map Φ is an isomorphism of C_m -representations in the homotopy category. That is, there are homotopies for $1 \leq i \leq m$*

$$\gamma_i \Phi \simeq \Phi \theta_i, \quad \gamma_i^\dagger \Phi \simeq \Phi \theta_i^*.$$

Proof. We prove that $\gamma_i \simeq \Phi \theta_i \Phi^{-1}$ and $\gamma_i^\dagger \simeq \Phi \theta_i^* \Phi^{-1}$. With the diagram (4.38) in mind, and following the terminology of Section 4.2, we define the transfers of θ_i, θ_i^* from closed operators on $S_m \otimes_k (Y \check{\otimes} X)$ to operators on the intermediate object $\check{K} \otimes_{k[[y]]} Y \check{\otimes} X$ by

$$\mathcal{T}(\theta_i) = \exp(-\delta) \theta_i \exp(\delta), \quad \mathcal{T}(\theta_i^*) = \exp(-\delta) \theta_i^* \exp(\delta).$$

Then we use Lemma 2.14, according to which there are homotopies

$$[\lambda_j, \lambda_k] = [\partial_{y_j}(d_X), \partial_{y_k}(d_X)] \simeq \partial_{y_j y_k}(V).$$

Therefore by Theorem 4.27 and Remark 4.31 we have $\mathcal{T}(\theta_i^*) = \theta_i^*$ and a homotopy

$$(4.40) \quad \mathcal{T}(\theta_i) \simeq \theta_i - \lambda_i - \frac{1}{2} \sum_q \partial_{y_q y_i}(V) \theta_q^*.$$

It remains to compute the further transfer of these operators to $Y|X$. The key point is proven in Lemma 4.36 below, where we show that there is a homotopy

$$(4.41) \quad \pi \theta_{q_1}^* \cdots \theta_{q_l}^* \sigma_\infty \simeq \text{At}_{q_1} \cdots \text{At}_{q_l}$$

for any indices q_1, \dots, q_l . Hence

$$\Phi \theta_i^* \Phi^{-1} \simeq \pi \mathcal{T}(\theta_i^*) \sigma_\infty = \pi \theta_i^* \sigma_\infty \simeq \text{At}_i = \gamma_i^\dagger.$$

Since π projects onto θ -degree zero,

$$\begin{aligned} \Phi \theta_i \Phi^{-1} &\simeq \pi \mathcal{T}(\theta_i) \sigma_\infty \\ &\simeq -\pi \lambda_i \sigma_\infty - \frac{1}{2} \sum_q \partial_{y_q y_i}(V) \pi \theta_q^* \sigma_\infty \\ &\simeq -\lambda_i - \frac{1}{2} \sum_q \partial_{y_q y_i}(V) \text{At}_q \\ &= \gamma_i \end{aligned}$$

which completes the proof. □

Lemma 4.36. *In the context of the diagram (4.38) there is a $k[x, z]$ -linear homotopy*

$$\pi \theta_{q_1}^* \cdots \theta_{q_l}^* \sigma_\infty \simeq \text{At}_{q_1} \cdots \text{At}_{q_l}$$

for any sequence of indices $1 \leq q_1, \dots, q_l \leq m$.

Proof. This is implicit in [15, Section 10]. The formula $\sigma_\infty = \sum_{m \geq 0} (-1)^m (Hd_{Y \otimes X})^m \sigma$ is given there, and with $t_i = \partial_{z_i} V$ the calculation that takes place in [15, (10.3)] yields

$$(4.42) \quad \sigma_\infty = \sum_{s \geq 0} \sum_{p_1, \dots, p_s} (-1)^{\binom{s+1}{2}} \frac{1}{s!} [d_{Y \otimes X}, \partial_{t_{p_1}}] \cdots [d_{Y \otimes X}, \partial_{t_{p_s}}] \theta_{p_1} \cdots \theta_{p_s} + (t \text{ terms}).$$

That is, σ_∞ is given modulo the submodule $(t_1, \dots, t_m)K \otimes Y \otimes X$ by the formula above. Since π vanishes on this submodule,

$$\begin{aligned} \pi \theta_{q_1}^* \cdots \theta_{q_l}^* \sigma_\infty &= \sum_{s \geq 0} \sum_{p_1, \dots, p_s} (-1)^{\binom{s+1}{2}} \frac{1}{s!} \pi \theta_{q_1}^* \cdots \theta_{q_l}^* \text{At}_{p_1} \cdots \text{At}_{p_s} \theta_{p_1} \cdots \theta_{p_s} \\ &= \sum_{s \geq 0} \sum_{p_1, \dots, p_s} (-1)^{\binom{s+1}{2} + sl} \frac{1}{s!} \text{At}_{p_1} \cdots \text{At}_{p_s} \pi \theta_{q_1}^* \cdots \theta_{q_l}^* \theta_{p_1} \cdots \theta_{p_s}. \end{aligned}$$

Now π is non-vanishing only on terms of θ -degree zero, so the sum restricts to $s = l$ and to \mathbf{p} a permutation of \mathbf{q} . Since

$$\theta_{q_1}^* \cdots \theta_{q_l}^* \theta_{q_{\sigma(1)}} \cdots \theta_{q_{\sigma(l)}} = (-1)^{|\sigma|} (-1)^{\binom{l}{2}}$$

we have, using the fact that the Atiyah classes anti-commute,

$$\begin{aligned} \pi \theta_{q_1}^* \cdots \theta_{q_l}^* \sigma_\infty &= \sum_{\sigma \in \mathfrak{S}_l} (-1)^{|\sigma|} \frac{1}{l!} \text{At}_{q_{\sigma(1)}} \cdots \text{At}_{q_{\sigma(l)}} \\ &\simeq \text{At}_{q_1} \cdots \text{At}_{q_l} \end{aligned}$$

which completes the proof. \square

Finally, we will need naturality of Theorem 4.34.

Proposition 4.37. *If $\phi : X \rightarrow X'$ and $\psi : Y \rightarrow Y'$ are morphisms of matrix factorisations the diagram*

$$(4.43) \quad \begin{array}{ccc} Y \mid X & \xleftarrow{\Phi} & S_m \otimes_k (Y \otimes X) \\ \psi \mid \phi \downarrow & & \downarrow \psi \otimes \phi \\ Y' \mid X' & \xleftarrow{\Phi} & S_m \otimes_k (Y' \otimes X') \end{array}$$

commutes up to $k[x, z]$ -linear homotopy.

Proof. Given the formula $\Phi = \pi \exp(-\delta) \epsilon$ it is clearly enough to show that $\psi \mid \phi$ commutes up to homotopy with δ , and for this it is enough to show that there is a homotopy

$$\partial_{y_i}(d_{X'}) \phi \simeq \phi \partial_{y_i}(d_X)$$

for each i . But this is the content of Lemma 2.13. \square

This achieves our aim: we took direct sums of copies of $Y \otimes X$ to form $S_m \otimes_k (Y \otimes X)$ and then used perturbation to find a finite model of this latter object. It is worth noting that, in the language of splitting homotopies, the cut $Y | X$ is the solution of a fixed point problem formulated in terms of linear operators on the initial data $S_m \otimes_k (Y \otimes X)$.

Remark 4.38. The isomorphism of C_m -representations in the homotopy category

$$\Phi : S_m \otimes_k (Y \otimes X) \longrightarrow Y | X$$

is straightforward to compute. Firstly, since $\Phi = \pi \exp(-\delta)\varepsilon$ and the operator $\delta = \sum_i \lambda_i \theta_i^*$ annihilates the identity of the exterior algebra $1 \in S_m$, the composite

$$(4.44) \quad (k \cdot 1) \otimes_k (Y \otimes X) \xrightarrow{\text{inc}} S_m \otimes_k (Y \otimes X) \xrightarrow{\Phi} Y | X$$

is just the quotient map $Y \otimes X \longrightarrow Y | X$, which we write as $\nu \mapsto \bar{\nu}$. Using

$$\exp(-\delta) = 1 - \sum_i \lambda_i \theta_i^* - \frac{1}{2} \sum_{i,j} \lambda_i \lambda_j \theta_i^* \theta_j^* + \dots$$

we have for example

$$\begin{aligned} \Phi(\theta_1 \theta_2 \otimes \nu) &= -\frac{1}{2} \sum_{i,j} \pi \lambda_i \lambda_j \theta_i^* \theta_j^* (\theta_1 \theta_2 \otimes \nu) \\ &= \frac{1}{2} (\lambda_1 \lambda_2 - \lambda_2 \lambda_1) (\bar{\nu}). \end{aligned}$$

4.4 The equivalence of \mathcal{C} and \mathcal{LG}

The bicategories \mathcal{LG} and \mathcal{C} have the same objects. Their morphism categories are

$$\begin{aligned} \mathcal{C}(W, V) &= \left(\text{hmf}(k[x, y], V - W)^\omega \right)^\bullet, \\ \mathcal{LG}(W, V) &= \text{hmf}(k[x, y], V - W)^\oplus \end{aligned}$$

with the notation from Section 2.2. These are equivalent supercategories, via the following chain of equivalences

$$(4.45) \quad \begin{array}{ccc} & \text{hmf}(k[x, y], V - W)^\omega & \\ \swarrow \iota & & \searrow F \\ \left(\text{hmf}(k[x, y], V - W)^\omega \right)^\bullet & \xleftarrow{\mathcal{Z}_{W,V}} & \text{hmf}(k[x, y], V - W)^\oplus \end{array}$$

where ι is the equivalence of a supercategory with its Clifford thickening, F is the functor given earlier in (2.3) and we define $\mathcal{Z}_{W,V} = \iota \circ F^{-1}$ to make the diagram commute.

The functor F is defined by choosing for each object (X, e) a (possibly infinite rank) matrix factorisation $F(X, e)$ which splits the idempotent e . That is, such that there are morphisms in the homotopy category

$$X \begin{array}{c} \xrightarrow{f} \\ \xleftarrow{g} \end{array} F(X, e)$$

with $fg = 1$ and $gf = e$. The inverse is defined by choosing, for each infinite rank matrix factorisation T which is a summand in the homotopy category of something finite rank, an actual finite rank X and morphisms f, g with $fg = 1_T$, and defining

$$F^{-1}(T) = (X, gf).$$

We may arrange these choices so that if T is *already* finite rank then $F^{-1}(T) = (T, 1_T)$.

Definition 4.39. A *strong superfunctor* $F : \mathcal{C} \rightarrow \mathcal{D}$ between supercategories without units is a strong superfunctor in the sense of Definition 2.9, but without the data related to units. We say further that F is an *equivalence* if

- for all $a, b \in \mathcal{C}$, $F_{a,b} : \mathcal{C}(a, b) \rightarrow \mathcal{D}(Fa, Fb)$ is an equivalence of categories, and
- for all $d \in \mathcal{D}$ there exists $a \in \mathcal{C}$ with $Fa \cong c$ in the sense that there are 1-morphisms $f : Fa \rightarrow c$ and $g : c \rightarrow Fa$ with $fg \cong 1$ and $gf \cong 1$.

Theorem 4.40. *There is an equivalence $\mathcal{Z} : \mathcal{LG} \rightarrow \mathcal{C}$ of supercategories without units.*

Proof. We define \mathcal{Z} to be the identity on objects and $\mathcal{Z}_{W,V}$ on 1- and 2-morphisms. The only remaining content is to produce a 2-isomorphism

$$(4.46) \quad \varphi_{Y,X} : \mathcal{Z}_{V,U}(Y) \mid \mathcal{Z}_{W,V}(X) \rightarrow \mathcal{Z}_{W,U}(Y \otimes X)$$

in the usual situation of Setup 4.1, which is natural in both Y and X . Once we have done this the rest follows by functoriality.

Since Y, X in this case are already finite rank $\mathcal{Z}(Y) = Y$ and $\mathcal{Z}(X) = X$. The calculation of $\mathcal{Z}(Y \otimes X)$ is more interesting, because $Y \otimes X$ is infinite rank over $k[x, z]$. However we know from Proposition 4.35 that there is an isomorphism in the homotopy category of (infinite rank) matrix factorisations of $U - W$ over $k[x, z]$

$$(4.47) \quad Y \mid X \begin{array}{c} \xleftarrow{\Phi} \\ \xrightarrow{\Phi^{-1}} \end{array} S_m \otimes_k (Y \otimes X)$$

where $Y \mid X$ is finite rank. Recall from Definition 2.18 that the idempotent

$$e_m = \gamma_1^\dagger \cdots \gamma_m^\dagger \gamma_m \cdots \gamma_1 \in C_m$$

represents projection onto $k \cdot 1$ in S_m . Since (4.47) is an isomorphism of C_m -representations the idempotent e_m acting on $Y \mid X$ must split to the same thing as the idempotent e_m

acting on the right hand side, which obviously splits to $Y \otimes X$. Put differently, there is a diagram in the homotopy category

$$Y \mid X \begin{array}{c} \xrightarrow{f} \\ \xleftarrow{g} \end{array} Y \otimes X$$

where g is the quotient map, and $fg = 1_{Y \otimes X}$ while $gf = e_m$.¹ Then we may take as our definition

$$\mathcal{Z}(Y \otimes X) = (Y \mid X, e_m).$$

So the problem is to define a 2-isomorphism (4.46), that is,

$$\varphi_{Y,X} : (Y \mid X, 1, \{\gamma_i, \gamma_i^\dagger\}_{i=1}^m) \longrightarrow (Y \mid X, e_m)$$

in the Clifford thickening of the idempotent completion of $\mathrm{hmf}(k[x, z], U - W)$, where the left hand side is equipped with the idempotent 1 and the action $\{\gamma_i, \gamma_i^\dagger\}_{i=1}^m$ of C_m while the right hand side has no Clifford action and idempotent e_m .

But by Lemma 2.27 the identity map $\varphi_{Y,X} = 1_{Y \mid X}$ is such an isomorphism, because the object $(Y \mid X, e_m)$ of the idempotent completion by definition splits the idempotent e_m on $Y \mid X$. Naturality of (4.46) in Y, X is a consequence of the naturality of (4.47), see Proposition 4.37. \square

Remark 4.41. Obviously the 1-morphisms $\Delta_W : W \longrightarrow W$ which are the units in \mathcal{LG} are, by virtue of the equivalence \mathcal{Z} , also units in \mathcal{C} , so that this is an honest bicategory and \mathcal{Z} an equivalence of bicategories. The reason we do not begin with units in \mathcal{C} is that the unitors are no simpler in \mathcal{C} than \mathcal{LG} , so there is no real point to introducing them.

4.5 Example: the Hom complex

Using the bicategory \mathcal{C} we can study the complex $\mathrm{Hom}_R(Y, X)$ for two finite rank matrix factorisations X, Y of a potential $V \in R = k[y_1, \dots, y_m]$. In particular, we will explain how the homotopy relation on morphisms of matrix factorisations is effectively computable.

The pair $(k, 0)$ is an object in \mathcal{LG} and we may view X and the dual matrix factorisation Y^\vee of $-V$ as a pair of 1-morphisms

$$(4.48) \quad 0 \xrightarrow{X} V \xrightarrow{Y^\vee} 0.$$

The composite is a 1-morphism $0 \longrightarrow 0$, that is, \mathbb{Z}_2 -graded complex of k -modules. This is the Hom complex of R -linear homogeneous maps $Y \longrightarrow X$,

$$Y^\vee \otimes_R X \cong \mathrm{Hom}_R(Y, X)$$

with the differential $\phi \mapsto d_X \circ \phi - (-1)^{|\phi|} \phi \circ d_Y$. The cut is

$$Y^\vee \mid X = Y^\vee \otimes_R J_V \otimes_R X \cong \mathrm{Hom}_{J_V}(\bar{Y}, \bar{X})$$

¹ f is Φ^{-1} followed by projection onto θ -degree zero. Not a very friendly map.

where $\bar{X} = X \otimes_R J_V$, $\bar{Y} = Y \otimes_R J_V$. There is by Theorem 4.34 a C_m -linear isomorphism in the homotopy category of \mathbb{Z}_2 -graded complexes of free k -modules

$$(4.49) \quad \Phi : S_m \otimes_k \text{Hom}_R(Y, X) \longrightarrow Y^\vee | X.$$

Example 4.42. Suppose $V = y^N$ in $k[y]$ for $N \geq 2$. Take matrix factorisations X, Y with the same underlying free $R = k[y]$ -module $X = Y = R \oplus R\theta$ and differentials

$$d_Y = y^i \theta + y^{N-i} \theta^* = \begin{pmatrix} 0 & y^{N-i} \\ y^i & 0 \end{pmatrix}, \quad d_X = y^j \theta + y^{N-j} \theta^* = \begin{pmatrix} 0 & y^{N-j} \\ y^j & 0 \end{pmatrix}$$

where $1 \leq i, j \leq N/2$. Using the basis $\text{Hom}_R(Y, X) = R\theta\theta^* \oplus R\theta^*\theta \oplus R\theta \oplus R\theta^*$,

$$(4.50) \quad d_{\text{Hom}} = \begin{pmatrix} 0 & 0 & y^{N-i} & y^j \\ 0 & 0 & y^{N-j} & y^i \\ -y^i & y^j & 0 & 0 \\ y^{N-j} & -y^{N-i} & 0 & 0 \end{pmatrix}.$$

Let us now compute the cut $Y^\vee | X$, according to Definition 3.8. With

$$(4.51) \quad J_V = k[y]/y^{N-1} = k \cdot 1 \oplus k \cdot y \oplus \cdots \oplus k \cdot y^{N-2}$$

the matrix of multiplication by y on J_V is $[y] = \begin{pmatrix} 0 & 0 \\ I_{N-2} & 0 \end{pmatrix}$. The differential on

$$(4.52) \quad Y^\vee | X = J_V \theta \theta^* \oplus J_V \theta^* \theta \oplus J_V \theta \oplus J_V \theta^*$$

is then obtained from (4.50) by replacing every y by the matrix $[y]$.

The Clifford algebra C_1 acts on $Y^\vee | X$ by generators γ, γ^\dagger . To determine their matrices in the basis determined by (4.51) and (4.52), note that with $t = \partial_y(V) = Ny^{N-1}$, $k[y]$ is a free $k[t]$ -module with basis $1, \dots, y^{N-2}$. Given a polynomial $f(y)$ we may uniquely write it as $f(y) = \sum_{l=0}^{N-2} f_l \cdot y^l$ with $f_l \in k[t]$. There is a connection $\nabla : k[y] \longrightarrow k[y] \otimes_{k[t]} \Omega_{k[t]/k}^1$ whose associated k -linear operator ∂_t is

$$\partial_t : k[y] \longrightarrow k[y], \quad \partial_t(f) = \sum_{l=0}^{N-2} \frac{\partial}{\partial t}(f_l) y^l$$

where $\frac{\partial}{\partial t}$ is the usual operator on $k[t]$. For $q < N-1$ we have $\partial_t(y^q) = 0$ and $\partial_t(y^{N-1+q}) = \frac{1}{N} y^q$, so the k -linear operator

$$R_a : J_V \longrightarrow J_V, \quad y^q \mapsto \partial_t(y^{a+q})$$

has the following block form for $0 \leq a \leq N-2$

$$[R_a] = \frac{1}{N} \begin{pmatrix} 0 & I_a \\ 0 & 0 \end{pmatrix}.$$

The Atiyah class is the operator $\text{At} = [d_{\text{Hom}}, \partial_t] = d_{\text{Hom}}\partial_t - \partial_t d_{\text{Hom}}$. For example,

$$\text{At}(y^q \theta \theta^*) = -\partial_t d_{\text{Hom}}(y^q \theta \theta^*) = \partial_t(y^{i+q})\theta - \partial_t(y^{N-j+q})\theta^*.$$

The matrix of the Atiyah class as an operator γ^\dagger on $Y^\vee | X$ is therefore

$$(4.53) \quad \gamma^\dagger = \begin{pmatrix} 0 & 0 & -[R_{N-i}] & -[R_j] \\ 0 & 0 & -[R_{N-j}] & -[R_i] \\ [R_i] & -[R_j] & 0 & 0 \\ -[R_{N-j}] & [R_{N-i}] & 0 & 0 \end{pmatrix}.$$

Whereas the matrix of $\partial_y(d_X) = jy^{j-1}\theta + (N-j)y^{N-j-1}\theta^*$ on $Y^\vee | X$ is

$$\partial_y(d_X) = \begin{pmatrix} 0 & 0 & 0 & j[y^{j-1}] \\ 0 & 0 & (N-j)[y^{N-j-1}] & 0 \\ 0 & j[y^{j-1}] & 0 & 0 \\ (N-j)[y^{N-j-1}] & 0 & 0 & 0 \end{pmatrix}.$$

By definition

$$\gamma = -\partial_y(d_X) - \frac{1}{2}N(N-1)y^{N-2}\gamma^\dagger$$

which completes the description of the cut $(Y^\vee | X, \{\gamma, \gamma^\dagger\})$.

A Tensor products in supercategories

Let \mathcal{C} be an idempotent complete supercategory.

Definition A.1. Let V be a \mathbb{Z}_2 -graded k -module and X an object of \mathcal{C} . An object of \mathcal{C} representing the functor $\text{Hom}_k^0(V, \mathcal{C}^*(X, -))$ is denoted $V \otimes_k X$ if it exists. That is to say, the tensor product consists of an object $V \otimes_k X$ and a natural isomorphism

$$\rho_Y : \mathcal{C}(V \otimes_k X, Y) \longrightarrow \text{Hom}_k^0(V, \mathcal{C}^*(X, Y)).$$

Such a pair is unique up to unique isomorphism, if it exists.

Remark A.2. From ρ we also obtain an isomorphism of \mathbb{Z}_2 -graded k -modules

$$\mathcal{C}^*(V \otimes_k X, Y) \cong \text{Hom}_k^*(V, \mathcal{C}^*(X, Y)).$$

The tensor product $V \otimes_k X$ is made functorial in V and X in such a way that ρ is natural in all its variables.

Example A.3. If $V = k^{\oplus n} \oplus k[1]^{\oplus m}$ is finite and free then

$$V \otimes_k X = X^{\oplus n} \oplus \Psi X^{\oplus m}$$

is a representing object, with the obvious isomorphism ρ .

Lemma A.4. *If \mathcal{C} is idempotent complete and V is a finitely generated projective \mathbb{Z}_2 -graded k -module, then $V \otimes_k X$ exists for any object X .*

Proof. Let V' be a finite free k -module with $f : V' \rightarrow V, g : V \rightarrow V'$ satisfying $fg = 1$. This determines an idempotent $1 \otimes e$ on $V' \otimes_k X$, which splits by hypothesis in \mathcal{C} . The object splitting this idempotent has the right property to be $V \otimes_k X$. \square

More generally, let A be a \mathbb{Z}_2 -graded k -algebra.

Definition A.5. Given a right \mathbb{Z}_2 -graded A -module V and an object X of \mathcal{C} , we denote by $V \otimes_A X$ the object representing the functor $\text{Hom}_A^0(V, \mathcal{C}^*(X, -))$ if it exists. That is, the tensor product is an object $V \otimes_A X$ together with a natural isomorphism

$$(A.1) \quad \rho_Y : \mathcal{C}(V \otimes_A X, Y) \rightarrow \text{Hom}_A^0(V, \mathcal{C}^*(X, Y)).$$

As before, the tensor product is functorial in both V and X .

Henceforth we concentrate our attention on \mathbb{Z}_2 -graded k -algebras A which are *Morita trivial* in the sense that they are isomorphic to an algebra of the form $\text{End}_k(P)$ for a finite rank free \mathbb{Z}_2 -graded k -module P .

Lemma A.6. *If P is a finite rank free \mathbb{Z}_2 -graded k -module and $A \cong \text{End}_k(P)$, then for any A -module X in \mathcal{C} there is an object \tilde{X} and an isomorphism of A -modules $X \cong P \otimes_k \tilde{X}$.*

Proof. The natural idempotents in A act as idempotents on X , which split. \square

Lemma A.7. *If A is Morita trivial and V is a \mathbb{Z}_2 -graded right A -module which is finitely generated and projective as a k -module, then $V \otimes_A X$ exists for any A -module X .*

Proof. Write $A \cong \text{End}_k(P)$ and $P^* = \text{Hom}_k(P, k)$ so that $V \cong \tilde{V} \otimes_k P^*$ as right A -modules for some \mathbb{Z}_2 -graded k -module \tilde{V} . Since V is finitely generated projective over k , so is \tilde{V} . By the previous lemma, $X \cong P \otimes_k \tilde{X}$ for some object \tilde{X} , so we might expect that

$$V \otimes_A X \cong (\tilde{V} \otimes_k P^*) \otimes_A (P \otimes_k \tilde{X}) \cong \tilde{V} \otimes_k \tilde{X}$$

Working backwards: $\tilde{V} \otimes_k \tilde{X}$ exists and has the right universal property since

$$\begin{aligned} \mathcal{C}(\tilde{V} \otimes_k \tilde{X}, Y) &\cong \text{Hom}_k(V \otimes_A P, \mathcal{C}^*(\tilde{X}, Y)) \\ &\cong \text{Hom}_A(V, P^* \otimes_k \mathcal{C}^*(\tilde{X}, Y)) \\ &\cong \text{Hom}_A(V, \mathcal{C}^*(X, Y)) \end{aligned}$$

as claimed. \square

From now on A, B, C are Morita trivial k -algebras. Recall \mathcal{C}_A denotes the category of left A -modules in \mathcal{C} .

Definition A.8. Given a left A -module X in \mathcal{C} , the object ΨX is a left A -module, where $a \in A_0$ acts by $\Psi(a)$ and $a \in A_1$ acts by

$$\Psi(X) \xrightarrow{\Psi(a)} \Psi\Psi(X) \xrightarrow{-1} \Psi\Psi(X) .$$

This determines a functor $\Psi : \mathcal{C}_A \longrightarrow \mathcal{C}_A$.

Lemma A.9. (\mathcal{C}_A, Ψ) is a supercategory.

Proof. The natural isomorphism $\xi_X : \Psi\Psi(X) \longrightarrow X$ is given by the identity in \mathcal{C} . \square

Let V be a B - A -bimodule which is finitely generated and projective over k . If X is a left A -module in \mathcal{C} then $V \otimes_A X$ is made into a left B -module in \mathcal{C} uniquely such that (A.1) is a natural isomorphism of B -modules. Hence the tensor product gives a functor

$$\Phi_V = V \otimes_A (-) : \mathcal{C}_A \longrightarrow \mathcal{C}_B .$$

Given a B - A -bimodule V and an object X for which $V \otimes_A X$ exists, there are natural isomorphisms of B -modules

$$\Psi V \otimes_A X \cong \Psi(V \otimes_A X) \cong V \otimes_A \Psi X$$

induced from the isomorphisms

$$\mathrm{Hom}_A(\Psi V, \mathcal{C}^*(X, -)) \cong \mathrm{Hom}_A(V, \Psi \mathcal{C}^*(X, -)) \cong \mathrm{Hom}_A(V, \mathcal{C}^*(\Psi X, -)) .$$

It follows that Φ_V is a superfunctor.

Lemma A.10. Let W be a C - B -bimodule and V a B - A -bimodule in \mathcal{C} . Then for any left A -module X in \mathcal{C} there is a natural isomorphism of C -modules

$$(W \otimes_B V) \otimes_A X \cong W \otimes_B (V \otimes_A X) .$$

Proof. This follows from the isomorphism

$$\begin{aligned} \mathrm{Hom}_A(W, \mathcal{C}^*(V \otimes_A X, -)) &\cong \mathrm{Hom}_A(W, \mathrm{Hom}_A^*(V, \mathcal{C}^*(X, -))) \\ &\cong \mathrm{Hom}_A(W \otimes_A V, \mathcal{C}^*(X, -)) . \end{aligned}$$

\square

Now we specialise to the Clifford algebras C_n and their modules.

Lemma A.11. Let X be a left C_n -module in \mathcal{C} . There is a k -linear isomorphism

$$\mathrm{Hom}_{C_n}^0(S_n, X) \cong \{x \in X^0 \mid \gamma_i^\dagger \cdot x = 0 \text{ for all } 1 \leq i \leq n\}$$

defined by $f \mapsto f(1)$. If X is a right C_n -module there is a k -linear isomorphism

$$\mathrm{Hom}_{C_n}^0((S_n)^*, X) \cong \{x \in X^0 \mid x \cdot \gamma_i = 0 \text{ for all } 1 \leq i \leq n\}$$

defined by $f \mapsto f(1^*)$.

Lemma A.12. *For an C_n -module Y there is a k -linear isomorphism*

$$\mathcal{C}_{C_n}(S_n \otimes_k X, Y) \cong \{\delta \in \mathcal{C}(X, Y) \mid \gamma_i^\dagger \circ \delta = 0 \text{ for all } 1 \leq i \leq n\}$$

defined by sending $\alpha : S_n \otimes_k X \longrightarrow Y$ to the composite

$$X \cong k \otimes_k X \xrightarrow{\iota \otimes 1} S_n \otimes_k X \xrightarrow{\alpha} Y$$

where $\iota : k \longrightarrow S_n$ is defined by $\iota(1) = 1$.

Proof. This follows from Lemma A.11 and the isomorphism

$$\mathcal{C}_{C_n}(S_n \otimes_k X, Y) \cong \text{Hom}_{C_n}(S_n, \mathcal{C}^*(X, Y)).$$

□

More generally

Lemma A.13. *Let $X \in C_{C_p}$ and $Y \in \mathcal{C}_{C_q}$. There is a k -linear isomorphism*

$$\begin{aligned} \mathcal{C}_{C_q}(S_{q,p} \otimes_{C_p} X, Y) &\cong \{\delta : X \longrightarrow Y \mid \delta \circ \gamma_i = 0 \text{ for } 1 \leq i \leq p \text{ and} \\ &\quad \gamma_i^\dagger \circ \delta = 0 \text{ for } 1 \leq i \leq q\} \end{aligned}$$

defined by sending $\alpha : S_{q,p} \otimes_{C_p} X \longrightarrow Y$ to the composite

$$X \cong C_p \otimes_{C_p} X \xrightarrow{\iota \otimes 1} S_{q,p} \otimes_{C_p} X \xrightarrow{\alpha} Y$$

where $\iota : C_p \longrightarrow S_{q,p}$ is defined by $\iota(1) = 1 \otimes 1^$.*

Proof. Using the previous lemma and Lemma A.11,

$$\begin{aligned} \mathcal{C}_{C_q}(S_{q,p} \otimes_{C_p} X, Y) &\cong \text{Hom}_{C_p - C_q - \text{bimod}}^0(S_q \otimes_k S_p^*, \mathcal{C}^*(X, Y)) \\ &\cong \text{Hom}_{C_q}^0(S_q, \text{Hom}_{C_p}^*(S_p^*, \mathcal{C}^*(X, Y))) \\ &\cong \{\delta \in \text{Hom}_{C_q}^0(S_p^*, \mathcal{C}^*(X, Y)) \mid \gamma_i^\dagger \circ \delta = 0 \text{ for } 1 \leq i \leq q\} \\ &\cong \{\delta : X \longrightarrow Y \mid \delta \circ \gamma_i = 0 \text{ for } 1 \leq i \leq p \text{ and } \gamma_i^\dagger \circ \delta = 0 \text{ for } 1 \leq i \leq q\} \end{aligned}$$

as claimed. □

B Constructing superbicategories

This appendix collects the data needed for constructing a superbicategory \mathcal{B} from a collection of supercategories $\mathcal{B}(a, b)$. Given a bicategory \mathcal{B} and for each pair of objects a, b the structure of a supercategory on $\mathcal{B}(a, b)$, we denote composition by

$$T : \mathcal{B}(a, b) \otimes_k \mathcal{B}(b, c) \longrightarrow \mathcal{B}(a, c).$$

We assume this is a superfunctor, i.e. that (??)-(??) commute. Since our main example is constructing the supercategory associated to a cut system, we write $T(X, Y)$ as $Y \mid X$. We denote the associator by α , the units in \mathcal{B} by $\Delta_a : a \longrightarrow a$, and the unitors for $X : a \longrightarrow b$ by $\lambda_X : \Delta_b \mid X \longrightarrow X$ and $\rho_X : X \mid \Delta_a \longrightarrow X$.

Suppose that for all objects a, b, c we are given natural isomorphisms

$$T \circ (\Psi \otimes 1) \longrightarrow \Psi \circ T, \quad T \circ (1 \otimes \Psi) \longrightarrow \Psi \circ T$$

both of which will be denoted τ . We define for an object a the 1-morphism $\Psi_a = \Psi(\Delta_a)$ and the 2-isomorphism ξ_a to be the following composite (using the structure maps of \mathcal{B} and isomorphisms τ)

$$\xi_a : \Psi_a \mid \Psi_a = \Psi(\Delta_a) \mid \Psi(\Delta_a) \cong \Psi(\Delta_a \mid \Psi(\Delta_a)) \cong \Psi^2(\Delta_a \mid \Delta_a) \cong \Delta_a \mid \Delta_a \cong \Delta_a.$$

Similarly given a 1-morphism $X : a \longrightarrow b$ we have a 2-isomorphism γ_X

$$\gamma_X : X \mid \Psi_a = X \mid \Psi(\Delta_a) \cong \Psi(X \mid \Delta_a) \cong \Psi(X) \cong \Psi(\Delta_b \mid X) \cong \Psi(\Delta_b) \mid X = \Psi_b \mid X.$$

Given a 1-morphism $X : a \longrightarrow b$ in \mathcal{B} there are diagrams

$$(B.1) \quad \begin{array}{ccc} \Delta_b \mid \Psi(X) & \xrightarrow{\tau} & \Psi(\Delta_b \mid X) \\ & \searrow \lambda_{\Psi_a} \quad \swarrow \Psi(\lambda) & \\ & \Psi(X) & \end{array}$$

$$(B.2) \quad \begin{array}{ccc} \Psi(X) \mid \Delta_a & \xrightarrow{\tau} & \Psi(X \mid \Delta_a) \\ & \searrow \rho \quad \swarrow \Psi(\rho) & \\ & \Psi(X) & \end{array}$$

whose commutativity expresses compatibility of the functors Ψ with the units, while for

a composable triple X, Y, Z the commutativity of the diagrams

$$\begin{array}{ccc}
\text{(B.3)} & (Z | Y) | \Psi(X) & \xrightarrow{\alpha} Z | (Y | \Psi(X)) \\
& \downarrow \tau & \downarrow \tau \\
& \Psi((Z | Y) | X) & \xrightarrow{\alpha} \Psi(Z | (Y | X)) \\
\text{(B.4)} & (Z | \Psi(Y)) | X & \xrightarrow{\alpha} Z | (\Psi(Y) | X) \\
& \downarrow \tau & \downarrow \tau \\
& \Psi(Z | Y) | X & \downarrow \tau \\
& \downarrow \tau & \downarrow \tau \\
& \Psi((Z | Y) | X) & \xrightarrow{\alpha} \Psi(Z | (Y | X)) \\
\text{(B.5)} & (\Psi(Z) | Y) | X & \xrightarrow{\alpha} \Psi(Z) | (Y | X) \\
& \downarrow \tau & \downarrow \tau \\
& \Psi(Z | Y) | X & \downarrow \tau \\
& \downarrow \tau & \downarrow \tau \\
& \Psi((Z | Y) | X) & \xrightarrow{\alpha} \Psi(Z | (Y | X))
\end{array}$$

expresses compatibility of Ψ with the associator of \mathcal{B} .

Lemma B.1. *Given a bicategory \mathcal{B} with the structure of a supercategory on $\mathcal{B}(a, b)$ for all pairs a, b , and natural isomorphisms τ , suppose that (B.1)-(B.5) commute. Then with Ψ, ξ and γ defined as above, \mathcal{B} is a superbicategory.*

Proof. The proof is an easy but somewhat lengthy exercise, which we omit. \square

References

- [1] D. W. Barnes and L. A. Lambe, *A fixed point approach to homological perturbation theory*, Proc. Amer. Math. Soc. **112** Number 3 (1991), 881–892.
- [2] J. Bénabou, *Introduction to bicategories*, Reports of the Midwest Category Seminar, Springer, 1967.
- [3] F. Borceux, *Handbook of categorical algebra 1*, volume 50 of *Encyclopedia of Mathematics and its Applications*, Cambridge University Press, Cambridge, 1994.

- [4] R.-O. Buchweitz and H. Flenner, *A semiregularity map for modules and applications to deformations*, *Compositio Math.* **137** (2003), 135–210.
- [5] A. Căldăraru and S. Willerton, *The Mukai pairing, I: a categorical approach*, *New York Journal of Mathematics* **16** (2010), 61–98, [[arXiv:0707.2052](#)].
- [6] N. Carqueville and D. Murfet, *Computing Khovanov-Rozansky homology and defect fusion*, *Algebraic & Geometric Topology* **14** (2014), 489–537, [[arXiv:1108.1081](#)].
- [7] N. Carqueville and D. Murfet, *Adjunctions and defects in Landau-Ginzburg models*, [[arXiv:1208.1481](#)].
- [8] N. Carqueville and D. Murfet, *A toolkit for defect computations in Landau-Ginzburg models*, [[arXiv:1303.1389](#)], contribution to the String-Math 2012 proceedings.
- [9] N. Carqueville, A. Ros Camacho and I. Runkel, *Orbifold equivalent potentials*, [[arXiv:1311.3354](#)].
- [10] N. Carqueville and I. Runkel, *On the monoidal structure of matrix bifactorisations*, *J. Phys. A: Math. Theor.* **43** (2010), 275401, [[arXiv:0909.4381](#)].
- [11] ———, *Orbifold completion of defect bicategories*, [[arXiv:1210.6363](#)].
- [12] M. Crainic, *On the perturbation lemma, and deformations*, [[arXiv:0403266](#)].
- [13] A. Davydov, L. Kong, and I. Runkel, *Field theories with defects and the centre functor*, *Mathematical Foundations of Quantum Field Theory and Perturbative String Theory*, *Proceedings of Symposia in Pure Mathematics*, AMS, 2011, [[arXiv:1107.0495](#)].
- [14] T. Dyckerhoff, *Compact generators in categories of matrix factorizations*, *Duke Math. J.* **159** (2011), 223–274, [[arXiv:0904.4713](#)].
- [15] T. Dyckerhoff and D. Murfet, *Pushing forward matrix factorisations*, *Duke Math. J.* **162** (2013), 1249–1311, [[arXiv:1102.2957](#)].
- [16] A. Ellis and A. Lauda, *An odd categorification of $U_q(\mathfrak{sl}_2)$* , [[arXiv:1307.7816](#)].
- [17] T. Friedrich, *Dirac operations in Riemannian geometry*, *Graduate studies in mathematics* **25**, AMS (1997).
- [18] J. W. Gray, *Formal category theory: adjointness for 2-categories*, Springer, Berlin, 1974.
- [19] S.-J. Kang, M. Kashiwara and S.-J. Oh, *Supercategorification of quantum Kac-Moody algebras*, *Adv. Math.* **242** (2013) 116–162, [[arXiv:1206.5933](#)].

- [20] ———, *Supercategorification of quantum Kac-Moody algebras II*, [\[arXiv:1303.1916\]](#).
- [21] A. Kapustin, *Topological Field Theory, Higher Categories, and Their Applications*, [\[arXiv:1004.2307\]](#).
- [22] G. M. Kelly and R. Street, *Review of the elements of 2-categories*, In Category Seminar (Proc. Sem., Sydney, 1972/1973), pages 75–103, Lecture Notes in Math., Vol. 420, Springer, Berlin, 1974.
- [23] M. Khovanov and L. Rozansky, *Matrix factorizations and link homology*, Fund. Math. **199** (2008), 1–91, [\[math/0401268\]](#).
- [24] S. Lack, *A 2-categories companion*, Towards Higher Categories, The IMA Volumes in Mathematics and its Applications **152** (2010), 105–191, [\[math/0702535\]](#).
- [25] C. I. Lazaroiu, *On the boundary coupling of topological Landau-Ginzburg models*, JHEP 0505 (2005), no. 037 [[hep-th/0312286](#)].
- [26] D. McNamee, *On the mathematical structure of topological defects in Landau-Ginzburg models*, MSc Thesis, Trinity College Dublin, 2009.
- [27] D. Murfet, *Computing with cut systems II*, in progress.
- [28] ———, *Logic and linear algebra: an introduction*, [\[arXiv:1407.2650\]](#).
- [29] I. Runkel and R. Suszek, *Gerbe-holonomy for surfaces with defect networks* Advances in Theoretical and Mathematical Physics 13.4 (2009): 1137–1219, [\[arXiv:0808.1419\]](#).
- [30] A. Vistoli, *Notes on Grothendieck topologies, fibered categories and descent theory*, [\[arXiv:0412512\]](#).
- [31] Y. Yoshino, *Tensor products of matrix factorizations*, Nagoya Math. J. **152** (1998), 39–56.