

Modelling malaria transmission and control

Sebastian Funk

Centre for the mathematical modelling of infectious diseases
London School of Hygiene & Tropical Medicine

Modelling malaria transmission and control

Objectives of this lecture

- Become familiar with the Ross-MacDonald model for malaria transmission
- Appreciate the key features of malaria epidemiology and how these affect transmission dynamics
- Understand the basic reproductive ratio of malaria and how its components are linked to control

Modelling malaria transmission and control

Ronald Ross



* 1857, Almora, India
† 1932, London

1897: Malaria in mosquito
1902: Nobel Prize

"...With tears and toiling breath,
I find thy cunning seeds,
O million-murdering Death."

Modelling malaria transmission and control

Ronald Ross

A priori method

"we assume a knowledge of the causes, construct our differential equations on that supposition, follow up the logical consequences, and finally test the calculated results by comparing them with the observed statistics."



A posteriori method


"we commence with observed statistics, endeavour to fit analytical laws to them, and so work backwards to the underlying cause."

Ross (1916-1917)

An application of the theory of probabilities to the study of a priori pathometry. I-III
Philos Trans R Soc Lond A

Modelling malaria transmission and control

George MacDonald



* 1903, Sheffield
† 1967, London

Modelling malaria transmission and control

An overview of malaria

- Malaria parasite = the protozoan *Plasmodium*
- 5 species which infect humans:
 - *P. falciparum*: most dangerous (cerebral malaria/severe anaemia)
 - *P. vivax*: milder disease but still significant (relapsing)
 - *P. ovale* (relapsing), *P. malariae*: rarer, mostly benign disease
 - *P. knowlesi*: zoonosis from monkeys (Singh & Cox-Singh 2004)
- Ancient disease, ~50,000-100,000 years old
 - Strong force in human evolution

Modelling malaria transmission and control

Equations of a basic Ross-MacDonald model



$$\begin{cases} \frac{dS_H}{dt} = rI_H - I_H S_H \\ \frac{dI_H}{dt} = I_H S_H - rI_H \end{cases}$$

λ_H : force of infection on humans
 r : recovery rate from Malaria

S_H, I_H, S_M, I_M are the **proportions** of humans/mosquitoes that are susceptible/infected, i.e., I_H is the prevalence of infection in humans.



$$\begin{cases} \frac{dS_M}{dt} = m - I_M S_M - mS_M \\ \frac{dI_M}{dt} = I_M S_M - mI_M \end{cases}$$

λ_M : force of infection on mosquitoes
 μ : mosquito death rate

Modelling malaria transmission and control

Force of infection: humans

 λ_H

From the viewpoint of a susceptible human:
The rate at which I'll get infected with malaria



= (rate at which infected mosquitoes bite me and transmit infection)

= (rate at which I get bitten by mosquitoes) ·
 (proportion of mosquitoes that are infected) ·
 (probability that I get infected when bitten by an infected mosquito)

= (rate at which one mosquito bites) · (number of mosquitoes) ·
 (probability that a biting mosquito chooses me) ·
 (proportion of mosquitoes that are infected) ·
 (probability that I get infected when bitten by an infected mosquito)

Modelling malaria transmission and control

Force of infection: humans

 λ_H

From the viewpoint of a susceptible human:
The rate at which I'll get infected with malaria



$$I_H = aM \frac{1}{H} I_M b$$

= (rate at which one mosquito bites) · (number of mosquitoes) ·
 (probability that a biting mosquito chooses me) ·
 (proportion of mosquitoes that are infected) ·
 (probability that I get infected when bitten by an infected mosquito)

Modelling malaria transmission and control

Force of infection: humans

Entomological
 inoculation rate
 (EIR)

$$\lambda_H = a m I_M b$$



Where

- a = biting rate
 (number of bites taken on humans per mosquito per time unit)
- m = number of mosquitoes per human ($m=M/H$)
- I_M = prevalence of infection in mosquitoes
- b = probability that a bite by an infectious mosquito leads to successful mosquito-to-human transmission

Force of infection on humans depends on prevalence in mosquitoes

Modelling malaria transmission and control

Equations of a basic Ross-MacDonald model



$$\begin{cases} \frac{dS_H}{dt} = rI_H - I_H S_H \\ \frac{dI_H}{dt} = I_H S_H - rI_H \end{cases}$$

λ_H : force of infection on humans
 r : recovery rate from Malaria

S_H, I_H, S_M, I_M are the **proportions** of humans/mosquitoes that are susceptible/infected, i.e., I_H is the prevalence of infection in humans.



$$\begin{cases} \frac{dS_M}{dt} = m - I_M S_M - mS_M \\ \frac{dI_M}{dt} = I_M S_M - mI_M \end{cases}$$

λ_M : force of infection on mosquitoes
 μ : mosquito death rate

Modelling malaria transmission and control

Force of infection: mosquitoes

 λ_M

From the viewpoint of a susceptible mosquito:
The rate at which I'll get infected with malaria



= (rate at which I bite infectious humans and become infected)

= (rate at which I bite humans) ·
 (probability that a human is infected) ·
 (probability that I become infected through the bite)

$$I_M = a I_H c$$

Modelling malaria transmission and control

Force of infection: mosquitoes

$$\lambda_M = acI_H$$



Where

- a = biting rate (as above)
- c = probability that a bite taken on infectious human leads to successful human-to-mosquito transmission
- I_H = prevalence of infectious humans

Force of infection on mosquitoes depends on prevalence in humans

Modelling malaria transmission and control

Equations of a basic Ross-MacDonald model



$$\begin{cases} \frac{dS_H}{dt} = rI_H - I_H S_H \\ \frac{dI_H}{dt} = I_H S_H - rI_H \end{cases}$$

S_H, I_H, S_M, I_M are the **proportions** of humans/mosquitoes that are susceptible/infected, i.e., I_H is the prevalence of infection in humans.



$$\begin{cases} \frac{dS_M}{dt} = m - I_M S_M - mS_M \\ \frac{dI_M}{dt} = I_M S_M - mI_M \end{cases}$$

Modelling malaria transmission and control

Equations of a basic Ross-MacDonald model



$$\begin{cases} \frac{dS_H}{dt} = rI_H - amI_M bS_H \\ \frac{dI_H}{dt} = amI_M bS_H - rI_H \end{cases}$$


S_H, I_H, S_M, I_M are the **proportions** of humans/mosquitoes that are susceptible/infected, i.e., I_H is the prevalence of infection in humans.



$$\begin{cases} \frac{dS_M}{dt} = m - acI_H S_M - mS_M \\ \frac{dI_M}{dt} = acI_H S_M - mI_M \end{cases}$$


Modelling malaria transmission and control

Model dynamics




Modelling malaria transmission and control

Model age at infection









Modelling malaria transmission and control

Model impact of control



Modelling malaria transmission and control



Using R_0 to assess control

$$R_0 = m \frac{a^2}{m} p^n bc \frac{1}{r}$$

Larval control
 Adult vector control
 Vaccination or prophylaxis to block infection
 Treating infection
 Drugs or vaccines that block transmission

$$R_0 = \frac{ma^2bcp^n}{rm}$$

Modelling malaria transmission and control

Objectives of this lecture

- Become familiar with the Ross-MacDonald model for malaria transmission
- Appreciate the key features of malaria epidemiology and how these affect transmission dynamics
- Understand the basic reproductive ratio of malaria and how its components are linked to control

Smith DL et al. (2012).
 Ross, Macdonald, and a Theory for the Dynamics and Control of Mosquito-Transmitted Pathogens.
 PLoS Pathog 8(4): e1002588. <http://dx.doi.org/10.1371/journal.ppat.1002588>

Modelling malaria transmission and control