

Math 105: Calculus with Algebra
Instructor: David Zureick-Brown (“DZB”)

All assignments

Gradescope code: **B5J8EN**

Gradescope instructions	2
1 Assignment 1, Due Friday, September 4	4
2 Assignment 1 Solutions	6
3 Assignment 2, Due Monday, September 7	9
4 Assignment 2 Solutions	10
5 Assignment 3, Due Friday, September 11	13
6 Assignment 3 Solutions	16
7 Assignment 4, Due Monday, September 14	21
8 Assignment 4 Solutions	23
9 Assignment 5, Due Friday, September 19	27
10 Assignment 5 Solutions	29
11 Assignment 6, Due Monday, September 21	34
12 Assignment 6 Solutions	36
13 Assignment 7, Due Friday, September 25	40
14 Assignment 7 Solutions	42
15 Assignment 8, Due Monday, September 28	46
16 Assignment 8 Solutions	48
17 Midterm 1 Review Packet	53

Gradescope Instructions for submitting work in Math 105

You will be using the online Gradescope program to submit your homework and exams. These instructions tell you how to sign up initially, and how to submit your written work.

Signing up for Gradescope the first time.

If you haven't used Gradescope for an **Amherst College** course before:

- Go to <http://www.gradescope.com>, click on "Sign up for free" (which may auto-scroll you to the bottom of the page), and select Sign up as [a] "Student".
- In the signup box:
 - Use the course entry code **B5J8EN**
 - Use your full name
 - Use your **Amherst College email** address. Or, if you are a Five-College student, use your email address from your own school.
 - Leave the "Student ID" entry blank.
- You will probably get an email asking to set a password for your account, so check your **amherst.edu** email inbox. (Or your email inbox through your own school, for Five-College students.)

Adding Math 105 to Gradescope.

If you **have** used Gradescope for an Amherst course before, and so you already have an account through your **amherst.edu** email, you still need to add Math 105, so:

- Go to <http://www.gradescope.com> and log in.
- Go to your Account Dashboard (click the Gradescope logo at upper left), and click "Add Course" at bottom right.
- Use the course code **B5J8EN**

(submission instructions on next page)

Submitting written work

First write it out on paper as you would normally. Then **scan it** to create a PDF. One method for scanning is the smartphone app **DropBox**. It makes nice clear scans, and it saves them directly into a folder so that you can have all your assignments in one place. **CamScanner** is another free scanning App, and there are others, too. **Gradescope** now has its own scanning app. You can also use a printer/scanner if you prefer.

Please be kind to our dear graders and make sure your submission is legible !

In particular, please leave some spacing between separate problems.

If you have a tablet computer, you may write your work there (instead of on paper) and save it as a PDF.

Some of you may know the math formatting package LaTeX and may want to use it in Math 105. That's fine, too; if so, you may write up your work in LaTeX and save the resulting PDF.

In short, any method is fine as long as it creates a legible PDF file and NOT a photo.

For example, if you use the DropBox app, then in your created *Math 105 Homework* Dropbox folder, you can select create (+) at the bottom of the screen and click the *Scan Document* option. Snap a shot of the first page of your homework, and then click [+] to snap shots of any subsequent pages. Do **not** use the *Take Photo* option.

After you have scanned/saved your work as a PDF, submit it on Gradescope as follows:

- Go to <http://www.gradescope.com> and log in.
- Select the course “Math 105, Fall 2026” and the appropriate assignment.
- Select “submit pdf” to submit your work in PDF format. Browse to find your PDF and upload.
- Now it is time to **tag** your problems. **This is an important step**, where you are telling Gradescope which problems are on which page(s).

For each problem, select the pages of your submission where your written solution appears.

I think the easiest thing to do is to click on the page of **your** homework upload where you wrote the given problem, and then click on the assigned problem listed. Repeat for each problem.

You must tag the problems or else you will not get credit for your work.

Gradescope will give you a warning when you go to submit your assignment if you have not selected the pages correctly. If you tag a problem incorrectly, you can fix it by clicking “More” and “Reselect Pages”.

- Click Save or Submit.

After your assignment is graded, you will be able to see your score on the written problems, along with comments, on Gradescope. You should receive an email notifying you when each homework set is graded.

MATH 105 CALCULUS WITH ALGEBRA FALL 2026
HOMEWORK 1

Due **Friday, September 4** at 11:55pm, on Gradescope

GOALS:

- Review of basic algebra.
- Review of absolute values and inequalities.
- Write complete solutions.

Justify your solution, writing down the steps that led you to the final answer. For a review on inequalities and absolute value, see Appendix A (Stewart Calculus book).

(1) Simplify:

$$4 - (5 - 3) + 3(4 - 7) - 4(1 + 2)$$

(2) In the following expressions, simplify and reduce to lowest terms. (Some answers may have more than one form)

(a)

$$\frac{\frac{4}{75}}{\frac{8}{25}}$$

(b)

$$\frac{7x}{3y} \cdot \frac{3y + 2}{x}$$

(c)

$$\frac{\frac{xy}{(x+y)}}{\frac{x^2y}{(x+y)^3}}$$

MORE PROBLEMS ON THE NEXT PAGE →

- (3) Express as a single fraction and simplify:

(a)

$$\frac{1}{2} + \frac{4}{3} - \frac{2}{5} - \frac{3}{15}$$

(b)

$$\frac{\frac{1}{3} + \frac{2}{5}}{\frac{3}{2}}$$

(c)

$$\frac{1}{x} - \frac{x+1}{xy} + \frac{x-2}{xz}$$

(d)

$$\frac{\frac{1}{st} - \frac{1}{w}}{\frac{1}{tw} - \frac{2}{s}}$$

- (4) Rewrite the expression without using the absolute-value symbol:

$$|x - 2| \text{ if } x < 2$$

- (5) Solve the inequalities with the **geometric interpretation** of absolute value.

(a)

$$|x| < 3$$

(b)

$$|x - 4| < 1$$

(c)

$$|x + 5| \geq 2$$

MATH 105 CALCULUS WITH ALGEBRA FALL 2026
HOMEWORK 1 SOLUTIONS

- (1) Simplify:

$$4 - (5 - 3) + 3(4 - 7) - 4(1 + 2)$$

Solution. We simplify the expressions inside parentheses first:

$$4 - (5 - 3) + 3(4 - 7) - 4(1 + 2) = 4 - 2 + 3(-3) - 4(3).$$

Therefore

$$4 - 2 - 9 - 12 = -19.$$

Hence the simplified value is $\boxed{-19}$.

- (2) In the following expressions, simplify and reduce to lowest terms. (Some answers may have more than one form)

- (a)

$$\frac{\frac{4}{75}}{\frac{8}{25}}$$

Solution. Dividing by a fraction is the same as multiplying by its reciprocal:

$$\frac{\frac{4}{75}}{\frac{8}{25}} = \frac{4}{75} \cdot \frac{25}{8} = \frac{1}{3} \cdot \frac{1}{2} = \boxed{\frac{1}{6}}.$$

- (b)

$$\frac{7x}{3y} \cdot \frac{3y+2}{x}$$

Solution. Cancel the common factor x :

$$\frac{7x}{3y} \cdot \frac{3y+2}{x} = \boxed{\frac{7(3y+2)}{3y}}.$$

- (c)

$$\frac{\frac{xy}{(x+y)}}{\frac{x^2y}{(x+y)^3}}$$

Solution. Multiply by the reciprocal of the denominator:

$$\frac{\frac{xy}{(x+y)}}{\frac{x^2y}{(x+y)^3}} = \frac{xy}{x+y} \cdot \frac{(x+y)^3}{x^2y}.$$

Cancelling common factors gives

$$\boxed{\frac{(x+y)^2}{x}}.$$

- (3) Express as a single fraction and simplify:

(a)

$$\frac{1}{2} + \frac{4}{3} - \frac{2}{5} - \frac{3}{15}$$

Solution. A common denominator is 30:

$$\frac{1}{2} + \frac{4}{3} - \frac{2}{5} - \frac{3}{15} = \frac{15}{30} + \frac{40}{30} - \frac{12}{30} - \frac{6}{30} = \boxed{\frac{37}{30}}.$$

(b)

$$\frac{\frac{1}{3} + \frac{2}{5}}{\frac{3}{2}}$$

Solution. First simplify the numerator:

$$\frac{1}{3} + \frac{2}{5} = \frac{5}{15} + \frac{6}{15} = \frac{11}{15}.$$

Then divide by $3/2$:

$$\frac{\frac{11}{15}}{\frac{3}{2}} = \frac{11}{15} \cdot \frac{2}{3} = \boxed{\frac{22}{45}}.$$

(c)

$$\frac{1}{x} - \frac{x+1}{xy} + \frac{x-2}{xz}$$

Solution. The common denominator is xyz . Thus

$$\frac{1}{x} - \frac{x+1}{xy} + \frac{x-2}{xz} = \frac{yz}{xyz} - \frac{z(x+1)}{xyz} + \frac{y(x-2)}{xyz}.$$

Combining the numerators,

$$\boxed{\frac{xy + yz - xz - 2y - z}{xyz}}.$$

(d)

$$\frac{\frac{1}{st} - \frac{1}{w}}{\frac{1}{tw} - \frac{2}{s}}$$

Solution. Put the numerator and denominator over the common denominator stw :

$$\frac{1}{st} - \frac{1}{w} = \frac{w-st}{stw}, \quad \frac{1}{tw} - \frac{2}{s} = \frac{s-2tw}{stw}.$$

Therefore

$$\frac{\frac{w-st}{stw}}{\frac{s-2tw}{stw}} = \boxed{\frac{w-st}{s-2tw}}.$$

(4) Rewrite the expression without using the absolute-value symbol:

$$|x-2| \text{ if } x < 2$$

Solution. If $x < 2$, then $x-2 < 0$. The absolute value of a negative number is its opposite, so

$$|x-2| = -(x-2) = \boxed{2-x}.$$

(5) Solve the inequalities with the **geometric interpretation** of absolute value.

(a)

$$|x| < 3$$

Solution. The quantity $|x|$ is the distance from x to 0. We want that distance to be less than 3, so x must lie between -3 and 3 :

$$\boxed{-3 < x < 3}.$$

(b)

$$|x - 4| < 1$$

Solution. The quantity $|x - 4|$ is the distance from x to 4. We want that distance to be less than 1, so

$$\boxed{3 < x < 5}.$$

(c)

$$|x + 5| \geq 2$$

Solution. Since $|x + 5| = |x - (-5)|$, this is the distance from x to -5 . We want that distance to be at least 2. Thus x must be at least 2 units to the left or right of -5 :

$$\boxed{x \leq -7 \quad \text{or} \quad x \geq -3}.$$

MATH 105 CALCULUS WITH ALGEBRA FALL 2026
HOMEWORK 2

Due **Monday, September 7** at 11:55pm, on Gradescope

GOALS:

- Review equations of lines.
- Write complete solutions.

Justify your solution, writing down the steps that led you to the final answer. For a review on equations of lines, see Appendix A (Stewart Calculus book).

Find the slope of the line through P and Q :

(1) $P(1, 5), Q(4, 11)$

(2) $P(-1, -4), Q(6, 0)$

Find the equation of the line that satisfies the given conditions.

(3) Through $(2, -3)$ with slope 6

(4) Through $(2, 1)$ and $(1, 6)$

(5) Slope 3, y -intercept -2

(6) x -intercept 1, y -intercept -3

(7) Through $(4, 5)$, parallel to x -axis

(8) Through $(4, 5)$, parallel to y -axis

(9) Through $(1, -6)$, parallel to the line $x + 2y = 6$

(10) Through $(-1, -2)$, perpendicular to the line $2x + 5y + 8 = 0$

MATH 105 CALCULUS WITH ALGEBRA FALL 2026
HOMEWORK 2 SOLUTIONS

Find the slope of the line through P and Q :

- (1) $P(1, 5), Q(4, 11)$

Solution. The slope through two points (x_1, y_1) and (x_2, y_2) is

$$m = \frac{y_2 - y_1}{x_2 - x_1}.$$

Thus

$$m = \frac{11 - 5}{4 - 1} = \frac{6}{3} = \boxed{2}.$$

- (2) $P(-1, -4), Q(6, 0)$

Solution. Using the slope formula,

$$m = \frac{0 - (-4)}{6 - (-1)} = \frac{4}{7}.$$

Therefore the slope is

$$\boxed{\frac{4}{7}}.$$

Find the equation of the line that satisfies the given conditions.

- (3) Through $(2, -3)$ with slope 6

Solution. Using point-slope form $y - y_1 = m(x - x_1)$,

$$y - (-3) = 6(x - 2).$$

Hence

$$y + 3 = 6x - 12,$$

so

$$\boxed{y = 6x - 15}.$$

- (4) Through $(2, 1)$ and $(1, 6)$

Solution. First find the slope:

$$m = \frac{6 - 1}{1 - 2} = \frac{5}{-1} = -5.$$

Using the point $(2, 1)$,

$$y - 1 = -5(x - 2).$$

Therefore

$$y - 1 = -5x + 10,$$

so

$$\boxed{y = -5x + 11}.$$

- (5) Slope 3, y -intercept -2

Solution. A line with slope m and y -intercept b has equation $y = mx + b$. Here $m = 3$ and $b = -2$, so

$$\boxed{y = 3x - 2}.$$

- (6) x -intercept 1, y -intercept -3

Solution. The x -intercept gives the point $(1, 0)$, and the y -intercept gives the point $(0, -3)$. The slope is

$$m = \frac{0 - (-3)}{1 - 0} = 3.$$

Since the y -intercept is -3 , the equation is

$$\boxed{y = 3x - 3}.$$

- (7) Through $(4, 5)$, parallel to x -axis

Solution. A line parallel to the x -axis is horizontal, so its y -coordinate is constant. Since the line passes through $(4, 5)$,

$$\boxed{y = 5}.$$

- (8) Through $(4, 5)$, parallel to y -axis

Solution. A line parallel to the y -axis is vertical, so its x -coordinate is constant. Since the line passes through $(4, 5)$,

$$\boxed{x = 4}.$$

- (9) Through $(1, -6)$, parallel to the line $x + 2y = 6$

Solution. Rewrite the given line in slope-intercept form:

$$x + 2y = 6 \implies 2y = -x + 6 \implies y = -\frac{1}{2}x + 3.$$

Its slope is $-\frac{1}{2}$. A parallel line has the same slope, so using point-slope form with $(1, -6)$,

$$y + 6 = -\frac{1}{2}(x - 1).$$

Therefore

$$\boxed{y = -\frac{1}{2}x - \frac{11}{2}}.$$

Equivalently, the equation can be written as $\boxed{x + 2y = -11}$.

- (10) Through $(-1, -2)$, perpendicular to the line $2x + 5y + 8 = 0$

Solution. First rewrite the given line in slope-intercept form:

$$2x + 5y + 8 = 0 \implies 5y = -2x - 8 \implies y = -\frac{2}{5}x - \frac{8}{5}.$$

Its slope is $-\frac{2}{5}$. A perpendicular line has slope equal to the negative reciprocal, so its slope is

$$\frac{5}{2}.$$

Using point-slope form with $(-1, -2)$,

$$y + 2 = \frac{5}{2}(x + 1).$$

Thus

$$y + 2 = \frac{5}{2}x + \frac{5}{2},$$

and therefore

$$\boxed{y = \frac{5}{2}x + \frac{1}{2}}.$$

MATH 105 CALCULUS WITH ALGEBRA FALL 2026
HOMEWORK 3

Due **Friday, September 11** at 11:55pm, on Gradescope

GOALS:

- Review of the essential ideas behind functions.
- Practice graphing of functions.
- Write complete solutions.

Justify your solution, writing down the steps that led you to the final answer.

(1) Let $f(x) = x^3 + 2x$. Determine:

(a) $f(2)$

(b) $f(2x)$

(c) $f(-x)$

(d) $f(2 + \Delta x)$

(2) Let $u(t) = \frac{1}{2}t^2 + t$. Determine:

(a) $u(-1)$

(b) $u(t + \Delta t)$


(c) $u(-t)$

(3) Sketch the graph of the function $g(t) = |1 - 3t|$


(4) Sketch the graph of the function

$$f(x) = \begin{cases} |x| & \text{if } |x| \leq 1 \\ 1 & \text{if } |x| > 1 \end{cases}$$


- (5) The graph of the function f is given.




- (a) State the value of $f(1.5)$.
- (b) Estimate the value of $f(-1)$.
- (c) For what values of x is $f(x) = 1$?
- (d) Estimate the value of x such that $f(x) = 0$.
- (6) Determine whether the curve is a graph of a function of x . If it is, state the domain and the range of the function.



- (7) State the domain and the range of the function graphed below.



- (8) Find the domain of $f(x) = \frac{x+4}{x^2-9}$
- (9) Match the each equation with its graph. Explain your choices. Don't use a computer or graphic calculator. (a) $y = x^2$ (b) $y = x^5$ (c) $y = x^8$



MATH 105 Calculus with Algebra – Fall 2026

Homework 3 – Solutions

1. Let $f(x) = x^3 + 2x$. Determine:

(a) $f(2)$.

Solution.

$$f(2) = 2^3 + 2(2) = 8 + 4 = \boxed{12}.$$

(b) $f(2x)$.

Solution.

$$f(2x) = (2x)^3 + 2(2x) = \boxed{8x^3 + 4x}.$$

(c) $f(-x)$.

Solution.

$$f(-x) = (-x)^3 + 2(-x) = \boxed{-x^3 - 2x} = -f(x).$$

(d) $f(2 + \Delta x)$.

Solution.

$$\begin{aligned} f(2 + \Delta x) &= (2 + \Delta x)^3 + 2(2 + \Delta x) \\ &= 8 + 12\Delta x + 6(\Delta x)^2 + (\Delta x)^3 + 4 + 2\Delta x \\ &= \boxed{12 + 14\Delta x + 6(\Delta x)^2 + (\Delta x)^3}. \end{aligned}$$

2. Let $u(t) = \frac{1}{2}t^2 + t$. Determine:

(a) $u(-1)$.

Solution.

$$u(-1) = \frac{1}{2}(-1)^2 + (-1) = \frac{1}{2} - 1 = \boxed{-\frac{1}{2}}.$$

(b) $u(t + \Delta t)$.

Solution.

$$\begin{aligned} u(t + \Delta t) &= \frac{1}{2}(t + \Delta t)^2 + (t + \Delta t) \\ &= \frac{1}{2}t^2 + t\Delta t + \frac{1}{2}(\Delta t)^2 + t + \Delta t \\ &= \boxed{\frac{1}{2}t^2 + t + (t + 1)\Delta t + \frac{1}{2}(\Delta t)^2}. \end{aligned}$$

(c) $u(-t)$.

Solution.


$$u(-t) = \frac{1}{2}(-t)^2 - t = \boxed{\frac{1}{2}t^2 - t}.$$

3. Sketch $g(t) = |1 - 3t|$.

Solution. Since $1 - 3t = 0$ when $t = \frac{1}{3}$,

$$g(t) = \begin{cases} 1 - 3t, & t \leq \frac{1}{3}, \\ 3t - 1, & t \geq \frac{1}{3}. \end{cases}$$


Thus the graph is a V with vertex $(\frac{1}{3}, 0)$, left slope -3 , and right slope 3 .




4. Sketch

$$f(x) = \begin{cases} |x|, & |x| \leq 1, \\ 1, & |x| > 1. \end{cases}$$

Solution. On $[-1, 1]$ the graph is the usual V-shaped graph $y = |x|$. Outside that interval the graph is the horizontal line $y = 1$. The pieces meet at $(-1, 1)$ and $(1, 1)$.



5. The graph of f is shown below.



(a) State $f(1.5)$.

Solution. From the graph, $f(1.5) \approx 2.5$.

(b) Estimate $f(-1)$.

Solution. From the graph, $f(-1) \approx -0.6$ (an estimate near -0.5 to -0.6 is reasonable from the hand-drawn graph).


(c) For what values of x is $f(x) = 1$?

Solution. The horizontal line $y = 1$ meets the graph at $x = 0$, so $x = 0$.

(d) Estimate the value of x such that $f(x) = 0$.


Solution. The x -intercept is approximately $x \approx -0.4$.

6. Determine whether the curve below is the graph of a function of x . If it is, state its domain and range.



Solution. It is not the graph of a function of x . A vertical line through some x -values intersects the red curve more than once, so it fails the vertical line test.

7. State the domain and range of the function graphed below.



Solution. The graph begins at the closed point with $x = -2$ and ends at the closed point with $x = 3$, so

$$\text{Domain} = [-2, 3].$$

Its lowest value is $y = -1$ and its highest value is $y = 2$, so

$$\boxed{\text{Range} = [-1, 2]}.$$

8. Find the domain of

$$f(x) = \frac{x+4}{x^2-9}.$$


Solution. The denominator cannot equal zero:

$$x^2 - 9 = (x-3)(x+3) \neq 0.$$

Thus $x \neq -3, 3$, and

$$\boxed{\text{Domain} = (-\infty, -3) \cup (-3, 3) \cup (3, \infty)}.$$

9. Match each equation with its graph: (a) $y = x^2$, (b) $y = x^5$, (c) $y = x^8$.



Solution. The red curve f takes negative values for negative x and positive values for positive x , so it is the odd power:

$$\boxed{f(x) = x^5}.$$

The blue curve h is the wider even-power graph, while the green curve g is flatter near 0 and grows more steeply for $|x| > 1$. Therefore

$$\boxed{h(x) = x^2, \quad g(x) = x^8}.$$

So the matching is

$$\boxed{\text{(a) } h, \quad \text{(b) } f, \quad \text{(c) } g}.$$

MATH 105 CALCULUS WITH ALGEBRA FALL 2026
HOMEWORK 4

Due **Monday, September 14** at 11:55pm, on Gradescope

GOALS:

- Practice graphing of functions.
- Practice shifting and stretching graphs.
- Simplify difference quotients.
- Practice compositions of functions.

- Write complete solutions.

Justify your solution, writing down the steps that led you to the final answer.

- (1) The graph of $y = f(x)$ is given. Match each equation with its graph and give reasons for your choices.

(a)

$$y = f(x - 4)$$

(b)

$$y = f(x) + 3$$

(c)


$$y = \frac{1}{3}f(x)$$

(d)

$$y = -f(x + 4)$$

(e)

$$y = 2f(x + 4)$$



- (2) Graph the following functions by hand, not by plotting points, but by starting with the graph of one of the standard functions (see Stewart, Section 1.2) and then applying the appropriate transformations.

(a) $y = -x^2$

(b) $1 - \frac{1}{x}$

(c) $|x - 2|$

- (3) Let $f(x) = \frac{x+1}{x}$.

(a) Compute $f(a + h)$

(b) Simplify

$$\frac{f(a + h) - f(a)}{h}$$

- (4) Find $f(g(h(x)))$:

$$f(x) = \sqrt{x - 3}, \quad g(x) = x^2, \quad h(x) = x^3 + 2$$


- (5) Express the function in the form $f(g(x))$:

$$F(x) = (2x + x^2)^4$$

(In other words, find functions $f(x)$ and $g(x)$ such that $F(x) = f(g(x))$).

MATH 105 CALCULUS WITH ALGEBRA FALL 2026
HOMEWORK 4 SOLUTIONS

- (1) The graph of $y = f(x)$ is given. Match each equation with its graph and give reasons for your choices.




- (a) $y = f(x - 4)$ is f shifted four units to the right, so it is (c).
- (b) $y = f(x) + 3$ is f shifted three units up, so it is (a).
- (c) $y = \frac{1}{3}f(x)$ is a vertical compression by a factor of $\frac{1}{3}$, so it is (d).
- (d) $y = -f(x + 4)$ shifts f four units left and reflects it across the x -axis, so it is (e).
- (e) $y = 2f(x + 4)$ shifts f four units left and stretches it vertically by a factor of 2, so it is (b).

- (2) Graph the following functions by hand, starting with a standard graph and applying the appropriate transformations.


(a) $y = -x^2$

Start with $y = x^2$ and reflect it across the x -axis. The vertex remains at $(0, 0)$ and the parabola opens downward.




(b) $y = 1 - \frac{1}{x}$

Start with $y = \frac{1}{x}$, reflect across the x -axis, and then shift up by 1. The vertical asymptote is $x = 0$ and the horizontal asymptote is $y = 1$.



(c) $y = |x - 2|$

Start with $y = |x|$ and shift it two units to the right. The vertex is $\boxed{(2, 0)}$.



(3) Let

$$f(x) = \frac{x+1}{x}.$$

(a) Replacing x by $a + h$ gives

$$\boxed{f(a+h) = \frac{a+h+1}{a+h}}.$$

(b) We simplify the difference quotient:

$$\begin{aligned} \frac{f(a+h) - f(a)}{h} &= \frac{1}{h} \left(\frac{a+h+1}{a+h} - \frac{a+1}{a} \right) \\ &= \frac{1}{h} \left(\frac{a(a+h+1) - (a+1)(a+h)}{a(a+h)} \right) \\ &= \frac{1}{h} \left(\frac{-h}{a(a+h)} \right) \\ &= \boxed{-\frac{1}{a(a+h)}}. \end{aligned}$$

The original expression requires $a \neq 0$, $a + h \neq 0$, and $h \neq 0$.

(4) Find $f(g(h(x)))$, where

$$f(x) = \sqrt{x-3}, \quad g(x) = x^2, \quad h(x) = x^3 + 2.$$

Work from the inside out:

$$h(x) = x^3 + 2,$$

so

$$g(h(x)) = (x^3 + 2)^2.$$

Therefore

$$f(g(h(x))) = \sqrt{(x^3 + 2)^2 - 3}.$$

Thus

$$\boxed{f(g(h(x))) = \sqrt{(x^3 + 2)^2 - 3}}.$$

(5) Express

$$F(x) = (2x + x^2)^4$$

in the form $f(g(x))$.

One natural choice is

$$\boxed{g(x) = 2x + x^2} \quad \text{and} \quad \boxed{f(x) = x^4}.$$

Indeed,

$$f(g(x)) = (g(x))^4 = (2x + x^2)^4 = F(x).$$

Other choices are possible.

MATH 105 CALCULUS WITH ALGEBRA FALL 2026
HOMEWORK 5

Due **Friday, September 18** at 11:55pm, on Gradescope

Justify your solutions, writing down the steps that led you to the final answer.

(1) The graph of f is given. Draw the graphs of the following functions.


(a) $y = f(x) - 3$

(b) $y = f(x + 1)$

(c) $y = \frac{1}{2}f(x)$

(d) $y = -f(x)$

(e) $y = f + f$.



(2) Graph the function $f(x) = -1 + \frac{1}{(2x)^2 - 1}$

(3) Let $f(x) = 1 - x^2$ and let $g(x) = \sqrt{x + 1}$. Compute (and simplify) $f + g$, fg , $f \circ g$, and $g \circ f$.

(4) Explain in your own words what is meant by the equation

$$\lim_{x \rightarrow 2} f(x) = 5.$$

Is it possible for this statement to be true and yet $f(2) = 3$? Explain.


- (5) Explain what it means to say that

$$\lim_{x \rightarrow 1^-} f(x) = 3 \quad \text{and} \quad \lim_{x \rightarrow 1^+} f(x) = 7.$$

In this situation is it possible that $\lim_{x \rightarrow 1} f(x)$ exists? Explain.

- (6) For the function f whose graph is given below, state the value of each quantity, if it exists. If it does not exist, explain why.

- (a) $\lim_{x \rightarrow 2^-} f(x)$
- (b) $\lim_{x \rightarrow 2^+} f(x)$
- (c) $\lim_{x \rightarrow 2} f(x)$
- (d) $f(2)$
- (e) $\lim_{x \rightarrow 4} f(x)$
- (f) $f(4)$
- (g) $\lim_{x \rightarrow 3} f(x)$



- (7) Determine the limits:

$$\lim_{x \rightarrow 5^+} \frac{x+1}{x-5}.$$

and


$$\lim_{x \rightarrow 5^-} \frac{x+1}{x-5}.$$

MATH 105 CALCULUS WITH ALGEBRA – FALL 2026
HOMEWORK 5 SOLUTIONS

(1) The graph of f is given by the piecewise-linear curve through


$$(-2, 2), \quad (0, 0), \quad (1, 3), \quad (3, 2).$$

We transform the entire graph, rather than transforming the individual points one at a time.




(a) $y = f(x) - 3$ shifts the graph of f **down 3 units**. The transformed vertices are

$$(-2, -1), \quad (0, -3), \quad (1, 0), \quad (3, -1).$$




(b) $y = f(x + 1)$ shifts the graph of f **left 1 unit**. The transformed vertices are

$$(-3, 2), \quad (-1, 0), \quad (0, 3), \quad (2, 2).$$




- (c) $y = \frac{1}{2}f(x)$ vertically compresses the graph by a factor of $\frac{1}{2}$. The x -coordinates stay the same, while all y -coordinates are halved:

$$(-2, 1), \quad (0, 0), \quad \left(1, \frac{3}{2}\right), \quad (3, 1).$$




- (d) $y = -f(x)$ reflects the graph across the x -axis. The transformed vertices are

$$(-2, -2), \quad (0, 0), \quad (1, -3), \quad (3, -2).$$



- (e) Since $f + f = 2f$, the graph is a vertical stretch by a factor of 2. The transformed vertices are

$$(-2, 4), \quad (0, 0), \quad (1, 6), \quad (3, 4).$$



(2) Graph

$$f(x) = -1 + \frac{1}{(2x)^2 - 1} = -1 + \frac{1}{4x^2 - 1}.$$

The denominator is zero when

$$4x^2 - 1 = 0,$$

so the vertical asymptotes are

$$\boxed{x = -\frac{1}{2} \quad \text{and} \quad x = \frac{1}{2}}.$$

As $|x|$ becomes large, the fraction $1/(4x^2 - 1)$ approaches 0, so the horizontal asymptote is

$$\boxed{y = -1}.$$

The function is even, so its graph is symmetric about the y -axis. Also,

$$f(0) = -2.$$


For the x -intercepts,

$$\begin{aligned} -1 + \frac{1}{4x^2 - 1} &= 0 \\ \frac{1}{4x^2 - 1} &= 1 \\ 4x^2 &= 2, \end{aligned}$$

so

$$\boxed{x = \pm \frac{1}{\sqrt{2}}}.$$

These features determine the graph:



(3) Let

$$f(x) = 1 - x^2, \quad g(x) = \sqrt{x + 1}.$$

$$\begin{aligned} (f + g)(x) &= f(x) + g(x) \\ &= \boxed{1 - x^2 + \sqrt{x + 1}}. \end{aligned}$$

This is defined for $x \geq -1$.

For the product,

$$(fg)(x) = \boxed{(1 - x^2)\sqrt{x + 1}},$$

again for $x \geq -1$.

For the first composition,

$$\begin{aligned}(f \circ g)(x) &= f(g(x)) \\ &= 1 - \left(\sqrt{x+1}\right)^2 \\ &= \boxed{-x}.\end{aligned}$$

Although the simplified formula is $-x$, the original composition requires $x \geq -1$ because $g(x)$ must be defined.

For the other composition,

$$\begin{aligned}(g \circ f)(x) &= g(f(x)) \\ &= \sqrt{(1-x^2)+1} \\ &= \boxed{\sqrt{2-x^2}}.\end{aligned}$$

For this to be defined we need

$$2 - x^2 \geq 0,$$

so its domain is $[-\sqrt{2}, \sqrt{2}]$.

(4) The statement

$$\lim_{x \rightarrow 2} f(x) = 5$$

means that the values of $f(x)$ can be made as close to 5 as we want by taking x sufficiently close to 2 (but with $x \neq 2$).

Yes, it is possible that $f(2) = 3$. A limit describes what happens to $f(x)$ *near* $x = 2$; it does not require the value at $x = 2$ itself to equal the limit.

(5) The statement

$$\lim_{x \rightarrow 1^-} f(x) = 3$$

means that $f(x)$ approaches 3 as x approaches 1 using values with $x < 1$. Similarly,

$$\lim_{x \rightarrow 1^+} f(x) = 7$$

means that $f(x)$ approaches 7 as x approaches 1 using values with $x > 1$.


The two-sided limit **does not exist**, because a two-sided limit exists only when the left-hand and right-hand limits agree. Here

$$3 \neq 7.$$

Thus

$$\boxed{\lim_{x \rightarrow 1} f(x) \text{ does not exist.}}$$

(6) For the graph below:



(a) As x approaches 2 from the left, the graph approaches $y = 3$, so

$$\lim_{x \rightarrow 2^-} f(x) = 3.$$

(b) As x approaches 2 from the right, the graph approaches the open point at $(2, 1)$, so

$$\lim_{x \rightarrow 2^+} f(x) = 1.$$

(c) Since the one-sided limits are different,

$$\lim_{x \rightarrow 2} f(x) \text{ does not exist.}$$

(d) The filled point at $x = 2$ is $(2, 3)$, so

$$f(2) = 3.$$

(e) From both sides of $x = 4$, the graph approaches the hole at $(4, 4)$, so

$$\lim_{x \rightarrow 4} f(x) = 4.$$

(f) There is no filled point at $x = 4$, so

$$f(4) \text{ is undefined.}$$

(g) The graph is continuous near $x = 3$ and passes through $(3, 2)$, so

$$\lim_{x \rightarrow 3} f(x) = 2.$$

(7) Consider

$$\frac{x+1}{x-5}.$$

As $x \rightarrow 5$, the numerator approaches 6, which is positive. The sign of the denominator determines the sign of the quotient.

For $x \rightarrow 5^+$, we have $x - 5 \rightarrow 0^+$, so

$$\lim_{x \rightarrow 5^+} \frac{x+1}{x-5} = +\infty.$$

For $x \rightarrow 5^-$, we have $x - 5 \rightarrow 0^-$, so

$$\lim_{x \rightarrow 5^-} \frac{x+1}{x-5} = -\infty.$$

MATH 105 CALCULUS WITH ALGEBRA FALL 2026
HOMEWORK 6

Due **Monday, September 21** at 11:55pm, on Gradescope

- (1) Factor the following expressions:

(a)

$$xy + 4x + 2xw$$

(b)

$$10x^8y^6 + 25x^2y^4 + 20x^3y^{10}$$

- (2) Explain in your own words what is meant by the equation

$$\lim_{x \rightarrow 2} f(x) = 5$$

Is it possible for this statement to be true and yet $f(2) = 3$?

- (3) Explain what it means to say that

$$\lim_{x \rightarrow 1^-} f(x) = 3 \text{ and } \lim_{x \rightarrow 1^+} f(x) = 7$$

In this situation is it possible that $\lim_{x \rightarrow 1} f(x)$ exists?

- (4) For the function f whose graph is given, state the value of each quantity, if it exists. If it does not exist, explain why.

(a) $\lim_{x \rightarrow 2^-} f(x)$


(c) $\lim_{x \rightarrow 2} f(x)$

(e) $\lim_{x \rightarrow 4} f(x)$

(b) $\lim_{x \rightarrow 2^+} f(x)$

(d) $f(2)$

(f) $f(4)$



- (5) For the function g whose graph is given, state the value of each quantity, if it exists. If it does not exist, explain why.

(a) $\lim_{t \rightarrow 0^-} g(t)$

(b) $\lim_{t \rightarrow 0^+} g(t)$

(c) $\lim_{t \rightarrow 0} g(t)$


(d) $\lim_{t \rightarrow 3^-} g(t)$

(e) $\lim_{t \rightarrow 3^+} g(t)$

(f) $\lim_{t \rightarrow 3} g(t)$

(g) $g(3)$

(h) $\lim_{t \rightarrow 2} g(t)$



- (6) Sketch the graph of the function and use it to determine the values of a for which $\lim_{x \rightarrow a} f(x)$ exists.

$$f(x) = \begin{cases} 1 + x & \text{if } x < -1 \\ x^2 & \text{if } -1 \leq x < 1 \\ 2 - x & \text{if } x \geq 1 \end{cases}$$

- (7) Guess the value of the limit (if it exists) by evaluating at the given numbers (using a calculator), correct to six decimal places.

$$\lim_{x \rightarrow 3} \frac{x^2 - 3x}{x^2 - 9},$$

where

$$x = 3.1, 3.05, 3.001, 2.9, 2.95, 2.995.$$

Note: I will not ask you to do this on quizzes/exams since calculators won't be available

MATH 105 CALCULUS WITH ALGEBRA FALL 2026
HOMEWORK 6 SOLUTIONS

- (1) Factor the following expressions.
(a) Every term has a factor of x , so

$$xy + 4x + 2xw = x(y + 4 + 2w).$$

- (b) The greatest common factor is $5x^2y^4$, so

$$10x^8y^6 + 25x^2y^4 + 20x^3y^{10} = 5x^2y^4(2x^6y^2 + 5 + 4xy^6).$$

- (2) Explain in your own words what is meant by the equation

$$\lim_{x \rightarrow 2} f(x) = 5.$$

Solution. This means that when x is close to 2 (but not necessarily equal to 2), the values of $f(x)$ are close to 5. The closer x gets to 2, the closer $f(x)$ gets to 5.

Yes, it is possible to have $\lim_{x \rightarrow 2} f(x) = 5$ while $f(2) = 3$. A limit describes what happens to $f(x)$ *near* $x = 2$; it does not require the value of the function at $x = 2$ itself to equal the limit.


- (3) Explain what it means to say that

$$\lim_{x \rightarrow 1^-} f(x) = 3 \quad \text{and} \quad \lim_{x \rightarrow 1^+} f(x) = 7.$$


Solution. The first statement means that as x approaches 1 through values less than 1, $f(x)$ approaches 3. The second means that as x approaches 1 through values greater than 1, $f(x)$ approaches 7.

Since the left-hand and right-hand limits are different, the two-sided limit $\lim_{x \rightarrow 1} f(x)$ does not exist.

- (4) For the function f whose graph is given, state the value of each quantity, if it exists.



- (a) As $x \rightarrow 2^-$, the graph approaches the filled point at height 3, so $\lim_{x \rightarrow 2^-} f(x) = 3$.
- (b) As $x \rightarrow 2^+$, the graph approaches the open point at height 1, so $\lim_{x \rightarrow 2^+} f(x) = 1$.
- (c) Since the two one-sided limits are different, $\lim_{x \rightarrow 2} f(x)$ does not exist.
- (d) The filled point at $x = 2$ is at height 3, so $f(2) = 3$.
- (e) From both sides of $x = 4$, the graph approaches the open point at height 4, so $\lim_{x \rightarrow 4} f(x) = 4$.
- (f) There is no filled point on the graph at $x = 4$, so $f(4)$ is undefined.
- (5) For the function g whose graph is given, state the value of each quantity, if it exists.




- (a) The graph is continuous near $t = 0$ and is at about height 3, so $\lim_{t \rightarrow 0^-} g(t) \approx 3$.
- (b) Similarly, $\lim_{t \rightarrow 0^+} g(t) \approx 3$.
- (c) Since the one-sided limits agree, $\lim_{t \rightarrow 0} g(t) \approx 3$.
- (d) Approaching $t = 3$ from the left, the graph approaches the open point at height 1, so $\lim_{t \rightarrow 3^-} g(t) = 1$.

- (e) Approaching $t = 3$ from the right, the graph approaches the open point at height 4, so $\lim_{t \rightarrow 3^+} g(t) = 4$.
- (f) Since the two one-sided limits are different, $\lim_{t \rightarrow 3} g(t)$ does not exist.
- (g) The filled point at $t = 3$ is at height 3, so $g(3) = 3$.
- (h) The graph is continuous near $t = 2$ and is at about height 1, so $\lim_{t \rightarrow 2} g(t) \approx 1$.

(6) Sketch the graph of

$$f(x) = \begin{cases} 1 + x & \text{if } x < -1, \\ x^2 & \text{if } -1 \leq x < 1, \\ 2 - x & \text{if } x \geq 1. \end{cases}$$

Solution. For $x < -1$, the graph is the line $y = x + 1$, with an open point at $(-1, 0)$. For $-1 \leq x < 1$, it is the parabola $y = x^2$, with a closed point at $(-1, 1)$ and an open point at $(1, 1)$. For $x \geq 1$, it is the line $y = 2 - x$, beginning with a closed point at $(1, 1)$.



At $x = -1$, the left-hand limit is 0 while the right-hand limit is 1, so the limit does not exist there. At $x = 1$, both sides approach 1, so the limit does exist. Everywhere else, the function is given by a polynomial or a line and has no break.

Therefore $\lim_{x \rightarrow a} f(x)$ exists for every real number a except $a = -1$; that is, for

$$a \in \mathbb{R} \setminus \{-1\}.$$

(7) Guess the value of

$$\lim_{x \rightarrow 3} \frac{x^2 - 3x}{x^2 - 9}$$

by evaluating the expression at the given values of x .

Solution. Using a calculator and rounding to six decimal places gives

x	$\frac{x^2 - 3x}{x^2 - 9}$
3.1	0.508197
3.05	0.504132
3.001	0.500083
2.9	0.491525
2.95	0.495798
2.995	0.499583

The values from both sides of 3 are getting closer to 0.5. Thus the data suggest that

$$\lim_{x \rightarrow 3} \frac{x^2 - 3x}{x^2 - 9} = \frac{1}{2}.$$

MATH 105 CALCULUS WITH ALGEBRA FALL 2026
HOMEWORK 7

Due **Friday, September 25** at 11:55pm, on Gradescope

(1) Determine the following limits.

(a) $\lim_{x \rightarrow 1} \frac{2-x}{(x-1)^2}$

(c) $\lim_{x \rightarrow 5^+} \frac{x^2 - 25}{|x - 5|}$

(b) $\lim_{x \rightarrow -2^+} \frac{x-1}{x^2(x+2)}$

(d) $\lim_{x \rightarrow 1} \frac{1 - \sqrt{x}}{x - 5}$

(2) Suppose that f and g are functions such that

$$\lim_{x \rightarrow 2} f(x) = 4 \quad \lim_{x \rightarrow 2} g(x) = -2 \quad \lim_{x \rightarrow 2} h(x) = 0$$

Which of the below limits exist, and if they exist, what are their values? Explain your work.

(a) $\lim_{x \rightarrow 2} (f(x) + 5g(x))$

(c) $\lim_{x \rightarrow 2} \sqrt{f(x)}$

(b) $\lim_{x \rightarrow 2} (g(x))^3$

(d) $\lim_{x \rightarrow 2} \frac{g(x)}{h(x)}$

(3) The *signum* (or sign) *function*, denoted by sgn , is defined by

$$\text{sgn} = \begin{cases} -1 & \text{if } x < 0 \\ 0 & \text{if } x = 0 \\ 1 & \text{if } x > 0 \end{cases}$$

(a) Sketch the graph of this function.

(b) Find each of the following limits or explain why it does not exist.



(i) $\lim_{x \rightarrow 0^+} \text{sgn } x$

(iii) $\lim_{x \rightarrow 0} \text{sgn } x$

(ii) $\lim_{x \rightarrow 0^-} \text{sgn } x$

(iv) $\lim_{x \rightarrow 0} |\text{sgn } x|$

- (4) The graphs of f and g are given. Use them to evaluate each limit, if it exists. If the limit does not exist, explain why.



(i) $\lim_{x \rightarrow 2} [f(x) + g(x)]$

(ii) $\lim_{x \rightarrow 2} [f(x) - g(x)]$

(iii) $\lim_{x \rightarrow -1} [f(x)g(x)]$

(iv) $\lim_{x \rightarrow 2} [x^2 f(x)]$

MATH 105 Calculus with Algebra – Fall 2026

Homework 7 Solutions

1.

$$\lim_{x \rightarrow 1} \frac{2 - x}{(x - 1)^2}$$

As $x \rightarrow 1$, the numerator approaches 1, while $(x - 1)^2$ approaches 0 through positive values. Therefore the quotient grows without bound:

$$\boxed{+\infty}.$$

2.

$$\lim_{x \rightarrow -2^+} \frac{x - 1}{x^2(x + 2)}$$

As $x \rightarrow -2^+$, the numerator approaches -3 , while x^2 stays positive and $x + 2 \rightarrow 0^+$. Thus the denominator approaches 0 through positive values, so

$$\boxed{-\infty}.$$

3.

$$\lim_{x \rightarrow 5^+} \frac{x^2 - 25}{|x - 5|}$$

Factor the numerator:

$$\frac{x^2 - 25}{|x - 5|} = \frac{(x - 5)(x + 5)}{|x - 5|}.$$

Since $x \rightarrow 5^+$, we have $x - 5 > 0$, so $|x - 5| = x - 5$. Therefore the expression simplifies to $x + 5$, and

$$\boxed{10}.$$

4.

$$\lim_{x \rightarrow 1} \frac{1 - \sqrt{x}}{x - 5}$$

The denominator is nonzero at $x = 1$, so direct substitution works:

$$\frac{1 - \sqrt{1}}{1 - 5} = \frac{0}{-4} = 0.$$

Thus

$$\boxed{0}.$$

5. Suppose that

$$\lim_{x \rightarrow 2} f(x) = 4, \quad \lim_{x \rightarrow 2} g(x) = -2, \quad \lim_{x \rightarrow 2} h(x) = 0.$$

(a)

$$\lim_{x \rightarrow 2} (f(x) + 5g(x))$$

By the limit laws,

$$\lim_{x \rightarrow 2} (f(x) + 5g(x)) = 4 + 5(-2) = -6.$$

Therefore

$$\boxed{-6}.$$

(b)

$$\lim_{x \rightarrow 2} (g(x))^3$$

Since the function $u \mapsto u^3$ is continuous,

$$\lim_{x \rightarrow 2} (g(x))^3 = (-2)^3 = -8.$$

Thus

$$\boxed{-8}.$$

(c)

$$\lim_{x \rightarrow 2} \sqrt{f(x)}$$

Since $f(x) \rightarrow 4$ and the square-root function is continuous at 4,

$$\lim_{x \rightarrow 2} \sqrt{f(x)} = \sqrt{4} = 2.$$

Thus

$$\boxed{2}.$$

(d)

$$\lim_{x \rightarrow 2} \frac{g(x)}{h(x)}$$


The quotient law cannot be applied because $\lim_{x \rightarrow 2} h(x) = 0$. Moreover, the given information does not tell us whether $h(x)$ approaches 0 through positive values, negative values, or with different signs from the two sides. Since the numerator approaches the nonzero number -2 , there is not enough information to determine this limit.

The limit cannot be determined from the given information.

6. The signum function is

$$\operatorname{sgn}(x) = \begin{cases} -1 & \text{if } x < 0, \\ 0 & \text{if } x = 0, \\ 1 & \text{if } x > 0. \end{cases}$$

(a) Its graph is:



(b) (i) As $x \rightarrow 0^+$, x is positive, so $\text{sgn}(x) = 1$. Hence

$$\lim_{x \rightarrow 0^+} \text{sgn}(x) = 1.$$

(ii) As $x \rightarrow 0^-$, x is negative, so $\text{sgn}(x) = -1$. Hence

$$\lim_{x \rightarrow 0^-} \text{sgn}(x) = -1.$$

(iii) The one-sided limits are different, so

$$\lim_{x \rightarrow 0} \text{sgn}(x) \text{ does not exist.}$$

(iv) For every $x \neq 0$, $|\text{sgn}(x)| = 1$. Therefore

$$\lim_{x \rightarrow 0} |\text{sgn}(x)| = 1.$$

7. The graphs of f and g are given below.

From the graph of f , as $x \rightarrow 2$ we have $f(x) \rightarrow -1$. From the graph of g , as $x \rightarrow 2$ we have $g(x) \rightarrow 2$.

(a)

$$\lim_{x \rightarrow 2} [f(x) + g(x)] = -1 + 2 = 1.$$

Thus

$$\boxed{1}.$$

(b)

$$\lim_{x \rightarrow 2} [f(x) - g(x)] = -1 - 2 = -3.$$

Thus

$$\boxed{-3}.$$

(c)

$$\lim_{x \rightarrow -1} [f(x)g(x)].$$

From the graph, $f(x) \rightarrow 1$ as $x \rightarrow -1$. The graph of g approaches a value a little above 2 at $x = -1$ (approximately 2.1 from the drawing). Therefore

$$\lim_{x \rightarrow -1} [f(x)g(x)] \approx 1(2.1) = 2.1.$$

Thus, from the graph,

$$\boxed{\text{approximately } 2.1}.$$

Note: The graph does not mark an exact value of the limit of $g(x)$ at $x = -1$, so only an estimate can be read from the drawing.

(d)


$$\lim_{x \rightarrow 2} [x^2 f(x)] = (2)^2(-1) = -4.$$

Thus


$$\boxed{-4}.$$

MATH 105 CALCULUS WITH ALGEBRA FALL 2026
HOMEWORK 8
DUE MONDAY, SEPTEMBER 28 AT 11:55PM, ON GRADESCOPE

- (1) (a) From the graph of f , state the numbers at which f is discontinuous and explain why.
 (b) For each of the numbers stated in part (a), determine whether f is continuous from the right, or from the left, or neither.



- (2) From the graph of g , state the intervals on which g is continuous.



- (3) Explain why the following function is discontinuous at the given number a . Sketch the graph of the function.

$$f(x) = \frac{1}{x+2}, \quad a = -2.$$

- (4) Explain why the following function is discontinuous at the given number a . Sketch the graph of the function.

$$f(x) = \begin{cases} 1 - x^2, & x < 1, \\ \frac{1}{x}, & x \geq 1, \end{cases} \quad a = 1.$$

- (5) Is the following function continuous at 0? What about at -1? Sketch its graph.

$$f(x) = \begin{cases} \frac{1}{x+2}, & x < -1, \\ 1, & x \geq -1, \end{cases}$$

- (6) Draw the graph of a function f whose domain is all real numbers and is continuous except for the following discontinuities:

- (a) f has a removable discontinuity at -2 and an infinite discontinuity at 2
- (b) f has a jump discontinuity at -3 and removable discontinuity at 2
- (c) f is discontinuous at 0 and 3 , but continuous from the right at 0 and from the left at 3

MATH 105 Calculus with Algebra — Fall 2026

Homework 8 Solutions

1. (a) From the graph, f is discontinuous at

$$\boxed{x = -4, -2, 2, 4}.$$

At $x = -4$ there is a hole, so this is a removable discontinuity.

At $x = -2$, the left-hand and right-hand limits are different, so there is a jump discontinuity.

At $x = 2$, the left-hand limit is the height of the open circle, while the right-hand limit is the height of the filled point. These are different, so there is a jump discontinuity.

At $x = 4$, the graph becomes unbounded as $x \rightarrow 4^-$, while the right-hand branch begins at a finite filled point. Thus $x = 4$ is an infinite discontinuity.

- (b) At $x = -4$, $f(-4)$ is not defined, so f is continuous from neither side.

At $x = -2$, the filled point lies on the branch coming from the left, so

$$\boxed{f \text{ is continuous from the left at } -2.}$$

At $x = 2$, the filled point lies on the branch coming from the right, so

$$\boxed{f \text{ is continuous from the right at } 2.}$$

At $x = 4$, the branch beginning at the filled point is continuous to the right, so

$$\boxed{f \text{ is continuous from the right at } 4.}$$

2. From the graph of g , the breaks occur at $x = -2, -1, 0$, and 1 .

The continuous pieces are

$$\boxed{[-3, -2), \quad (-2, -1), \quad (-1, 0], \quad (0, 1), \quad (1, 3].}$$

The endpoints -3 , 0 , and 3 can be included on the indicated intervals because continuity at an endpoint of an interval only requires the appropriate one-sided limit.

- 3.

$$f(x) = \frac{1}{x+2}, \quad a = -2.$$


The function is not defined at $x = -2$. In fact, $x = -2$ is a vertical asymptote:

$$\lim_{x \rightarrow -2^-} \frac{1}{x+2} = -\infty, \quad \lim_{x \rightarrow -2^+} \frac{1}{x+2} = +\infty.$$

Therefore

$$\boxed{f \text{ is discontinuous at } x = -2.}$$

A sketch is:



4.

$$f(x) = \begin{cases} 1 - x^2, & x < 1, \\ \frac{1}{x}, & x \geq 1. \end{cases}$$

At $x = 1$,

$$\lim_{x \rightarrow 1^-} f(x) = 1 - 1^2 = 0, \quad \lim_{x \rightarrow 1^+} f(x) = 1, \quad f(1) = 1.$$


The one-sided limits are different, so

$$\boxed{f \text{ is discontinuous at } x = 1.}$$

It is continuous from the right at $x = 1$, since

$$\lim_{x \rightarrow 1^+} f(x) = f(1) = 1.$$

A sketch is:



5.

$$f(x) = \begin{cases} \frac{1}{x+2}, & x < -1, \\ 1, & x \geq -1. \end{cases}$$

At $x = 0$, the function is simply $f(x) = 1$ in a neighborhood of 0, so

f is continuous at 0.


At $x = -1$,

$$\lim_{x \rightarrow -1^-} f(x) = \frac{1}{-1 + 2} = 1, \quad \lim_{x \rightarrow -1^+} f(x) = 1, \quad f(-1) = 1.$$

Therefore

f is also continuous at -1 .

Notice that the graph does have a vertical asymptote at $x = -2$.




6. Many answers are possible. Here are three examples.

- (a) We want a removable discontinuity at -2 and an infinite discontinuity at 2 , while keeping the domain equal to all real numbers. One example is

$$f(x) = \begin{cases} \frac{1}{x-2}, & x \neq -2, 2, \\ 0, & x = -2, \\ 0, & x = 2. \end{cases}$$


At $x = -2$, the limit exists and equals $-1/4$, but $f(-2) = 0$, so the discontinuity is removable. At $x = 2$, the graph has a vertical asymptote, even though we have assigned a value to $f(2)$.



- (b) We want a jump at -3 and a removable discontinuity at 2 . For example,

$$f(x) = \begin{cases} 0, & x < -3, \\ 1, & x \geq -3, \ x \neq 2, \\ 0, & x = 2. \end{cases}$$

At $x = -3$, the left-hand limit is 0 and the right-hand limit is 1 , so there is a jump. At $x = 2$, the limit is 1 but the function value is 0 , so the discontinuity is removable.




- (c) We want discontinuities at 0 and 3 , with right continuity at 0 and left continuity at 3 . One example is

$$f(x) = \begin{cases} 0, & x < 0, \\ 1, & 0 \leq x \leq 3, \\ 2, & x > 3. \end{cases}$$

At $x = 0$, $f(0) = 1$ and the right-hand limit is 1 , so the function is continuous from the right there, but the left-hand limit is 0 .

At $x = 3$, $f(3) = 1$ and the left-hand limit is 1 , so the function is continuous from the left there, but the right-hand limit is 2 .



Math 105 ~ Calculus with Algebra: Review Packet

- These are practice problems, and are not to be turned in.
- On Wednesday in class, I will work any problems you ask about, and answer any questions.
- On Thursday, there will be a practice exam during the worksheet session.
- *Good luck!*

What you need to know for Exam 1

You should know everything from the beginning of the course up to (and including). The following is a list of most of the topics covered. **THIS IS NOT A COMPREHENSIVE LIST, BUT MERELY AN AID.** Remember, no calculators or electronics that can connect to the internet on any exams.

- Appendices A and B: Review equations of lines, and inequalities. Parallel lines have same slope. Perpendicular lines have opposite reciprocal slopes. Know the difference between *point-slope form*, and *slope-intercept form*. Know how to use *slope-intercept form* to sketch lines quickly and efficiently.
- 1.1: What is a function? (Vertical line test.) Domain and Range of a function. Know how to quickly sketch all the standard functions like x^2 , x^3 , $\frac{1}{x}$, $\sqrt{x}$, and $|x|$. Piecewise-defined functions, including using all these functions.
- 1.2: Standard functions: polynomials (including constant and linear), power functions (including root functions and things like $\frac{1}{x}$), rational functions.
- 1.3: Adding, subtracting, multiplying, and dividing functions. Composition of functions. Given the graph of $y = f(x)$, what do $y = f(x) \pm c$, $y = f(x \pm c)$, $y = cf(x)$, $y = f(cx)$, $y = -f(x)$, $y = f(-x)$, and $y = |f(x)|$ look like? (See boxes on pp. 36–37, and Example 5.)
- 1.5: Limits. Intuitive idea of what a limit is. Estimating limits using tables and computations. Be familiar with the various ways a limit can fail to exist. Right- and left-hand limits; and Box 3, page 55. Infinite limits. Vertical asymptotes.
- 1.6: Limit laws: sum, difference, constant multiple, product, quotient, power, root laws. Computing limits of polynomials, rational functions, and other functions. Computing harder limits, like $0/0$, using algebraic tricks. What to do in each case of a DSP (Direct substitution) attempt:

When you get

$$\frac{\text{constant, including } 0}{\text{non-zero constant}}$$

then it's just DSP.

When you get

$$\frac{\text{non-zero constant}}{0}$$

then you do *sign analysis*. Study the RHL and LHL (Right and Left Hand Limit) cases carefully. Yes it will be an infinite limit, but what sign is attached? Pay attention to your two-point argument: 1. size and 2. sign.

When you get

$$\frac{0}{0}$$

then you use an algebraic technique to cancel the zero factor. Use factoring, common denominator, or conjugate techniques. It means **MORE WORK!**

Finally, if you see absolute values, certainly for a $\frac{0}{0}$ case, then you must explore the two RHL and LHL cases.

- 1.8: Continuity: at a number a , or on an interval I . Know both the official definition

$$\lim_{x \rightarrow a} f(x) = f(a)$$

and the intuitive idea (you can draw it without lifting your pencil off the paper). The three ways continuity can fail ($f(a)$ not defined; or $\lim_{x \rightarrow a} f(x)$ not defined; or both defined, but not equal). Continuity from the right or left.

Tips

- You will be allowed to do the problems in any order, but **make sure you don't accidentally skip any** if you like to jump around.
- Make sure you know how to manipulate functions. If any of the problems from Sections 1.1–1.3 gave you serious trouble, make sure you get things cleared up.
- You will need to compute some limits carefully with full justification. Be ready. Know the algebra tricks you need for $\frac{0}{0}$ limits. Know when to use LHL and RHL sign analysis. Also know how to examine the two cases for absolute value pieces of your functions.
- Make sure you are clear when making a statement about discontinuity. You must tell me which of the three pieces of the definition fail(s).
- Do not write $\lim_{x \rightarrow a}$ by itself without the function.
- If you make a declaration about a RHL or LHL limit, then write it out formally using math notation. Look to put forward your *best presentation* for each solution.
- For your piece-wise defined graphs, make sure that when you piece together the puzzle of individual function pieces, that you pay attention to the end-point output values where those cuts are. That will help you determine whether the function pieces line up. Indeed that might change a check of continuity at a given point.
- Justify all of your work!! Don't just use random words to explain answers; I will look for clear mathematical statements.

Limit Practice Problems

Evaluate the following limits. Be clear if the limit equals a finite value, Does Not Exist, or is $+\infty$ or $-\infty$. Always justify your work:

1. $\lim_{w \rightarrow 0} \frac{16}{w} =$

2. $\lim_{t \rightarrow 2} \frac{3-t}{t-2} =$

3. $\lim_{t \rightarrow 2} \frac{3-t}{(t-2)^2} =$

4. $\lim_{x \rightarrow 4} \frac{(x+2)^2}{x^2-3x-4} =$

5. $\lim_{x \rightarrow 4} \frac{x-4}{x^2-3x-4} =$

6. $\lim_{x \rightarrow 4} \frac{x^2-2x-8}{x^2-3x-4} =$

7. $\lim_{x \rightarrow 1} \frac{x^2-4x-12}{x^2-3x-18} =$

8. $\lim_{x \rightarrow 0} \frac{x^2-4x-12}{x^2-3x-18} =$

9. $\lim_{x \rightarrow -3} \frac{x+2}{x+3} =$

10. $\lim_{x \rightarrow -2} \frac{x^2-4x-12}{x^2-3x-18} =$

11. $\lim_{x \rightarrow 0} \frac{x^2-4x}{x^2-7x} =$

12. $\lim_{x \rightarrow 3} \frac{x^2-9}{|x-3|} =$

13. $\lim_{x \rightarrow -5} \frac{x^2+6x+5}{|x+5|} =$

14. $\lim_{t \rightarrow 1} \frac{t^2-1}{t^2-11t+10} =$

15. $\lim_{t \rightarrow 1} \frac{t^2}{t^2+t-1} =$

$$16. \lim_{t \rightarrow -1} \frac{2009(t^2 + 6t + 5)}{t^2 + t} =$$

$$17. \lim_{x \rightarrow 9} \frac{x^2 - 10x + 9}{x^2 + x - 90} =$$

$$18. \lim_{t \rightarrow 1} (t^{500} + t^{400} + t^{300} + t^{200} + t^{100}) =$$

$$19. \lim_{x \rightarrow 3} \frac{x^2 - x - 6}{x^2 - 2x - 3} =$$

$$20. \lim_{x \rightarrow 1} \frac{x - 1}{\sqrt{x + 3} - 2} =$$

$$21. \lim_{x \rightarrow 9} \frac{9x - x^2}{3 - \sqrt{x}} =$$

$$22. \lim_{x \rightarrow 1} \frac{5}{1 - x} =$$

$$23. \lim_{x \rightarrow 5} \frac{6x}{5 - x} =$$

$$24. \lim_{x \rightarrow 2} \frac{x^2 - 9x + 14}{x^2 - 4x + 4} =$$

$$25. \lim_{x \rightarrow 2} \frac{x^2 - 4}{|x - 2|} =$$

$$26. \lim_{x \rightarrow 3} \frac{\sqrt{x + 6} - 3}{x^2 - x - 6} =$$

$$27. \lim_{x \rightarrow 7} \frac{\frac{1}{7} - \frac{1}{x}}{x - 7} =$$

$$28. \lim_{x \rightarrow -6} \frac{\frac{1}{2-x} - \frac{1}{8}}{x + 6} =$$

$$29. \lim_{x \rightarrow 3} \frac{\sqrt{x + 1} - 2}{3 - x} =$$

$$30. \lim_{x \rightarrow 7} \frac{x^2 - 49}{2 - \sqrt{x - 3}} =$$

$$31. \lim_{x \rightarrow 5} \frac{\frac{1}{\sqrt{x+20}} - \frac{1}{5}}{x - 5} = \quad \text{Challenge!}$$

Functions and Limit Practice Problems Evaluate the following limits:

32. Let $g(x) = 2x + 1$. Compute $\lim_{x \rightarrow 1} \frac{x - 1}{g(x^2) - 3} =$

33. Let $G(u) = u^2 + u$. Compute $\lim_{u \rightarrow 2} \frac{u^2 - 2u}{G(u - 3)} =$

34. Let $h(y) = y^2 - 3$. Compute $\lim_{x \rightarrow -2} \frac{x + 2}{h(2x) - h(x + 6)} =$

35. Let $f(t) = \frac{1}{t}$. Compute $\lim_{t \rightarrow 4} \frac{f(t - 3) - 4f(t)}{t - 4} =$

More Functions

36. Let $f(x) = \sqrt{x}$, $g(x) = x^2 + 4$, and $h(x) = \frac{1}{x}$. Compute (and simplify, if possible) the following:

(a) $f \circ g(x) =$

(b) $g \circ f(x) =$

(c) $h \circ g \circ f(x) =$

(d) $g \circ g(x) =$

Piece-wise defined functions

Consider each of the following piecewise defined functions. Answer the related questions. *Justify your answers* please.

37. Let

$$f(x) = \begin{cases} x, & x < 0, \\ x^2, & 0 \leq x \leq 2, \\ 8 - x, & x > 2. \end{cases}$$

Sketch the graph. Find the numbers at which f is discontinuous. Evaluate:

$$\lim_{x \rightarrow 2} f(x) = \quad \lim_{x \rightarrow 0} f(x) = \quad f(2) = \quad f(4) =$$

38. Let

$$f(x) = \begin{cases} x - 1, & x < 2, \\ 1, & 2 < x < 4, \\ 3, & x = 4, \\ \sqrt{x}, & x > 4. \end{cases}$$

Sketch the graph. Find the numbers at which f is discontinuous. Evaluate:

$$\lim_{x \rightarrow 0} f(x) = \quad \lim_{x \rightarrow 2} f(x) = \quad \lim_{x \rightarrow 4} f(x) = \quad f(4) =$$

39. Let

$$h(x) = \begin{cases} \frac{8}{x+2}, & x < 0, \\ 2, & x = 0, \\ \frac{1}{2}x - 4, & 0 < x \leq 16, \\ \sqrt{x}, & x > 16. \end{cases}$$

Sketch the graph. Find the numbers at which h is discontinuous. Evaluate:

$$\lim_{x \rightarrow -2} h(x) = \quad \lim_{x \rightarrow 0} h(x) = \quad \lim_{x \rightarrow 16} h(x) = \quad f(0) = \quad f(16) =$$

40. Let

$$F(x) = \begin{cases} \frac{1}{x-4}, & x > 4, \\ 3-x, & 0 < x < 4, \\ 3-x^2, & x < 0. \end{cases}$$

Sketch the graph. Find the numbers at which F is discontinuous. Evaluate:

$$\lim_{x \rightarrow 0} F(x) = \quad \lim_{x \rightarrow 4} F(x) = \quad F(-3) = \quad F(4) =$$


41. Consider the line L given by $2x + 4y = 12$.

- (a) Sketch this line L .
- (b) Find the equation of the new line M that is **parallel** to the first line L , and passes through the point $(0, 1)$.
- (c) Find the equation of the new line M' that is **perpendicular** to the first line L , and passes through the point $(1, -2)$.
- (d) Sketch this new line M found in (b).

42. Sketch the graphs of the following functions. Label a few points on the graph.

- (a) $f(x) = \sqrt{x-3}$
- (b) $f(x) = 1 + \frac{1}{x+1}$
- (c) $f(x) = x|x|$
- (d) $f(x) = \frac{x^2-9}{|x-3|}$

43. Let $f(x)$ be the function with the following graph.



- (a) Graph $-f(x)$.
- (b) Graph $1 + f(2x + 2)$.
- (c) Write a (piecewise) formula for $f(x)$.

44. Let $f(x) = x^2 + 3x$.

- (a) Compute $f(x + h)$
- (b) Simplify

$$\frac{f(x + h) - f(x)}{h}$$

45. Simplify the following expressions:

- (a)

$$\frac{1}{x-1} - \frac{1}{x}$$

- (b)

$$\frac{x}{x-1} - \frac{x}{x+1}$$

- (c)

$$\frac{\frac{s+1}{s-1} + \frac{s-1}{s+1}}{\frac{1}{s^2-1}}$$