Generative models and how it can be used in bioinformatics

2022-05-03 Ping-Han Hsieh

Let us assume we all live in a virtual reality.


- Let us assume we all live in a virtual reality.
- Where I am the creator of this world.
 - I get to decide the physical law
 - I get to decide what people can observe.

- Let us assume we all live in a virtual reality.
- Where I am the creator of this world.
 - I get to decide the physical law
 - I get to decide what people can observe.
- I am trying to make sure nobody realizes they live in the virtual reality.
 - The physical law need to be understandable, so people find this world interesting.
 - But not too simple, so no one realizes they live in the virtual reality.


- Let us assume we all live in a virtual reality.
- Where I am the creator of this world.
 - I get to decide the physical law
 - I get to decide what people can observe.
- I am trying to make sure nobody realizes they live in the virtual reality.
 - The physical law need to be understandable, so people find this world interesting.
 - But not too simple, so no one realizes they live in the virtual reality.
- So, I can keep my reign forever.

- Let me put the hint to infer the physical law in the virtual universe.
- And the law is very simple:
 - Depend on the expression of Transcription Factor A, the expression of Gene A will change in a linear way.
 - There are only 16 cells in this universe, each has only Transcription Factor A, Transcription Factor B, Transcription Factor C and Gene A and you can observe the exact expression of each molecule.
 - And I let you know transcription factors regulate genes.
 - Gene expression follows Gaussian distribution.


- Let me put the hint to infer the physical law in the virtual universe.
- And the law is very simple:
 - Depend on the expression of Transcription Factor A, the expression of Gene A will change in a linear way.
 - There are only 16 cells in this universe, each has only Transcription Factor A, Transcription Factor B, Transcription Factor C and Gene A and you can observe the exact expression of each molecule.
 - And I let you know transcription factors regulate genes.
 - Gene expression follows Gaussian distribution.
- Hope no one will find out the secret of this universe.




Term Definition




Linear Generative Process




$$x = f_{decoder}(z)$$
$$x = zw + b$$
$$\theta = \{w, b\}$$



From the Data Analysist Point of View





- Gather information about the molecular systems (features)
- Build machine learning model or perform correlation analysis.
- Found that TF A can be used to predict the expression of Gene A (or the highest correlation coefficient).
- Conclude that in this molecular system, TF A might relate to the production of Gene A.


• Soon people will find out they are controlled by me. Let me make the world more complicated.

 Soon people will find out they are controlled by me. Let me make the world more complicated.


- The new law:
 - Depend on the expression of Transcription Factor A, the expression of Gene A will change in a non-linear way (but in groups).
 - There are only 16 cells in this universe, each has only Transcription Factor Aand Gene A and you can observe the exact expression of each molecule.
 - And I let you know transcription factors regulate genes.
 - Gene expression follows Gaussian distribution.

 Soon people will find out they are controlled by me. Let me make the world more complicated.

- The new law:
 - Depend on the expression of Transcription Factor A, the expression of Gene A will change in a non-linear way (but in groups).
 - There are only 16 cells in this universe, each has only Transcription Factor A and Gene A and you can observe the exact expression of each molecule.
 - And I let you know transcription factors regulate genes.
 - Gene expression follows Gaussian distribution.
- I guess no one can figure this out.




Non-linear Generative Process



$$x = f_{decoder}(z)$$

$$x = 1_{\{z < t_1\}}[zw_1 + b_1] + 1_{\{t_1 \le z \le t_2\}}[zw_2 + b_2] + 1_{\{z > t_2\}}[zw_3 + b_3]$$

$$\theta = \{w_1, w_2, w_3, b_1, b_2, b_3\}$$



Maybe I should just remove all the researchers in this universe?

Maybe I should just remove all the researchers in this universe?

• This time:


- Depend on the expression of Molecule A, the expression of Gene A will change in a non-linear way (but in groups).
- There are only 16 cells in this universe, each has only Molecule A and Gene A and you can only observe the exact expression of Gene A in part of these cells.
- You do not know which molecule regulates Gene A, you can not observe expression of Molecule A neither.
- Gene expression follows Gaussian distribution.

Maybe I should just remove all the researchers in this universe?


• This time:

- Depend on the expression of Molecule A, the expression of Gene A will change in a non-linear way (but in groups).
- There are only 16 cells in this universe, each has only Molecule A and Gene A and you can only observe the exact expression of Gene A in part of these cells.
- You do not know which molecule regulates Gene A, you can not observe expression of Molecule A neither.
- Gene expression follows Gaussian distribution.


Perfect!




Gaussian Mixture Model (1)




Gaussian Mixture Model (2)



Gaussian Mixture Model (3)





Gaussian Mixture Model (4)

• Assumptions:

- There are different clusters in the samples.
- Samples from the same cluster undergo similar generative process.


Limitations:

- Gaussian mixture model can be used to infer the generative process for a group of samples, but not for each individual sample.
- Additional analysis for each clusters need to be done to reconstruct the generative process.
- Not effective on high-dimensional data (curse of dimensionality).
- Latent variable is a discrete variable.
- Data distribution needs to be a mixture of Gaussian distribution.


• Finally:


- Depend on the expression of Molecule A, the expression of Gene A will change in a non-linear way (even the variability within each group is generated using a specific rule).
- There are only 16 cells in this universe, each has only Molecule A and Gene A and you can only observe the expression of Gene A in part of these cells.
- You do not know which molecule regulates Gene A, you can not observe expression of Molecule A neither.
- The observed expression of Gene A is uncertain.
- Gene expression follows Negative Binomial distribution.

Improve Gaussian Mixture Model (1)





Improve Gaussian Mixture Model (2)







Improve Gaussian Mixture Model (3)






Improve Gaussian Mixture Model (4)





Improve Gaussian Mixture Model (5)



Improve Gaussian Mixture Model (5)

- The expectation-maximization becomes unstable when the number of component increases.
- How to address this issue?
 - Assume K = N.
 - For each point we observe, train a function and map them into a Gaussian distribution.
 - Since it is a function, now we can map any input (even if they are not in this universe) to a Gaussian distribution.


$$\mu = \frac{1}{N}(x_1 + x_2 + \dots + x_N)$$

$$\sigma = \sqrt{\frac{1}{N}[(x_1 - \mu)^2 + (x_2 - \mu)^2 + \dots + (x_N - \mu)^2]}$$

$$\mu = f_{encoder,\mu}(x)$$

$$\sigma = f_{encoder,\sigma}(x)$$

Improve Gaussian Mixture Model (6)



Improve Gaussian Mixture Model (7)


• Assumptions:


- There are different clusters in the samples.
- Samples from the same cluster undergo similar generative process.

Limitations:

- Gaussian mixture model can be used to infer the generative process for a group of samples, but not for each individual sample.
- Additional analysis for each clusters need to be done to reconstruct the generative process.
- Not effective on high-dimensional data (curse of dimensionality).
- Latent variable is a discrete variable.
- Data distribution needs to be a mixture of Gaussian distribution.

Improve Gaussian Mixture Model (6)





Improve Gaussian Mixture Model (7)


• Assumptions:

- There are different clusters in the samples.
- Samples from the same cluster undergo similar generative process.


Limitations:

- Gaussian mixture model can be used to infer the generative process for a group of samples, but not for each individual sample.
- Additional analysis for each clusters need to be done to reconstruct the generative process.
- Not effective on high-dimensional data (curse of dimensionality).
- Latent variable is a discrete variable.
- Data distribution needs to be a mixture of Gaussian distribution.

Variational Autoencoder (1)



Variational Autoencoder (2)



Conclusion

Generative Models

- Allow us to reconstruct the generative process of the system we observe.
- Allow us to identify the potential causal factors that generative the data we observe.
- Allow us to understand the cluster of samples
- Allow us to generative new data based on the system we observe.

Variational Autoencoder

• A flexible framework that allow us to model the generative process of the system we observe if the data distribution is continuous (and differentiable).

Thanks