

Adult Cardiac Arrest – PEA/Asystole

Key Findings: No shock advised. Not a shockable rhythm. Consider possible causes such as hyperkalemia, hypovolemia or toxins.

EMT/AEMT

- Follow Adult Cardiac Arrest – Initial Care Guideline

Paramedic

- Continuous High Quality CPR
 - Follow Adult Cardiac Arrest – Initial Care Guideline
 - **Epinephrine** 0.5 mg IO/IV
 - Start Fluid Bolus **NS/LR**
 - Endotracheal Intubation (2 initial attempts) if adequate ventilation is not achieved by SGA
 - **Epinephrine** 0.5 mg IO/IV
 - **Bicarbonate** 1 mEq/kg, IO/IV (If suspected hyperkalemia, ingestion, toxins or prolonged down-time)
 - Consider Cardiac Arrest Additional Treatments
 - **Epinephrine** 0.5 mg IO/IV
 - **Epinephrine** 0.5 mg IO/IV
-
- Epinephrine (1:10,000) 0.5 mg IO/IV max 4 doses (2 mg total)
 - Dosing intervals every 5 to 10 minutes
 - DO NOT use Calcium Chloride unless hyperkalemia is suspected

Hyperkalemia should be expected in:

- Rhabdomyolysis
- Severe dehydration
- Avalanche
- Known renal disease