

# Adult Cardiac Arrest – VF / Pulseless VT

## EMT/AEMT

- Follow Adult Cardiac Arrest – Initial Care Guideline

## Paramedic

### For Patient in VF or Pulseless VTach (Shockable Rhythm)

- Continuous high-quality CPR
- **Defibrillation** 200 J
- Follow Adult Cardiac Arrest – Initial Care Guideline
- Supraglottic airway (SGA)
- **Epinephrine** 0.5 mg IO/IV
- Start Fluid Bolus **NS** IO/IV
- **Defibrillation** 200 J
- **Amiodarone** 300mg IO/IV
- **Change Vector of Defibrillation Pads or Double Sequential Defibrillation** (Max energy both devices) - Double Sequential Defibrillation requires 2 AEDs, a monitor and an AED, or 2 monitors and therefore may not be possible in all environments
- **Epinephrine** 0.5 mg IO/IV
- **Amiodarone** 150 mg IO/IV
- **Change Vector of Defibrillation Pads or Double Sequential Defibrillation** (Max energy both devices)
- **Epinephrine** 0.5 mg IO/IV
- Continue ACLS with epinephrine doses 0.5mg until Epinephrine 2mg total (4 doses of 0.5mg) is achieved unless ROSC is achieved prior to 2mg given.
- Continue Defibrillation until ROSC is achieved, resources are exhausted, the scene becomes unsafe, or advanced care providers arrive on location