

Aerie Backcountry Medicine EMT/AEMT Expanded Scope

These guidelines are intended to expand the general scope of EMTs and AEMTs working with Timberline Medics. We recognize that some of these expanded scopes are not allowable in particular states. When operating in more restricted states without an Emergency Declaration Order, we will notify the EMT or AEMT that the expanded scope protocols are not in use.

In the Guidelines the expanded scope is denoted by a Green Box between the EMT/AEMT blue box and the Paramedic red box. If doses are missing, please see the Paramedic red box for guidance.

Below is a summary of the expanded scope for EMT and AEMT providers:

1. Supraglottic airway – iGel – in individual unable to protect their airway
2. IO placement and administration of medications
 - a. D10W
 - b. LR
 - c. NS
 - d. TXA
3. Nebulizer administration of Albuterol, Ipratropium bromide, Epinephrine, and TXA
4. IM Injections of Epinephrine 1:1000, Diphenhydramine, Toradol, and Ondansetron
5. Eye emergency (without concern for open globe)
 - a. Tetracaine – 2 drops in affected eye prior to irrigation
 - b. Irrigation with Morgan Lens
6. Blood Glucose Testing
7. Treatment of hypoglycemia
 - a. D10W via IO in patient who cannot protect their airway
8. Pain management – Can give Acetaminophen and either Ibuprofen or Ketorolac together (do not give ibuprofen and Ketorolac together)
 - a. Acetaminophen PO
 - b. Ibuprofen PO (400mg) – Take with 1L water
 - c. Toradol 15mg IM – Take with 1L water
 - d. Lidocaine patch
9. Famotidine 20mg PO for GERD and allergy symptoms
10. Loratadine 10mg PO for allergy symptoms
11. Guafenesin in pre-dosed packets for benign cough
12. Hydrocortisone cream for skin rash
13. Naloxone 0.5mg-2mg IN or IM for respiratory depression with concern of overdose on opiates

14. Steroids

- a. Prednisone 60mg PO
- b. Solumedrol 125mg IM
- c. Dexamethasone 10mg PO/IM (can give IV solution PO)

15. Ondansetron (Zofran) 4mg ODT or IM every 5 min x 2 doses (8mg total)

16. Sennosides (Senokot) 17.2mg daily for constipation

17. TXA

- a. TXA 2g IO for significant trauma and tolerating IO
- b. TXA 500mg Nebulized or IN with atomizer for epistaxis or airway bleeding