

General Medical Care

General History to Obtain in All Adult Medical Patients:

- **Age of the patient**
- **Chief complaint**
- **Timing** of events / factors leading up to medical personnel's contact with the patient
- **Other symptoms or complaints** (review of systems)
- **Patient's Past Medical History**
- **Patient Medications and Allergies**
- **USE BYSTANDERS** to gain more information if possible
- **Has the patient contacted or been treated by medical in this deployment? Other deployments?**
- **Specific to environment: Has the patient had difficulty with heat, cold, or altitude in the past?**

Remember to Assess the Following in all Medical Patients:

- **General Appearance** – If the patient looks ill, they probably are!
- **Vital signs** – Obtain all necessary vital signs on every patient encounter.
- **Mental Status** – Alert and oriented x person, place, time, event (spell it out). If a trauma, you may utilize the Glasgow Coma Score as well.
- **Breath sounds** – Clear? Rales? Wheezing? Diminished?
- **Heart sounds**
- Abdominal exam
- Skin – Color? Cool or warm? Diaphoretic? Clammy?
- Neurologic exam – Is the speech normal? Are grip strengths and facial movements symmetrical? Is the patient moving all 4 extremities? Is the patient noted to have numbness in his/her trunk or extremities?

EMT/AEMT

General Care

- Scene safety
- Primary assessment / Vital signs (including blood glucose / Temperature)
- Primary respiratory assessment – include breath sounds, pulse oximetry, EtCO₂, Blood glucose if indicated
- Supplemental Oxygen if available and SpO₂ <94%
- Airway management as indicated –
 - Open airway, clear airway, suction
 - Basic ventilatory (BVM) airway and basic adjunct (OPA vs NPA)
- Secondary assessment and treat any concerning findings as indicated
- Look for evidence of trauma
- Spinal motion restriction if indicated

EMT/AEMT - Expanded Scope

- Supraglottic airway if needed for ventilation
- **NS** IO 500ml x 1 bolus per Medical Shock Guidelines
- **Zofran** 4mg IM/ODT every 5 min x 2 doses (8mg max) for nausea/vomiting
- Pain management as per Pain Management Guideline
- Hypoglycemia per Diabetic Emergency Guideline

Paramedic

- 12-Lead EKG if accessible
- EtCO₂, as indicated and if accessible
- **NS** IV/IO 500-1000 fluid bolus per Medical Shock Guideline, if in non-hemorrhagic shock
- Consider **Zofran** 4mg IV/IO/IM or ODT for nausea / vomiting
- Pain management as per Pain Management Guideline
- Secondary survey and treat any concerning findings as indicated

