

Ketorolac

(Toradol)

I. Classification – Non-steroidal anti-inflammatory, Antipyretic

II. Action(s)

- COX-1 and COX-2 inhibitor

III. Indication(s)

- Mild – Moderate pain
- Fever

IV. Relative Contraindication

- Known Coronary Artery Disease
- Known Diabetes Mellitus

V. Contraindications

- Known Hypersensitivity
- Bleeding disorders
- Anti-coagulant use
- Anti-platelet use
- Renal failure
- Rhabdomyolysis
- Severe dehydration
- Max dose < 4hrs prior (see notes)

VI. Drug interactions

- Antiplatelets: may enhance the antiplatelet effect
- Alcohol (Ethyl): May enhance the adverse/toxic effects of NSAIDs – specifically the risk of GI bleeding
- Anticoagulants: May enhance the anticoagulant effect of anticoagulants
- Aspirin: May enhance the adverse/toxic effect of Aspirin
- Beta-blockers: May diminish the antihypertensive effect of Beta-blockers
- Corticosteroids: May enhance the adverse/toxic effect of NSAIDs
- Metformin: May enhance the adverse/toxic effect of Metformin

VII. Adult Dosing and Administration

- 15 mg every 6-8 hours IV/IO/IM

VIII. Pediatric Dosing and Administration

- 0.5 mg/kg IV/IO/IM every 6-8 hours
- Max dose 15mg

IX. Onset

- 30 min

X. Duration

- 4-6 hrs

XI. Possible Adverse Effects

- Acute Myocardial Infarction
- Cerebrovascular event
- Hypertension

- GI bleed/ulcer/perforation
- Hypersensitivity reaction
- Acute kidney injury
- Aplastic anemia, other blood dyscrasias

XII. Other Notes

- Ketorolac and acetaminophen given together have been shown to have the same effect on pain reduction as opioids