

Metoprolol

(Lopressor, Toprol)

I. Classification

- Antihypertensive, Beta-blocker

II. Action(s)

- Preferentially blocks beta₁ receptors

III. Indication(s)

- Refractory VF / pulseless VT
- Narrow Complex Tachycardia with pulses
- Hypertension

IV. Contraindication(s)

- Allergy
- Bradycardia, 2° or 3° heart block, cardiogenic shock, cardiac failure
- Sick-sinus syndrome
- Severe peripheral arterial disease

V. Precautions

- Use very cautiously in patients with congestive heart failure
- Use cautiously in patients with history of asthma / COPD

VI. Drug Interactions

- Digitalis and other beta blockers – may increase risk of bradycardia

VII. Adult Dosing and Administration

- Narrow Complex Tachycardia: 5 mg slow IV / IO push, may repeat x1 after 5 minutes
- SBP >220 or DBP >120 WITH signs of end-organ damage:
 - ☐ 5 mg IV / IO every 2 minutes x 3 doses (max total dose 15 mg)
 - ☐ Consult On-Call Medical Director for Order

VIII. Pediatric Dosing and Administration

- Not indicated for pediatric patients

IX. Onset

- 10-20 minutes

X. Duration

- 5-8 hours

XI. Possible Adverse Effects

- Allergy
- Bradycardia and hypotension
- Respiratory difficulty, wheezing, bronchospasm
- Congestive Heart Failure