

Ondansetron

(Zofran, Zofran ODT)

- I. **Classification** – Antiemetic
- II. **Action(s)**
 - Antagonizes 5-HT₃ serotonin receptors
- III. **Indication(s)**
 - Nausea
 - Vomiting
- IV. **Contraindications**
 - Known hypersensitivity
 - Apomorphine use – induces profound hypotension
- V. **Precautions**
 - Caution in hepatic failure
 - Caution with antipsychotic use
- VI. **Drug interactions**
 - Apomorphine – induces profound hypotension, contraindicated
 - Antipsychotics – QT prolongation, cardiac dysrhythmias
 - Amiodarone – QT prolongation, cardiac dysrhythmias
 - Antivirals – QT prolongation, cardiac dysrhythmias
- VII. **Adult Dosing and Administration**
 - 4 mg IV (slow) / IO (ODT)
 - Repeat in 10 min
 - Max dose 8mg
- VIII. **Pediatric Dosing and Administration**
 - 0.15 mg / kg slow IV / IO (ODT)
 - < 15kg: 2mg IV/IO/IM/PO/SL (do not give PO/SL in less than 4 month old)
 - > 15kg: 4mg IV/IO/IM/PO/SL
 - Max dose 4 mg
- IX. **Onset**
 - 5-10 min
- X. **Duration**
 - 4-6 hrs
- XI. **Possible Adverse Effects**
 - Hypersensitivity / allergic reaction
 - Q-T prolongation
 - Transient blindness
 - Extrapyrimal reactions
- XII. **Other Notes**
 - Oral Dissolving Tablets (ODT) and IV administration have similar efficacy