

Sodium Bicarbonate

(Baking Soda, Citrocarbonate, Neut)

- I. Classification**
 - Alkalinizing agent
- II. Action(s)**
 - Acts as an alkalinizing agent by increasing bicarbonate ions, raising blood pH
- III. Indication(s)**
 - Hyperkalemia
 - Metabolic acidosis
 - Prolonged Cardiac Arrest
 - Tricyclic antidepressant overdose
- IV. Contraindications**
 - Metabolic or Respiratory alkalosis
 - Known hypocalcemia
- V. Precautions**
 - CHF / severe peripheral edema
 - Renal insufficiency
 - Concurrent corticosteroid therapy
 - Children with DKA (may increase risk of cerebral edema)
- VI. Drug interactions**
 - Must be cautious when pushing through IV line with several other medications (calcium chloride, Amiodarone) – flush thoroughly or start second IV line
- VII. Adult Dosing and Administration**
 - 1 mEq/kg IV / IO, max 100 mEq dose
- VIII. Pediatric Dosing and Administration**
 - Age >1 year: 1 mEq/kg IV / IO, max dose 100 mEq
- IX. Onset**
 - Immediate
- X. Duration**
 - Variable depending on patient situation
- XI. Possible Adverse Effects**
 - Fluid Overload
 - Alkalosis
 - Hypokalemia
 - Hyponatremia (elevated blood sodium levels)
- XII. Other Notes**
 - Watch IV site closely. Extravasation may cause tissue irritation or cellulitis.