

Burns Thermal

IMPORTANT NOTES ABOUT TREATMENT OF BURN PATIENTS:


- Consider intubation EARLY in burn patients with any signs of inhalation injury:
 - Soot around the nose or mouth / sooty sputum
 - Any wheezing or respiratory difficulty
 - Facial burns / singed facial hair
 - Hoarse voice / Cough / Stridor
- Use the "Rule of Nines" to estimate the percent of body surface burned (includes only 2° and 3° burns)
- Use the patient's palm as an estimate of 1% of that patient's BSA
- Avoid placing an IV in an injured / burned extremity if possible
- Note the patient injuries that require transport to a burn center

Burn Center Criteria

- Partial thickness burns greater than 10% total body surface area
- Burns that involve the face, hands, feet, genitalia, perineum, or joints.
- Third degree burns in any age group.
- Electrical burns, including lightning injury.
- Chemical burns.
- Inhalation injury.

INJURIES REQUIRING TRANSPORT TO A BURN CENTER

- Any % BSA 3° burns
- > 5 % BSA 2° burns
- Evidence of inhalation injury
- Significant burns to face, groin, hands feet, joints
- Chemical, Electrical, Radiation Burns
- Circumferential burns to thorax, hands, feet, wrists, fingers, other joints
- Persons with significant medical history
- Burns associated with other traumatic injuries



EMT/AEMT

- Follow Trauma General Patient Care Guideline
- Stop burning process
- Assess for Inhalation Injury
- Establish Basic Airway
- High flow oxygen with tight fitting non rebreather mask or BVM
- Consider Spinal Motion Restriction for explosion or history of fall
- Remove clothing and jewelry and prevent hypothermia
- Dry Dressing to areas of burn
- Plan extrication to Medic Unit

Paramedic

- Load patient and Continue High Flow Oxygenation
- Apply Monitor and evaluate for Carbon Monoxide
- Consider early intubation for inhalation or neck burns
- Start IV/IO and administer fluids according to Trauma General Patient Care Guideline
 - Modified Brooke Formula:
 - $[2\text{ml/kg} \times (\text{total \% body surface area burned})]/24\text{hrs} = \text{ml/hr in first 24 hours}$
- Pain Management Guideline (**Do not use Toradol in severe or diffuse burns**)
- Transport to Burn Center - Do not pass a trauma center if other trauma mechanisms are involved

****Bacitracin or Aquafor (preferred)** may be placed on superficial/1st Degree burns**