

Spivak, *Physics for Mathematicians: Mechanics I*

Danny Nygård Hansen

27th February 2023

5 • Rigid Bodies

Isometries

If (S, ρ) and (T, δ) are metric spaces, then a map $f: S \rightarrow T$ is an **isometry** if

$$\delta(f(x), f(y)) = \rho(x, y)$$

for all $x, y \in S$.

Let X and Y be normed vector spaces, and let $A \subseteq X$ be a subset. Since e.g. the norm on X induces a metric ρ_X given by $\rho_X(\mathbf{x}, \mathbf{x}') = \|\mathbf{x} - \mathbf{x}'\|_X$, a (not necessarily linear) map $\varphi: A \rightarrow Y$ is an isometry if

$$\|\varphi(\mathbf{x}) - \varphi(\mathbf{x}')\|_Y = \|\mathbf{x} - \mathbf{x}'\|_X$$

for all $\mathbf{x}, \mathbf{x}' \in A$. If X and Y are also inner product spaces, then we have the following:

LEMMA 5.1

Let X, Y be inner product spaces, and let $A \subseteq X$ be a subset containing $\mathbf{0}$. Let $\varphi: A \rightarrow Y$ be a map such that $\varphi(\mathbf{0}) = \mathbf{0}$. Then φ is an isometry if and only if

$$\langle \varphi(\mathbf{x}), \varphi(\mathbf{x}') \rangle_Y = \langle \mathbf{x}, \mathbf{x}' \rangle_X \quad (5.1)$$

for all $\mathbf{x}, \mathbf{x}' \in A$.

PROOF. Notice the identities

$$\|\mathbf{x} - \mathbf{x}'\|_X^2 = \|\mathbf{x}\|_X^2 - 2\langle \mathbf{x}, \mathbf{x}' \rangle_X + \|\mathbf{x}'\|_X^2$$

and

$$\|\varphi(\mathbf{x}) - \varphi(\mathbf{x}')\|_Y^2 = \|\varphi(\mathbf{x})\|_Y^2 - 2\langle \varphi(\mathbf{x}), \varphi(\mathbf{x}') \rangle_Y + \|\varphi(\mathbf{x}')\|_Y^2.$$

First assume that (5.1) holds. If we let $\mathbf{x}' = \mathbf{0}$ so that $\varphi(\mathbf{x}') = \mathbf{0}$, then $\|\varphi(\mathbf{x})\|_Y = \|\mathbf{x}\|_X$, and we similarly have $\|\varphi(\mathbf{x}')\|_Y = \|\mathbf{x}'\|_X$. The above identities then imply that φ is an isometry.

If conversely φ is an isometry, notice that

$$\|\varphi(\mathbf{x})\|_Y = \|\varphi(\mathbf{x}) - \varphi(\mathbf{0})\|_Y = \|\mathbf{x} - \mathbf{0}\|_X = \|\mathbf{x}\|_X,$$

and we similarly have $\|\varphi(\mathbf{x}')\|_Y = \|\mathbf{x}'\|_X$. The above identities then imply (5.1). \square

PROPOSITION 5.2

Every isometry $\varphi: \mathbb{R}^d \rightarrow \mathbb{R}^d$ is on the form

$$\varphi(\mathbf{x}) = A\mathbf{x} + \mathbf{b}, \quad \mathbf{x} \in \mathbb{R}^d,$$

for a unique $d \times d$ matrix A and vector $\mathbf{b} \in \mathbb{R}^d$. Furthermore, A is orthogonal.

In particular, any isometry that preserves the origin is linear.

PROOF. We first show that $\varphi(\mathbf{x}) = \varphi_0(\mathbf{x}) + \mathbf{b}$ for some $\mathbf{b} \in \mathbb{R}^d$, where $\varphi_0: \mathbb{R}^d \rightarrow \mathbb{R}^d$ is an isometry fixing the origin. Simply let $\mathbf{b} = \varphi(\mathbf{0})$ and $\varphi_0(\mathbf{x}) = \varphi(\mathbf{x}) - \mathbf{b}$. Then

$$\varphi_0(\mathbf{0}) = \varphi(\mathbf{0}) - \mathbf{b} = \mathbf{b} - \mathbf{b} = \mathbf{0},$$

and for $\mathbf{x}, \mathbf{y} \in \mathbb{R}^d$ we have

$$\|\varphi_0(\mathbf{x}) - \varphi_0(\mathbf{y})\| = \|\varphi(\mathbf{x}) - \varphi(\mathbf{y})\| = \|\mathbf{x} - \mathbf{y}\|,$$

so φ_0 is an isometry.

Next we show that φ_0 is linear; since it is an isometry, this will imply the existence of an orthogonal matrix A as above. For $\mathbf{x}, \mathbf{y}, \mathbf{z} \in \mathbb{R}^d$ we have by Lemma 5.1

$$\begin{aligned} \langle \varphi_0(\mathbf{x} + \mathbf{y}), \varphi_0(\mathbf{z}) \rangle &= \langle \mathbf{x} + \mathbf{y}, \mathbf{z} \rangle \\ &= \langle \mathbf{x}, \mathbf{z} \rangle + \langle \mathbf{y}, \mathbf{z} \rangle \\ &= \langle \varphi_0(\mathbf{x}), \varphi_0(\mathbf{z}) \rangle + \langle \varphi_0(\mathbf{y}), \varphi_0(\mathbf{z}) \rangle \\ &= \langle \varphi_0(\mathbf{x}) + \varphi_0(\mathbf{y}), \varphi_0(\mathbf{z}) \rangle. \end{aligned}$$

Since φ_0 preserves orthogonality, it maps an orthogonal basis into an orthogonal set. Since φ_0 is also injective (since it is an isometry), this set is in fact a basis. Replacing \mathbf{z} by the elements in such a basis, it follows that

$\langle \varphi_0(\mathbf{x} + \mathbf{y}), \mathbf{w} \rangle = \langle \varphi_0(\mathbf{x}) + \varphi_0(\mathbf{y}), \mathbf{w} \rangle$ for all $\mathbf{w} \in \mathbb{R}^d$. Hence $\varphi_0(\mathbf{x} + \mathbf{y}) = \varphi_0(\mathbf{x}) + \varphi_0(\mathbf{y})$. We similarly find that, for $\alpha \in \mathbb{R}$,

$$\begin{aligned} \langle \varphi_0(\alpha \mathbf{x}), \varphi_0(\mathbf{z}) \rangle &= \langle \alpha \mathbf{x}, \mathbf{z} \rangle \\ &= \alpha \langle \mathbf{x}, \mathbf{z} \rangle \\ &= \alpha \langle \varphi_0(\mathbf{x}), \varphi_0(\mathbf{z}) \rangle \\ &= \langle \alpha \varphi_0(\mathbf{x}), \varphi_0(\mathbf{z}) \rangle, \end{aligned}$$

implying that $\varphi_0(\alpha \mathbf{x}) = \alpha \varphi_0(\mathbf{x})$. Thus φ_0 is linear.

To show uniqueness, assume that $A\mathbf{x} + \mathbf{b} = \varphi(\mathbf{x}) = A'\mathbf{x} + \mathbf{b}'$ for all $\mathbf{x} \in \mathbb{R}^d$. Then $\mathbf{b} = \varphi(\mathbf{0}) = \mathbf{b}'$, so $A\mathbf{x} = A'\mathbf{x}$. Since this holds for all \mathbf{x} , it follows that $A = A'$ as desired. \square



Consider points $\mathbf{b}_1, \dots, \mathbf{b}_K \in \mathbb{R}^3$ and let $\mathbf{b} = (\mathbf{b}_1, \dots, \mathbf{b}_K)$. Let further $\mathbf{c} = (\mathbf{c}_1, \dots, \mathbf{c}_K)$ be a rigid motion of \mathbf{b} . That is, there is a one-parameter family of isometries $\varphi_t: \{\mathbf{b}_1, \dots, \mathbf{b}_K\} \rightarrow \mathbb{R}^3$ such that $\varphi_t(\mathbf{b}_i) = \mathbf{c}_i(t)$. However, to apply [Proposition 5.2](#) to φ_t it must be defined on \mathbb{R}^3 . A natural question is then whether it is possible to extend an isometry from a subset of \mathbb{R}^3 to all \mathbb{R}^3 . The answer is affirmative, as we shall see below.

Another natural question is whether such an extension is unique. If the points $\mathbf{b}_1, \dots, \mathbf{b}_K$ do not lie in a plane, then by renumbering we may assume that $\mathbf{b}_2 - \mathbf{b}_1$, $\mathbf{b}_3 - \mathbf{b}_1$, and $\mathbf{b}_4 - \mathbf{b}_1$ are linearly independent and hence span \mathbb{R}^3 . The proof below will show that this implies that an extension of φ_t is unique.

LEMMA 5.3

Let $A \subseteq \mathbb{R}^d$ and let $\varphi: A \rightarrow \mathbb{R}^d$ be an isometry. Then φ extends to an isometry $\tilde{\varphi}: \mathbb{R}^d \rightarrow \mathbb{R}^d$.

Furthermore, we have the following uniqueness results:

- (i) If A contains $d + 1$ affinely independent elements, then φ has a unique isometric extension to \mathbb{R}^d .
- (ii) If A contains d affinely independent elements, then φ has a unique isometric extension to \mathbb{R}^d that is also orientation preserving.¹

PROOF. By translation we may assume that $\mathbf{0} \in A$ (if A is empty then the claim is obvious) and that $\varphi(\mathbf{0}) = \mathbf{0}$. First extend φ by linearity to $\text{span } A$. This is well-defined, since if $\mathbf{x}_1, \dots, \mathbf{x}_n \in A$ and $\alpha_1, \dots, \alpha_n \in \mathbb{R}$, then by [Lemma 5.1](#) we

¹ By this we mean that if $\tilde{\varphi}(\mathbf{x}) = A\mathbf{x} + \mathbf{b}$ in accordance with [Proposition 5.2](#), then $A \in \text{SO}(d)$.

have

$$\left\| \sum_{i=1}^n \alpha_i \varphi(\mathbf{x}_i) \right\|^2 = \sum_{i,j=1}^n \alpha_i \alpha_j \langle \varphi(\mathbf{x}_i), \varphi(\mathbf{x}_j) \rangle = \sum_{i,j=1}^n \alpha_i \alpha_j \langle \mathbf{x}_i, \mathbf{x}_j \rangle = \left\| \sum_{i=1}^n \alpha_i \mathbf{x}_i \right\|^2.$$

This also shows that φ extends to an isometry on $\text{span } A$, and in particular φ is linear by [Proposition 5.2](#).

Since φ is linear it maps $\text{span } A$ to a subspace $\varphi(\text{span } A)$ of \mathbb{R}^d with the same dimension. Let U and V be the orthogonal complements of $\text{span } A$ and $\varphi(\text{span } A)$ respectively, and let $\psi: U \rightarrow V$ be an isometry. Letting $\tilde{\varphi} = \varphi \oplus \psi$, it follows easily by Pythagoras' theorem that $\tilde{\varphi}$ is also an isometry as desired.

For the first uniqueness claim, notice that if A contains a collection of $d + 1$ affinely independent elements and \mathbf{a} is one element in such a collection, then $A - \mathbf{a}$ contains d linearly independent vectors. Hence its span, which is contained in $\text{span } A$, is d -dimensional.

To prove the second uniqueness claim, notice that an argument similar to the above shows that $\text{span } A$ has dimension at least $d - 1$. Then notice that if $\dim U = 1$, then there are exactly two isometries $U \rightarrow V$, only one of which makes $\tilde{\varphi}$ orientation preserving. \square

Motion relative to the centre of mass

In the section ‘The inertial tensor’, Spivak considers a rigid motion $\mathbf{c} = (\mathbf{c}_1, \dots, \mathbf{c}_K)$ of a collection of points $\mathbf{b}_1, \dots, \mathbf{b}_K$. These are given by $\mathbf{c}_i(t) = B(t)\mathbf{b}_i + \mathbf{w}(t)$ for some $B: \mathbb{R} \rightarrow \text{SO}(3)$ and $\mathbf{w}: \mathbb{R} \rightarrow \mathbb{R}^3$.

Spivak then claims that we may choose \mathbf{w} to be the centre of mass \mathbf{C} given by

$$\mathbf{C} = \frac{1}{M} \sum_{i=1}^K m_i \mathbf{c}_i, \quad \text{where} \quad M = \sum_{i=1}^K m_i.$$

Given the uniqueness part of [Proposition 5.2](#), it is not immediately clear that this is possible, even if it is intuitively reasonable. Inserting the rigid motion in the expression for \mathbf{C} , we get

$$\begin{aligned} \mathbf{C}(t) &= \frac{1}{M} \sum_{i=1}^K m_i (B(t)\mathbf{b}_i + \mathbf{w}(t)) = \frac{1}{M} \sum_{i=1}^K m_i B(t)\mathbf{b}_i + \mathbf{w}(t) \\ &= B(t) \frac{\sum_{i=1}^K m_i \mathbf{c}_i(0)}{M} + \mathbf{w}(t) = B(t)\mathbf{C}(0) + \mathbf{w}(t). \end{aligned}$$

That is, the centre of mass has the exact same time evolution as each of the individual particles. If we choose our coordinate system such that $\mathbf{C}(0) = \mathbf{0}$, we thus get $\mathbf{C}(t) = \mathbf{w}(t)$.

Notice that this does not mean that we consider the motion as seen from the centre of mass system. We simply choose an inertial system in which $\mathbf{C}(0) = \mathbf{0}$.