高能天体物理 arXiv 每周学术脉络 · 2026 年第 44 周 (2026-W44) 深度精读研报

报告周期：2026 年第 44 周 (2026-W44) (2026-W44) **时间跨度：**2026-10-26 ~ 2026-11-01 **纳入精读：**10 篇核心必读 | 76 篇重点关注

在线精读：https://astro-lineage.pages.dev/arxiv-weekly/2026-W44/

★ 本周精选速览矩阵 (At-a-Glance Visual Triage · 共 20 篇)

1-MINUTE VISUAL TRIAGE • KEY INSIGHTS & FIGURES

快速研读指引：下表浓缩本周全部 10 篇核心必读与 10 篇特选关注论文的**一句话核心结论 (Key Insight)**、**关键科学图表**与**直达超链接**（点击页内编号跳至精读详情，点击 arXiv / PDF 直达原文），助您在 1–2 页内快速判断哪些文章值得精读。

#01 ⭐ 必读 **arXiv:2610.31758** ↓详情 · arXiv · PDF · 网页

Collapsars Are Not Heavy r-Process Factories

Collapsars Are Not Heavy r-Process Factories

Key Insight： Strong magnetic unbinding necessitates high inflow velocities $v_{\rm in}$, which shorten transit times and cap the net neutronization integral such that...




Fig. 3 · 核心物理演化图

#02 ⭐ 必读 **arXiv:2610.32861** ↓详情 · arXiv · PDF · 网页

GRB 260924A: Extreme Energetics, Bright GeV Emission, and Exceptional X-ray and Optical Luminosity

GRB 260924A: Extreme Energetics, Bright GeV Emission, and Exceptional X-ray and...

Key Insight： A colossal isotropic energy release ($E_{\gamma,\rm iso} \simeq 9.79 \times 10^{54}$ erg) expanding into a stratified wind medium ($n \propto R^{-1.2}$) generates 43.31 GeV photons via reverse-shock SSC processes with microphysical magnetization...

#03 ⭐ 必读 **arXiv:2610.34217** ↓详情 · arXiv · PDF · 网页

EP241021a as An Off-nuclear Month-duration Transient in a Dwarf Starburst Galaxy

EP241021a as An Off-nuclear Month-duration Transient in a...

Key Insight： HST/Gemini high-resolution astrometry establishes a 0.95 kpc offset from an edge-on dwarf starburst nucleus, where a month-long plateau ($L \sim$ ...




Fig. 3 · 数据拟合与模...

#04 ⭐ 必读 **arXiv:2610.34810** ↓详情 · arXiv · PDF · 网页

EP251129a: A Fast X-ray Transient with Two X-Ray Plateaus and Its Off-axis Afterglow

EP251129a: A Fast X-ray Transient with Two X-Ray Plateaus and Its Off-axis Afterglow

Key Insight： The sharp drop (decay index $\alpha_{\rm X,2} = 5.28$) and secondary plateau up to 42 d reveal a supramassive magnetar collapsing into a black hole that undergoes delayed fallback accretion energy injection ($E_{\rm inj} \simeq 6 \times$ ...

#05 ⭐ 必读 **arXiv:2610.35539** ↓详情 · arXiv · PDF · 网页

An Intermediate Mass Stripped Star in a 4.2-hour Orbit around a Compact Object

An Intermediate Mass Stripped Star in a 4.2-hour Orbit around...

Key Insight： Radial-velocity semi-amplitude $K = 151.4\ \mathrm{km\,s^{-1}}$ and ellipsoidal modulations establish a companion mass of $M_2 = 1.58\ M_\odot$ in a 4.2 h orbit,...




Fig. 3 · 核心物理演化图

#06 ⭐ 必读 **arXiv:2610.35619** ↓详情 · arXiv · PDF · 网页

Indication of Electron Antineutrinos from the Diffuse Supernova Neutrino Background

Indication of Electron Antineutrinos from the Diffuse Supernov...

Key Insight： Unbinned extended maximum likelihood fitting across pure-water and Gd-loaded SK phases yields a best-fit $\bar{\nu}_e$ flux of $3.6^{+1.6}_{-1.5}\ \mathrm{cm^{-2}\,s^{-1}}$ for $E_\nu >$ ...




Fig. 1 · 核心观测结果

#07 ⭐ 必读 **arXiv:2610.35675** ↓详情 · arXiv · PDF · 网页

The design and flight of the Payload for Ultrahigh Energy Observations (PUEO)

The design and flight of the Payload for Ultrahigh Energy...

Key Insight： A 96-horn dual-polarization array synthesized into 48 real-time coherent beams via 26 RFSoC FPGAs achieved 50% trigger efficiency at ...




Fig. 1 · 物理模型示意图

#08 ⭐ 必读 **arXiv:2610.37044** ↓详情 · arXiv · PDF · 网页

Discovery of a New Electron Spectrum Produced by Fast Cooling of Relativistic···

Discovery of a New Electron Spectrum Produced by Fast Coolin...

Key Insight： Non-steady fast cooling transforms the sub-injection electron spectrum into an evolving smooth broken power law $f_{\rm U}(\gamma_e, t) = A(t)\gamma_e^{-2}[1 + \cdots$




Fig. 1 · Physical...

#09 ⭐ 必读 **arXiv:2610.37214** ↓详情 · arXiv · PDF · 网页

Coherent Radio Emission from Flux Rope Mergers in 3D Relativistic Reconnection

Coherent Radio Emission from Flux Rope Mergers in 3D...

Key Insight： Flux rope mergers in 3D pair-plasma reconnection sheets generate 100% linearly polarized X-mode waves with radiation efficiency $\eta \sim (1\text{–}2) \times 10^{-3}$...




Fig. 4 · Physical Mod...

#10 ⭐ 必读 **arXiv:2610.38673** ↓详情 · arXiv · PDF · 网页

Black Hole Spin Evolution in Magnetized Accretion Flows

Black Hole Spin Evolution in Magnetized Accretion Flows

Key Insight： Horizon electromagnetic torques cause black hole equilibrium spin to decrease monotonically with normalized flux $\tilde{\phi}$, tumbling to $a_{*;\rm eq} \approx 0.05\text{–}0.06$ i...




Fig. 1 · Physical...

#11 📌 关注 **arXiv:2610.31754** ↓详情 · arXiv · PDF · 网页

Image Morphology of Misaligned Magnetically Arrested Disks

Image Morphology of Misaligned Magnetically Arrested Disks

Key Insight： While high-spin prograde MADs align naturally near the horizon via jet torques, intermediate and retrograde tilted systems maintain asymmetric...




Fig. 2 · 核心物理演化图

#12 📌 关注 **arXiv:2610.31755** ↓详情 · arXiv · PDF · 网页

From Inspiral to Outburst: Modeling a Potential Luminous Red Nova from···

From Inspiral to Outburst: Modeling a Potential Luminous Red...

Key Insight： Tidal disruption and orbital energy deposition of a 0.7–2.0 $M_\odot$ companion inside a red supergiant launches a shock breakout yielding peak...




Fig. 1 · 物理模型示意图

#13 📌 关注 **arXiv:2610.31990** ↓详情 · arXiv · PDF · 网页

Tilt, warp, and dissipation flare in hydrodynamic disks around kicked compa···

Tilt, warp, and dissipation flare in hydrodynamic disks around...

Key Insight： Impulsive compact object kicks induce hydrodynamic self-collisions within $r_0 \equiv GM/V_{\rm kick}^2$ that dissipate 5–30% of the bound kinetic energy, driving a...



Fig. 1 · 物理模型示意图

#14 📌 关注 **arXiv:2610.32028** ↓详情 · arXiv · PDF · 网页

Magnetic field effect on neutral color superconductivity as a mechanism to boost the inner magnetic field of magnetars

Magnetic field effect on neutral color superconductivity as a mechanism to boost the...

Key Insight： In the neutral 2SC phase, Fermi surface mismatch triggers tachyon gluon Meissner masses and paramagnetic anti-screening, naturally boosting internal magnetic fields to $\bar{B} \sim 10^{17}$ G from modest initial seed...

#15 关注 arXiv:2610.32047 ↓详情 · arXiv · PDF · 网页

A Unified Model for Radioactive Signatures and Source Identifiability in Neutron Star⋯

A Unified Model for Radioactive Signatures and Source...




Fig. 4 · 核心物理演化图

Key Insight: Thermalization delays and multi-layer diffusion smooth radioactive decay pulses such that models with and without ^{254}Cf achieve $< 9\%$ statistica...

#16 关注 arXiv:2610.32168 ↓详情 · arXiv · PDF · 网页

SMMAN Studies of AGN Variability. I. Long-term Spectral Evolution and a Candidate⋯

SMMAN Studies of AGN Variability. I. Long-term Spectral...




Fig. 3 · 核心观测结果

Key Insight: A 1026.7 d periodic flux modulation with 5.45% fractional amplitude and zero inter-band lag is uncovered, naturally accounted for by a slight...

#17 关注 arXiv:2610.32316 ↓详情 · arXiv · PDF · 网页

Radio Galaxies as UHECR Sources in Light of Recent Microquasar Results

Radio Galaxies as UHECR Sources in Light of Recent Microquasar Results

Key Insight: Microquasar SS 433 synchrotron cutoffs confirm near-Bohm electron acceleration, implying that radio galaxy jet shear layers can accelerate UHECR protons to > 10 EeV during episodic bursts like Cen A's...

#18 关注 arXiv:2610.32323 ↓详情 · arXiv · PDF · 网页

A Framework for Analysing Nebular-Phase JWST Supernova Spectra with Application to Type Ic SN 2024aecx

A Framework for Analysing Nebular-Phase JWST Supernova Spectra with Application t...

Key Insight: Self-consistent analysis of [Ni II] and [Co II] lines reveals a low iron-zone filling factor ($f_{\rm FeHe} = 0.059$), proving that radioactive ^{56}Ni bubble expansion was negligible, while chlorine yields favor an O-C shell merger.

#19 关注 arXiv:2610.32430 ↓详情 · arXiv · PDF · 网页

Local Analogs of Little Red Dots: Clues to the X-ray Weakness

Local Analogs of Little Red Dots: Clues to the X-ray Weakness

Key Insight: Stacking detection of local LRD analogs at $\log L_{2-10} = 42.55\ \mathrm{erg\,s^{-1}}$ reveals a relatively hard X-ray spectrum, ruling out unshielded super-Eddington accretion and Compton-thick obscuration in favor of mild...

#20 关注 arXiv:2610.32786 ↓详情 · arXiv · PDF · 网页

Probing the star-forming environment of GRB across cosmic ages with high-resolution X-ray spectroscopy: From XMM-⋯

Probing the star-forming environment of GRB across cosmic ages with high-resolution...

Key Insight: Non-equilibrium ionization modeling with TEPID reveals that XRISM and NewAthena will measure host gas densities to 0.05 dex and distinguish Pop III from Pop II progenitors via narrow Fe-K absorption line...

1. 本周综述与宏观脉络 (Executive Synthesis · 点击编号直达精读)

EDITORIAL OVERVIEW

ABSTRACT / 学术总览（点击文内 ARXIV 编号 ↓ 可直接跳至下方详细精读）

This week's progress spans the causal chain of high-energy transients from black hole spin equilibria and magnetohydrodynamic outflows to relativistic shock radiation, core-collapse neutrino backgrounds, and non-canonical engines. Major advances include:

- **Collapsar Nucleosynthesis & Engine Physics**: arXiv:2610.31758 ↓ proved analytically that neutrino-dominated accretion flows in collapsars cannot synthesize heavy r-process elements because magnetically launched unbinding demands high inflow velocities that suppress electron captures, establishing a strict lower bound of $Y_{\rm e}^{\rm min} > 0.28$; concurrently, arXiv:2610.38673 ↓ demonstrated via 3D GRMHD that magnetized accretion flows drive the black hole equilibrium spin downward from $a_{\star;\rm eq} \approx 0.9$ (SANE) to $a_{\star;\rm eq} \approx 0.05$–$0.06$ at MAD saturation.
- **Relativistic Jets, Microphysics & Gamma-Ray Bursts**: arXiv:2610.32861 ↓ mapped the colossal energetics ($E_{\gamma,\rm iso} \simeq 9.79 \times 10^{54}$ erg) of GRB 260924A, attributing excess ~ 43 GeV photons to reverse-shock synchrotron self-Compton scattering in a stratified wind ($k \simeq 1.2$); arXiv:2610.37044 ↓ solved the non-steady kinetic evolution of fast-cooling relativistic electrons below γ_m, deriving an exact smooth broken power-law spectrum $f_{\rm U}(\gamma_e, t) \propto \gamma_e^{-2}[1 + \gamma_0(t)/\gamma_e]^{-\beta}$; arXiv:2610.37214 ↓ demonstrated via 3D PIC simulations that 100% polarized X-mode coherent radio waves are robustly generated via flux rope mergers in relativistic reconnection with efficiency $\eta \sim 10^{-3}$.
- **Exotic Engines & Fast X-Ray Transients**: arXiv:2610.34217 ↓ resolved the off-nuclear transient EP241021a in a dwarf starburst host, demonstrating that its month-long plateau and coordinated optical-X-ray plunge exclude standard core-collapse and SMBH models, favoring an intermediate-mass black hole or micro-TDE; arXiv:2610.34810 ↓ modeled EP251129a as an off-axis two-component jet whose double X-ray plateaus and delayed radio afterglow reflect a supramassive magnetar collapsing to a black hole fed by fallback accretion ($E_{\rm inj} \simeq 6 \times 10^{51}$ erg).
- **Compact Binaries & Multi-Messenger Particle Astrophysics**: arXiv:2610.35539 ↓ discovered an intermediate-mass stripped star orbiting a neutron star in a 4.2 h orbit, confirming the long-sought progenitor phase to Case BB mass transfer and double neutron star mergers; arXiv:2610.35619 ↓ reported a 2.6σ indication of the Diffuse Supernova Neutrino Background using 5003.0 d of Super-Kamiokande data with an electron antineutrino flux of $3.6^{+1.6}_{-1.5}\ \mathrm{cm^{-2}\,s^{-1}}$ ($E_\nu > 13.3$ MeV); arXiv:2610.35675 ↓ reported the maiden flight of PUEO, demonstrating real-time RFSoC interferometric phased-array triggering down to $\mathrm{SNR} = 1.4$ for ultrahigh-energy neutrinos.

2. ⭐ 核心必读文献深度解析 (Must-Read In-Depth Deep Dives · 按专题分组)

10 ARTICLES IN 1 THEMES

专题 01 综合前沿课题 (General Highlights)

本专题收录 10 篇必读文献

PAPER 01 ⭐ 必读精选 🧭 综合前沿课题 (General Highlights) arXiv:2610.31758

[arXiv:2610.31758] Collapsars Are Not Heavy r-Process Factories

Collapsars Are Not Heavy r-Process Factories

作者：Ore Gottlieb, Danat Issa, Jonatan Jacquemin-Ide arXiv 原文 | PDF 直达 | SciX (ADS) | 站内交互雷达

RESEARCH HIGHLIGHT & BREAKTHROUGH / 核心突破与推荐理由

【核心突破】： Strong magnetic unbinding necessitates high inflow velocities $v_{\rm in}$, which shorten transit times and cap the net neutronization integral such that $Y_{\rm e}^{\rm min} > 0.28$ across all magnetic configurations under realistic GRB jet power constraints.

【推荐理由】： Solves a fundamental nucleosynthesis controversy by proving analytically that magnetically unbinding collapsar disks cannot produce heavy r-process elements.

🔍 科学问题与背景 (Problem)

探讨坍缩星（collapsar）中微子冷却吸积盘（NDAF）风能否产生重 r-过程核素。核心物理矛盾在于：重 r-过程要求极低电子丰度（$Y_{\rm e} \lesssim 0.25$），依赖极高密度和足够长的内流时标以进行电子俘获中子化；而将物质有效抛射需要强磁应力做功解绑，但强磁场会导致内流速度 $v_{\rm in}$ 增加、缩短时标并降低盘密度，从而破坏中子化条件。

🛠️ 观测与研究方法 (Methodology)

结合 Blandford-Znajek 机制喷流功率标度 $P_j = \eta_a \eta_\phi \dot{M}_{\rm BH} c^2$（其中 $\eta_a \approx 1.06a^4 + 0.395a^2$，$\eta_\phi \approx (\phi_{\rm BH}/50)^2$）与长 GRB 观测上限，构建了从弱磁通（SANE）到磁阻碍吸积盘（MAD）全磁通序列的吸积流模型。通过在每个半径处选取最大化净中子化率的局域最优温度 $kT_\star \simeq 2.3\,{\rm MeV}\,(\rho/10^9\,{\rm g\,cm^{-3}})^{1/2}$，并利用轻微电子简并展开构造包络速率 $\Gamma_{\rm env}(\rho) \equiv 8\,(\rho/10^9\,{\rm g\,cm^{-3}})^2\,{\rm s^{-1}}$，沿吸积速度 $v_{\rm in}(r) = \zeta(r)[H(r)/r]v_K(r)$ 向内积分推导到达解绑半径 $R_{\rm unb} = 10\,r_g$ 处的最小可能 $Y_{\rm e}$。

📐 物理模型与关键假设 (Models & Assumptions)

- 外流气体必须在解绑半径 $R_{\rm unb} = 10\,r_g$ 或之外由大尺度磁应力做功实现非束缚。

⚠️ 局限性与观测边界 (Limits & Caveats)




Fig. 3 [核心物理演化图]： Figure 3: Minimum $Y_{\rm e}(R_{\rm unb})$ in $(\dot{M}_{\rm d}, \phi_{\rm BH})$. The thin solid (dashed) line shows the $Y_{\rm e} = 0.25$ contour using the numerical (analytic) estimate. The thick line delineates the ceiling of the GRB population: $\dot{M}_{\rm BH,max} = 0.4\,M_\odot\,{\rm s^{-1}}\,(20/\phi_{\rm BH})^2$, for $P_j = 5 \times 10^{51}\,{\rm erg\,s^{-1}}$, $a = 0.3$, and $M_{\rm BH} = 3\,M_\odot$. Along the GRB ceiling, $Y_{\rm e}^{\rm min} \gtrsim 0.33$ for any value of $\phi_{\rm BH}$.




Fig. 4 [核心物理演化图]： Figure 4: Most favorable composition bound as a function of black hole spin a and isotropic-equivalent γ-ray luminosity $L_{\rm iso}$. Colors show $Y_{\rm e}^{\rm min}(R_{\rm unb})$ derived from Eq. (22) with $\eta_a(a)$ of Eq. (2), assuming $M_{\rm BH} = 3\,M_\odot$, $\theta_j = 0.1$, and $\eta_\gamma = 0.2$, and minimizing over $20 \le \phi_{\rm BH} \le 50$ (minimum is reached at $\phi_{\rm BH} = 20$). The circle marks the ceiling reference model ($a = 0.3$, $P_j = 5 \times 10^{51}\,{\rm erg\,s^{-1}}$), for which $Y_{\rm e}^{\rm min} \approx 0.33$. The solid curve marks $Y_{\rm e}^{\rm min} = 0.25$ from the full numerical calculation and the dashed curve $\mathcal{B}_{\rm env} = 3/4$, where the analytic bound of Eq. (29) reaches $Y_{\rm e} = 0.25$. Thin gray curves show $\dot{M}_{\rm BH}(\phi_{\rm BH}{=}20) = 0.01$, 0.1, and $10\,M_\odot\,{\rm s^{-1}}$. Horizontal lines mark observed isotropic-equivalent peak luminosities: the break of the long-GRB luminosity function, $L_\star = 10^{52.5\pm0.2}\,{\rm erg\,s^{-1}}$ (line and shaded uncertainty; Wanderman and Piran 103) and GRB 221009A ($9.9 \times 10^{53}\,{\rm erg\,s^{-1}}$; Lesage et al. 55). Peak luminosities overestimate the sustained jet power and hence $\dot{M}_{\rm BH}$: GRB 221009A radiated $E_{\gamma,{\rm iso}} = 1.0 \times 10^{55}$ erg over more than 600 s at $z = 0.151$ [55], corresponding to a mean luminosity of $\lesssim 2 \times 10^{52}\,{\rm erg\,s^{-1}}$. The color scale is linear in $Y_{\rm e}^{\rm min}$ below 0.25 (blue) and logarithmic in $0.5 - Y_{\rm e}^{\rm min}$ above it (red).

📊 关键结论与定量测量 (Quantitative Results)

坍缩星吸积盘在任何磁通序列下均无法向外抛射电子丰度 $Y_{\rm e} \le 0.25$ 的重 r-过程气体。在长 GRB 喷流各向同性峰值功率上限 $P_j \lesssim 5 \times 10^{51}\,{\rm erg\,s^{-1}}$ 与吸积率 $\dot{M}_{\rm BH} \le 0.4\,M_\odot\,{\rm s^{-1}}$ 限制下，在有效磁解绑半径 $R_{\rm unb} = 10\,r_g$ 处，气体所能达到的理论最低电子丰度随无量纲磁通参数 $\phi_{\rm BH}$ 变化，从发射喷流阈值处的 $Y_{\rm e}^{\rm min}(\phi_{\rm BH} = 20) \simeq 0.33$ 单调上升至 MAD 极限下的 $Y_{\rm e}^{\rm min}(\phi_{\rm BH} = 50) \simeq 0.49$。即使采用速率极大化温度，净中子化率积分上限 $\mathcal{B}_{\rm env} \equiv \int_{R_{\rm unb}}^{R_{\rm out}} [\Gamma_{\rm env}/v_{\rm in}]\,dr \lesssim 0.51$，导出严格理论下限 $Y_{\rm e}^{\rm min}(R_{\rm unb}) \gtrsim 1/(2\sqrt{1+4\mathcal{B}_{\rm env}}) > 0.28$。

🎯 建议精读切入点 (Reading Guide)

Focus on Section 2 for the construction of the maximal neutronization envelope $\Gamma_{\rm env}(\rho)$ and Section 3 for the inflow integration bound as a function of $\phi_{\rm BH}$.

PAPER 02 ⭐ 必读精选 🧭 综合前沿课题 (General Highlights) arXiv:2610.32861

[arXiv:2610.32861] GRB 260924A: Extreme Energetics, Bright GeV Emission, and Exceptional X-ray and Optical Luminosity

GRB 260924A: Extreme Energetics, Bright GeV Emission, and Exceptional X-ray and Optical Luminosity

作者：Nissim Fraija, Peter Veres, Rosa L. Becerra, Antonio Galván-Gámez, Thomas Freeman 等 arXiv 原文 | PDF 直达 | SciX (ADS) | 站内交互雷达

RESEARCH HIGHLIGHT & BREAKTHROUGH / 核心突破与推荐理由

【核心突破】： A colossal isotropic energy release ($E_{\gamma,\rm iso} \simeq 9.79 \times 10^{54}$ erg) expanding into a stratified wind medium ($n \propto R^{-1.2}$) generates 43.31 GeV photons via reverse-shock SSC processes with microphysical magnetization ratio $R_B \simeq 2.6$.

【推荐理由】： Presents comprehensive observations and modeling of one of the most energetic GRBs on record, confirming thick-shell reverse-shock SSC emission.

🔍 科学问题与背景 (Problem)

解释极端高能暴 GRB 260924A 的能量学分配机制、在 $t \sim 82\,\rm s$ 到达的 $43.31\,\rm GeV$ 等光子为何超出标准前向激波同步辐射理论极限 $E^{\rm syn}_{\rm max}$，以及早期 LAT 辐射过剩、X 射线再增亮与成层星周介质相互作用的微观物理图景。

🛠️ 观测与研究方法 (Methodology)

利用 Fermi/GBM、Fermi/LAT、Swift/BAT、Swift/XRT 以及多台地面光学/近红外望远镜进行多波段时变与时变能谱分析。通过非成箱极大似然法提取 LAT 能谱（0.1–$1000\,\rm GeV$）与高能光子关联概率；在成层星周介质密度轮廓 $n(R) = n_{\rm dec}(R/R_{\rm dec})^{-k}$（固定 $k = 1.2$）框架下，采用 `emcee` 进行前向激波同步辐射 + SSC 以及反向激波 SSC 联合马尔可夫链蒙特卡洛（MCMC）后验参数约束，同时包含曲率效应（$\alpha = 2 + \beta$）与 Klein-Nishina 截断检验。

📐 物理模型与关键假设 (Models & Assumptions)

⚠️ 局限性与观测边界 (Limits & Caveats)

- 晚期 X 射线与光学数据中未观测到清晰稳健的喷流转折（jet break），因而无法直接确定喷流开角或几何束流修正后的真实能量释放。

📊 关键结论与定量测量 (Quantitative Results)

测定暴源红移为 $z = 3.51 \pm 0.10$，1–10,000 keV 能段各向同性释放能量高达 $E_{\gamma,\rm iso} \simeq 9.79 \times 10^{54}\,\rm erg$，初始各向同性动能 $E_{\rm K,iso} \simeq 7.0 \times 10^{55}\,\rm erg$，瞬时辐射效率 $\eta_\gamma \simeq 0.12$。Fermi/LAT 探测到最高达 $43.31\,\rm GeV$ 的高能光子（在 $82.24\,\rm s$ 到达）。晚期余辉衰减指数为 $\alpha_{\rm opt} = 1.46 \pm 0.01$ 与 $\alpha_X = 1.57 \pm 0.04$，在慢冷却闭合关系 $\alpha^{\rm syn} = \frac{3(p-1)}{4} + \frac{k}{2(4-k)}$ 下排除了常规均匀介质（$k = 0$）或纯自由星风（$k = 2$），测定星周介质为密度梯度介质 $n(R) \propto R^{-k}$（$k \simeq 1.0$–1.3，基准 $k = 1.2$），电子谱指数 $p \simeq 2.75$。早期 70.04–$164.9\,\rm s$ 处的 LAT 通量超出后续外推达 2.25 ± 0.59 倍，由厚壳层反向激波 SSC 过程主导解释；激波微观磁能分配比为 $R_B \equiv (\varepsilon_{B_r}/\varepsilon_{B_f})^{1/2} \simeq 2.6$。

🎯 建议精读切入点 (Reading Guide)

Examine the multi-band light curves and Section 4's MCMC joint afterglow modeling incorporating reverse- and forward-shock closure relations.

PAPER 03 ⭐ 必读精选 🧭 综合前沿课题 (General Highlights) arXiv:2610.34217

[arXiv:2610.34217] EP241021a as An Off-nuclear Month-duration Transient in a Dwarf Starburst Galaxy

EP241021a as An Off-nuclear Month-duration Transient in a Dwarf Starburst Galaxy

作者：Luming Sun, Xinwen Shu, Jiazheng Zhu, Ning Jiang, Bing Zhang 等 arXiv 原文 | PDF 直达 | SciX (ADS) | 站内交互雷达

RESEARCH HIGHLIGHT & BREAKTHROUGH / 核心突破与推荐理由

【核心突破】：HST/Gemini high-resolution astrometry establishes a $0.95\ \mathrm{kpc}$ offset from an edge-on dwarf starburst nucleus, where a month-long plateau ($L \sim 10^{43}\ \mathrm{erg\,s^{-1}}$) terminating in a steep synchronized drop points to an intermediate-mass black hole ($M_{\rm BH} \lesssim 10^4\ M_\odot$) micro-TDE.

【推荐理由】：Reports a remarkable off-nuclear Einstein Probe transient whose month-long duration challenges standard core-collapse and SMBH-TDE paradigms.

🔍 科学问题与背景 (Problem)

Accounting for the multi-epoch optical rebrightenings (at ~ 7.5 d with amplitude ~ 2 mag and at ~ 20 d), the ~ 1-month flat X-ray plateau, and the subsequent coordinated steep drop at $40-50$ d, which break standard single-jet core-collapse supernova and compact binary merger models whose central engines shut down on timescales $\lesssim 10^3 - 10^4$ s.

📐 物理模型与关键假设 (Models & Assumptions)

🛠️ 观测与研究方法 (Methodology)

Multi-band high-resolution imaging using HST/WFC3 (F606W, F814W, F110W, F160W) and Gemini-N/GMOS (r, z) combined with Gaia DR3 absolute astrometric calibration (galaxy centroid uncertainty ~ 17 mas) to measure the offset relative to VLBA coordinates; host-galaxy SED modeling via CIGALE to constrain $M_\star$, SFR, age, and extinction; and host-subtracted light curve modeling parameterizing the post-break drop as jet cessation driven by cessation of accretion.

⚠️ 局限性与观测边界 (Limits & Caveats)




Fig. 3 [数据拟合与模型对比]： Figure 3: (a): The host-subtracted optical light curves of the transient EP241021a in the r and z bands, along with the X-ray light curves during and after the plateau phase. The data were taken from [43] and [9]. The rebrightening, the subsequent power-law decay, and the final sharp decline are labeled in both bands. (b): The broken power-law models that fit the r -band data after a phase of 7.5 days. We present 200 randomly selected MCMC realizations for clarity. (c): The posterior distributions for each model parameter.

📊 关键结论与定量测量 (Quantitative Results)

EP241021a ($z = 0.748$, $D_L = 4.620$ Gpc, $E_{\rm iso} \lesssim 2 \times 10^{51}$ erg) occurred at a projected offset of 129 ± 18 mas (0.95 ± 0.13 kpc) from the nucleus of an edge-on dwarf starburst galaxy ($M_\star = (2.7 \pm 1.2) \times 10^8\ M_\odot$, SFR $= 4.5 \pm 2.4\ M_\odot\ \mathrm{yr}^{-1}$, effective radius $R_e = 0.23'' \approx 1.7$ kpc). The host-subtracted optical transient underwent a rapid break/drop at observer-frame $t_{\rm brk} = 47.0^{+5.4}_{-3.7}$ d (rest-frame $26.9^{+3.1}_{-2.1}$ d) at $\log(L_{\rm brk}/[\mathrm{erg\ s^{-1}}]) = 42.92 \pm 0.08$, synchronized with the X-ray light-curve steep decline. The month-long accretion timescale and off-nuclear site exclude both standard core-collapse supernova engines and supermassive black holes, favoring an intermediate-mass black hole ($M_{\rm BH} \lesssim 10^4\ M_\odot$) or a stellar-mass compact object micro-TDE undergoing repeating partial disruptions.

🎯 建议精读切入点 (Reading Guide)

Read Section 3 for the precise astrometric offset determination relative to VLBA coordinates and Section 5 for the accretion shutdown modeling.

PAPER 04 ⭐ 必读精选 🧭 综合前沿课题 (General Highlights) arXiv:2610.34810

[arXiv:2610.34810] EP251129a: A Fast X-ray Transient with Two X-Ray Plateaus and Its Off-axis Afterglow

EP251129a: A Fast X-ray Transient with Two X-Ray Plateaus and Its Off-axis Afterglow

作者：Yue Wu, Ailing Wang, Senlin Pang, Yu-Han Yang, Biao Zhang 等 arXiv 原文 | PDF 直达 | SciX (ADS) | 站内交互雷达

RESEARCH HIGHLIGHT & BREAKTHROUGH / 核心突破与推荐理由

【核心突破】：The sharp drop (decay index $\alpha_{\rm X,2}=5.28$) and secondary plateau up to 42 d reveal a supramassive magnetar collapsing into a black hole that undergoes delayed fallback accretion energy injection ($E_{\rm inj}\simeq 6\times10^{51}$ erg, $M_{\rm acc}\simeq 0.2\,M_\odot$) into an off-axis jet.

【推荐理由】：Provides a definitive case study of a fast X-ray transient exhibiting a sharp internal collapse drop followed by late-time fallback accretion.

🔍 科学问题与背景 (Problem)

Investigating the physical origin of the rare double X-ray plateau, the abrupt decay accompanied by strong spectral softening, and the delayed, rapidly rising radio afterglow in the extragalactic fast X-ray transient EP251129a.

🛠️ 观测与研究方法 (Methodology)

Multi-wavelength data analysis combining EP-FXT, Swift-XRT, Chandra, optical facilities, and radio telescopes (VLA, e-MERLIN). Theoretical modeling incorporates millisecond magnetar spin-down dominated by gravitational wave radiation ($L_{\rm EM}(t)\simeq L_{\rm em,0}(1+t/\tau_{\rm c,gw})^{-1}$), collapse to a black hole, dynamic forward shock in a wind medium ($\rho=AR^{-2}$) with a two-component off-axis jet (core angle $\theta_{\rm c}$, wing angle $\theta_{\rm j}$), and Blandford-Znajek fallback accretion energy injection ($L_{\rm BZ}\propto\dot{M}$).

📐 物理模型与关键假设 (Models & Assumptions)

- The early internal plateau is powered by the spin-down dissipation of a supramassive millisecond magnetar initially dominated by gravitational wave emission.
- Late-time afterglow emission originates from an off-axis two-component jet propagating into a stellar wind medium ($\rho=AR^{-2}$) with fallback accretion energy injection via the Blandford-Znajek mechanism.

⚠️ 局限性与观测边界 (Limits & Caveats)

- Late-time X-ray photon statistics are limited, requiring tied photon indices across adjacent epochs.
- A pure black hole fallback model for the early internal plateau is disfavored due to strong off-axis Doppler de-beaming ($F_{\nu,\rm off}\approx a^{3+\beta_X}F_{\nu,\rm on}$) with $a\sim10^{-3}$.

📊 关键结论与定量测量 (Quantitative Results)

EP251129a at $z=0.505$ exhibits an early internal X-ray plateau ($\alpha_{\rm X,1}=0.93$) terminating at $t_{\rm X,1}=1.27\times10^4$ s with a steep drop ($\alpha_{\rm X,2}=5.28$) and spectral softening ($\Gamma_{\rm X}$ softening from $\simeq2.1$ to $\simeq3.6$), followed by a second plateau ($\alpha_{\rm X,3}=0.42$) up to ~42 days in rest frame. Radio afterglow shows a steep rise ($F_\nu\propto t^{1.9-2.5}$, with $\alpha_{5.5,1}=-2.46$, $\alpha_{7.4,1}=-1.89$) peaking at rest-frame $\sim24-27$ days with $\nu L_\nu\simeq3\times10^{40}\,{\rm erg\,s^{-1}}$. Modeling yields fallback accretion energy injection $E_{\rm inj}\simeq6\times10^{51}$ erg and accreted mass $M_{\rm acc}\simeq0.2\,M_\odot$ onto the newly formed black hole.

🎯 建议精读切入点 (Reading Guide)

Inspect the multi-instrument X-ray and radio light curve in Figure 2 and review Section 4 for the two-component off-axis jet and spin-down collapse model.

PAPER 05 ⭐ 必读精选 🧭 综合前沿课题 (General Highlights) arXiv:2610.35539

[arXiv:2610.35539] An Intermediate Mass Stripped Star in a 4.2-hour Orbit around a Compact Object

An Intermediate Mass Stripped Star in a 4.2-hour Orbit around a Compact Object

作者：Bethany Ludwig, Maria R. Drout, Ylva Götberg, Alexander Laroche, Katie Auchettl 等 arXiv 原文 | PDF 直达 | SciX (ADS) | 站内交互雷达

RESEARCH HIGHLIGHT & BREAKTHROUGH / 核心突破与推荐理由

【核心突破】：Radial-velocity semi-amplitude $K = 151.4\ \mathrm{km\,s^{-1}}$ and ellipsoidal modulations establish a companion mass of $M_2 = 1.58\ M_\odot$ in a $4.2\ \mathrm{h}$ orbit, setting the stage for Case BB mass transfer, a Type Ibn supernova, and GW coalescence in $220\ \mathrm{Myr}$.

【推荐理由】：Identifies the crucial missing evolutionary link in double neutron star formation: an intermediate-mass stripped helium star orbiting a neutron star in a ultra-compact orbit.

🔍 科学问题与背景 (Problem)

Identifying the intermediate evolutionary stage of double neutron star progenitors—specifically verifying the existence of stripped helium stars in tight orbits with a first-born compact object prior to Case BB mass transfer and core collapse—and determining companion nature against main-sequence or double-stripped star alternatives.

🛠️ 观测与研究方法 (Methodology)

Multi-epoch Magellan/MagE optical spectroscopy analyzed with *The Joker* for orbital parameter estimation and radial velocity semi-amplitude K; NTT/ULTRACAM high-speed photometry fitted with `ellc` for ellipsoidal modulation and orbital inclination constraints; Chandra ACIS-S and XMM-Newton X-ray upper limits evaluated against Bondi-Hoyle-Littleton accretion models ($L_{\rm BHL} = \epsilon \frac{GM_{\rm NS}\dot{M}_{\rm acc}}{R_{\rm NS}}$); and detailed MESA forward binary evolution tracking non-conservative Case BB mass transfer ($dj/dt = \gamma j$) through core collapse and gravitational-wave orbital decay.

📐 物理模型与关键假设 (Models & Assumptions)

- The stripped star rotates synchronously with the orbital period due to rapid tidal synchronization in close binaries.

⚠️ 局限性与观测边界 (Limits & Caveats)

- DGL-3 is not significantly detected in individual or stacked XMM-Newton/Chandra observations, limiting the 2–$7\ \mathrm{keV}$ unabsorbed luminosity to $< 7.8 \times 10^{32}\ \mathrm{erg\ s^{-1}}$.




Fig. 3 [核心物理演化图]： Figure 3: Evolution from stripped star binary to merging double neutron star. Mass and orbital evolution of the stripped star binary model before (a) and after (b) the supernova. a, Top: masses of the stripped star, its remnant, and the companion. The surface is marked by a line; the interior is colored by the leftover hydrogen layer (yellow), the helium core (light blue) and the carbon-oxygen core (dark blue). Grey shading indicates the phase of Roche-lobe overflow referred to as case BB mass transfer and during which all hydrogen is lost. Bottom: orbital period evolution, with circular orbits assumed until core collapse. 1M ⊙ of material is lost in the helium-rich supernova, which makes the resulting double neutron star orbit eccentric. Gravitational-wave radiation then shrinks and circularizes the orbit, producing a neutron-star merger after $\sim 220\,\mathrm{Myr}$. b, Predicted future evolutionary sequence of the observed system. The stripped star fills its Roche lobe during helium-shell burning (case BB mass transfer), explodes as a stripped-envelope supernova, and leaves a bound neutron-star binary that merges through gravitational-wave emission.




Fig. 1 [数据拟合与模型对比]： Figure 1: The light curve and orbit of DGL-3. The light curve (a) and RV curve (b) are plotted as a function of orbital phase, folded so that conjunctions occur at phases 0 and 0.5. Normalized and detrended ULTRACAM photometry is shown as the grey scatter points and binned in 5-minute intervals as the dark pink points. MagE radial velocities are shown as the black points. Thick teal lines show the best-fit ellc model in (a) and the best-fit Joker model in (b). Thin teal curves show other models allowed by the data: ellc solutions within 2σ of the best fit in (a), and random draws from the Joker MCMC posterior in (b). The light curve shows double-peaked variability consistent with ellipsoidal modulation. The RV curve shows large-amplitude variations from the short-period binary. (c) and (d) show the residuals of the binned photometry and radial velocities relative to the best-fit models.

📊 关键结论与定量测量 (Quantitative Results)

The spectroscopic and photometric analysis shows the binary has orbital period $P \approx 4.2\,\mathrm{h}$, semi-amplitude $K = 151.4 \pm 2.5\,\mathrm{km\,s^{-1}}$, and circular orbit ($e < 0.045$ at 95% credible limit). The mass function $f(m) = \frac{PK^3}{2\pi G} = 0.062 \pm 0.003\,M_\odot$ yields a minimum companion mass $M_{2,\mathrm{min}} = 1.38 \pm 0.25\,M_\odot$ ($i = 90°$), with ellipsoidal light-curve modeling yielding a posterior peak of $M_2 = 1.58\,M_\odot$ (68% HDI: $[1.40 - 2.41]\,M_\odot$), identifying the companion as a compact object (most likely a neutron star). Chandra limits unabsorbed 2–7 keV X-ray luminosity to $L_{2-7\mathrm{keV}} < 7.8 \times 10^{32}\,\mathrm{erg\,s^{-1}}$. MESA evolutionary models show the system undergoes non-conservative Case BB mass transfer, exploding as an interaction-powered Type Ibn supernova and leaving a bound double neutron star that merges via gravitational waves in $\tau_{\mathrm{delay}} \approx 220\,\mathrm{Myr}$.

🎯 建议精读切入点 (Reading Guide)

Focus on the spectroscopic radial velocity fit with The Joker in Section 2 and the MESA forward binary evolution tracks in Section 4.

PAPER 06 ⭐ 必读精选 🧭 综合前沿课题 (General Highlights) arXiv:2610.35619

[arXiv:2610.35619] Indication of Electron Antineutrinos from the Diffuse Supernova Neutrino Background

Indication of Electron Antineutrinos from the Diffuse Supernova Neutrino Background

作者：The Super-Kamiokande Collaboration, :, K. Abe, Y. Asaoka, M. Harada 等 arXiv 原文 | PDF 直达 | SciX (ADS) | 站内交互雷达

RESEARCH HIGHLIGHT & BREAKTHROUGH / 核心突破与推荐理由

【核心突破】： Unbinned extended maximum likelihood fitting across pure-water and Gd-loaded SK phases yields a best-fit $\bar{\nu}_e$ flux of $3.6^{+1.6}_{-1.5}\ \mathrm{cm}^{-2}\,\mathrm{s}^{-1}$ for $E_\nu > 13.3$ MeV, providing direct empirical bounds on cosmic star formation and stellar core collapse.

【推荐理由】： Reports a milestone 2.6σ indication of the long-sought diffuse supernova neutrino background using over 5000 days of Super-Kamiokande observations.

🔍 科学问题与背景 (Problem)

Detecting electron antineutrinos from the cumulative past core-collapse supernovae (DSNB) against overwhelming backgrounds from atmospheric neutrino neutral-current and charged-current interactions as well as cosmic-ray muon spallation products.

🛠️ 观测与研究方法 (Methodology)

Analysis of 5003.0 days of Super-Kamiokande data (pure water and Gd-loaded SK-VI to SK-VIII phases) detecting inverse beta decay $\bar{\nu}_e + p \to e^+ + n$ via an unbinned extended maximum likelihood fit across 6 subregions defined by Cherenkov angle θ_C and tagged neutron multiplicity N_n, incorporating machine learning for NC background rejection and an optimized threshold of $E_{rec} = 11.5$ MeV for $N_n = 1$.

📐 物理模型与关键假设 (Models & Assumptions)

- DSNB signal follows core-collapse supernova neutrino emission models (such as Horiuchi+09 6 MeV, Kresse+21, or Ashida+23).

⚠️ 局限性与观测边界 (Limits & Caveats)

- Signal significance stands at 2.6σ, which represents an observational indication rather than a definitive 5σ discovery.




Fig. 1 [核心观测结果]： Figure 1: Energy spectra of the data and expected backgrounds in the $N_n = 1$ signal region. The grey hatch represents the total uncertainty on the expected backgrounds. Background-only p -values for each 2 MeV bin for $E_{rec} \in [11.5, 29.5]$ MeV are 0.12, 0.14, 0.77, 0.039, 0.24, 0.32, 0.46, 0.27, and 1.00. The nominal Horiuchi+09 DSNB model prediction [60] is overlaid for reference.




Fig. 5 [数据拟合与模型对比]： Figure 5: Profile likelihood ratio as a function of the DSNB flux for $E_\nu > 13.3$ MeV. The solid black curve represents the joint fit. The colored dashed curves represent the separate per-phase fits, shown only for comparison.

📊 关键结论与定量测量 (Quantitative Results)

Using a representative DSNB model (Horiuchi+09 with $T = 6\ \mathrm{MeV}$), the best-fit $\bar{\nu}_e$ flux is $3.6^{+1.6}_{-1.5}\ \mathrm{cm^{-2}\,s^{-1}}$ for $E_\nu > 13.3\ \mathrm{MeV}$ (corresponding to an event rate of $4.8^{+2.2}_{-2.0}\ \mathrm{events\,yr^{-1}}$), rejecting the null hypothesis at 2.6σ. For $E_\nu > 17.3\ \mathrm{MeV}$, the flux is $1.8^{+0.8}_{-0.7}\ \mathrm{cm^{-2}\,s^{-1}}$.

🎯 建议精读切入点 (Reading Guide)

Examine Section 3 for Cherenkov angle and neutron tagging selection cuts, and Figure 4 for the final reconstructed energy spectrum and likelihood contours.

PAPER 07 ⭐ 必读精选 🧭 综合前沿课题 (General Highlights) arXiv:2610.35675

[arXiv:2610.35675] The design and flight of the Payload for Ultrahigh Energy Observations (PUEO)

The design and flight of the Payload for Ultrahigh Energy Observations (PUEO)

作者：PUEO Collaboration, Q. Abarr, S. Agarwal, J. Alfaro, S. Ali 等 arXiv 原文 | PDF 直达 | SciX (ADS) | 站内交互雷达

RESEARCH HIGHLIGHT & BREAKTHROUGH / 核心突破与推荐理由

【核心突破】： A 96-horn dual-polarization array synthesized into 48 real-time coherent beams via 26 RFSoC FPGAs achieved 50% trigger efficiency at $\mathrm{SNR} = 1.4$, demonstrating an order-of-magnitude sensitivity leap over ANITA-IV.

【推荐理由】： Details the flight validation of an ultra-sensitive digital phased-array trigger architecture for discovering ultrahigh-energy Askaryan neutrinos above $1\ \mathrm{EeV}$.

🔍 科学问题与背景 (Problem)

Detecting extremely low flux ultrahigh-energy neutrinos ($E \gtrsim 1\ \mathrm{EeV}$) requires monitoring massive target volumes ($\sim 10^6\ \mathrm{km}^3$) of Antarctic ice while achieving trigger sensitivity near the thermal noise floor and discriminating genuine impulsively coherent Askaryan or air shower signals from anthropogenic RF interference and thermal fluctuations.

🛠️ 观测与研究方法 (Methodology)

PUEO incorporates a 96-horn dual-polarization main antenna array ($300 - 1200\ \mathrm{MHz}$) coupled to a real-time digital interferometric phased-array trigger implemented on 26 Xilinx RFSoC ZU47DR modules sampling at $3\ \mathrm{GSa/s}$, synthesizing 48 simultaneous coherent beams. Signal classification on the science computer uses coherent summation and Hilbert envelope energy concentration ($< 10\%$ envelope duration containing 30% total power for high-priority signals) alongside calibration pulses from ground pulsers and twin HiCal-3 balloons.

📐 物理模型与关键假设 (Models & Assumptions)

⚠️ 局限性与观测边界 (Limits & Caveats)

- Instrument duty cycle was restricted to an overall average of $\sim 55\%$ to maintain RFSoC FPGA temperatures below the $80^\circ\mathrm{C}$ threshold.




Fig. 1 [物理模型示意图]： Figure 1: Classes of particle-shower induced radio signatures observable by PUEO. UHECR showers can be observed through either the direct (left) or reflected (right) channels of geomagnetic and Askaryan emission in the air. Neutrinos can be detected via the channel of tau-decay on the left or in-ice Askaryan emission on the right. An inset diagram of the payload as flown is shown on the top right.




Fig. 40 [核心观测结果]： Figure 40: The coherently summed waveforms (top row) and their reconstructed pointing directions (bottom row) as seen by the PUEO MI for a HiCal-3a direct (a, c) and reflected (b, d) event pair. Time of flight and reconstruction identify this event pair as originating from the HiCal-3a pulser. The reconstruction and coherent sum are performed using preliminary estimates for the PUEO antenna phase centers from photogrammetry.

📊 关键结论与定量测量 (Quantitative Results)

PUEO successfully operated during a 23-day Antarctic flight with an average livetime fraction of $\sim 55\%$ (limited by thermal constraints to keep RFSoC temperatures below $80^{\circ}\mathrm{C}$), collecting $114,046,211$ RF trigger events. Ground-testing confirmed a 50% trigger efficiency threshold at $\mathrm{SNR} = 1.4$ (over twice as sensitive as ANITA-IV). The multi-antenna GNSS orientation achieved smoothed pitch and roll standard deviations of 0.015° and 0.024°, and heading residuals of 0.070° after applying a spin-dependent time-lag correction $\mathrm{Heading} = \mathrm{Heading} - \tau \cdot \omega$.

🎯 建议精读切入点 (Reading Guide)

Read Section 2 for the phased-array RFSoC trigger synthesis architecture and Section 5 for the Antarctic flight operational performance.

PAPER 08 ⭐ 必读精选 🧭 综合前沿课题 (General Highlights) arXiv:2610.37044

[arXiv:2610.37044] Discovery of a New Electron Spectrum Produced by Fast Cooling of Relativistic Electrons

Discovery of a New Electron Spectrum Produced by Fast Cooling of Relativistic Electrons

作者：Z. Lucas Uhm arXiv 原文 | PDF 直达 | SciX (ADS) | 站内交互雷达

RESEARCH HIGHLIGHT & BREAKTHROUGH / 核心突破与推荐理由

【核心突破】：Non-steady fast cooling transforms the sub-injection electron spectrum into an evolving smooth broken power law $f_{\rm U}(\gamma_e,t) = A(t)\gamma_e^{-2}[1+\gamma_0(t)/\gamma_e]^{-\beta}$, which invalidates the universal γ_e^{-2} steady-state approximation.

【推荐理由】：Overturns decades-old assumptions in relativistic jet astrophysics by deriving the exact analytical kinetic spectrum of fast-cooling electrons below γ_m.

🔍 科学问题与背景 (Problem)

In astrophysical relativistic jets such as GRBs and blazars, electrons below the minimum injection Lorentz factor γ_m evolve in a non-steady state, causing their energy distribution to deviate significantly from the canonical steady-state power law $f(\gamma_e) \propto \gamma_e^{-2}$.

🛠️ 观测与研究方法 (Methodology)

Analytical kinetic integration and differential equation solving of electron cooling trajectories $\dot{\gamma}_e = -g(t)\gamma_e^2 - \frac{2}{3t}\gamma_e$ combined with exact verification against the source-free Fokker-Planck equation $\frac{\partial f}{\partial t} + \frac{\partial}{\partial \gamma_e}[\dot{\gamma}_e f] = 0$ for $\gamma_e < \gamma_m$.

📐 物理模型与关键假设 (Models & Assumptions)

- Conical flow expanding radially with a constant bulk Lorentz factor for adiabatic cooling
- Continuous relativistic electron injection occurs only above γ_m with a power-law distribution $\propto \gamma_e^{-p}$

⚠️ 局限性与观测边界 (Limits & Caveats)




Fig. 1 [Physical Evolution]：Figure 1: Comparison of our electron spectrum $f_{\rm U}$ (thick-dotted lines) to a detailed numerical calculation of a blazar electron spectrum (solid lines) as a function of observer time $t_{\rm obs}$. The solid lines show the model [a] presented in [24]. The model [a] has a minimum injection Lorentz factor $\gamma_m = 10^3$.

📊 关键结论与定量测量 (Quantitative Results)

In the fast-cooling regime below the minimum injection Lorentz factor ($\gamma_e < \gamma_m$), the time-evolving cooled electron spectrum follows the smooth broken power-law form $f_{\rm U}(\gamma_e,t) = A(t)\gamma_e^{-2}[1+\gamma_0(t)/\gamma_e]^{-\beta}$, scaling asymptotically as γ_e^{-2} for $\gamma_e \gg \gamma_0(t)$ and $\gamma_e^{-2+\beta}$ for $\gamma_e \ll \gamma_0(t)$. For pure synchrotron cooling with $B \propto t^{-b}$ and injection rate $R_{\rm inj} \propto t^q$, $A(t) = \frac{b_0}{g_0}t^{2b+q}$, $\gamma_0(t) = \frac{2b-1}{g_0}t^{2b-1}$, and $\beta = \frac{2b+q}{2b-1}$; with adiabatic cooling included (expansion $\dot{\gamma}_{\rm ad} = -\frac{2}{3t}\gamma_e$), $\gamma_0(t) = \frac{6b-1}{3g_0}t^{2b-1}$ and $\beta = \frac{6b+3q+2}{6b-1}$.

🎯 建议精读切入点 (Reading Guide)

Review Section 2 for the kinetic integration of the cooling trajectories and the differential verification against the Fokker-Planck equation.

PAPER 09 ⭐ 必读精选 🧭 综合前沿课题 (General Highlights) arXiv:2610.37214

[arXiv:2610.37214] Coherent Radio Emission from Flux Rope Mergers in 3D Relativistic Reconnection

Coherent Radio Emission from Flux Rope Mergers in 3D Relativistic Reconnection

作者：Jasmine Parsons, Lorenzo Sironi, Anatoly Spitkovsky, Alexander Chernoglazov arXiv 原文 | PDF 直达 | SciX (ADS) | 站内交互雷达

RESEARCH HIGHLIGHT & BREAKTHROUGH / 核心突破与推荐理由

【核心突破】： Flux rope mergers in 3D pair-plasma reconnection sheets generate 100% linearly polarized X-mode waves with radiation efficiency $\eta \sim (1\text{–}2) \times 10^{-3}$, establishing reconnection as a self-consistent physical engine for FRBs.

【推荐理由】： Delivers the first conclusive 3D PIC proof that coherent fast magnetosonic waves survive realistic 3D kink instabilities in relativistic reconnection.

🔍 科学问题与背景 (Problem)

Whether 3D relativistic instabilities (e.g., relativistic drift kink instability and flux rope kink instability) and realistic outflow boundaries suppress coherent radio wave emission found in simplified 2D periodic relativistic reconnection simulations.

📐 物理模型与关键假设 (Models & Assumptions)

🛠️ 观测与研究方法 (Methodology)

3D particle-in-cell (PIC) simulations of pair plasmas using TRISTAN-MP with open outflow boundaries along x ($L_x = 1600\,c/\omega_p$, $L_z = 816\,c/\omega_p$), cold magnetization $\sigma = B_0^2/(4\pi n_0 mc^2) = 10$, and guide fields $B_g/B_0 \in \{0, 0.1, 0.3\}$, diagnosing wave Poynting flux $S_{\rm wave} = (c/4\pi)(\delta E_z \delta B_x - \delta E_x \delta B_z)$ and efficiency $\eta \equiv S_{\rm wave}/S_0$.

⚠️ 局限性与观测边界 (Limits & Caveats)




Fig. 4 [Physical Model Schematic]： Figure 4: Large-scale, coherent structures in the $B_g/B_0 = 0.3$ simulation at $ct/L = 6.75$. Top row: Volume rendering of the average particle outflow velocity $\langle v_x \rangle/c$. Purple and red field lines show the upstream field above and below the sheet, respectively, and the tubes on each side of the sheet follow the corresponding field direction. The left panels zoom in on the interface between a tube above and a tube below the sheet, with a sample of reconnected field lines overlaid. The green and gold field lines originate from two different diffusion regions (shown in black, i.e., regions of strong $E_\parallel$), and are pulled in opposite directions along x. Both of these bundles of field lines cross the sheet, with one end above it and one end below it. Bottom row: Same as above, but showing particle density. The linkage point between the two bundles appears in the left-hand zoom-in as a large overdense region. The diffusion region that produced the green bundle is indicated by a black-outlined arrow, while a newly created flux rope is indicated by a purple-outlined arrow. A green bar indicates that the density structure is $\sim 200\,c/\omega_p$ wide.




Fig. 5 [Physical Evolution]： Figure 5: Polarization angle (PA) of the outgoing waves, for $B_g/B_0 = 0$ (top row) and $B_g/B_0 = 0.3$ (bottom row), at $ct/L = 5.6$. The four plots show $x - z$ slices near the top of the box (left column) and bottom of the box (right column), $\sim 640\, c/\omega_p$ from the current sheet, showing the outgoing Poynting flux carried by the coherent waves, normalized by the inflowing flux S_0. The yellow-purple areas are patches of high-intensity waves. The light blue ticks denote the direction of the oscillating electric vector in these bright patches, whose angle from the z axis is the PA. In the $B_g/B_0 = 0$ case, the waves escaping on either side of the box have their electric field aligned with the z axis. For the $B_g/B_0 = 0.3$ case, on the other hand, because the PA of the outgoing waves is set by the direction of the upstream field, there is a rotation across the sheet by $\Delta\mathrm{PA} \sim 2\ \arctan(B_g/B_0)$.

📊 关键结论与定量测量 (Quantitative Results)

Coherent fast magnetosonic waves are stably excited via flux rope mergers in 3D relativistic reconnection, yielding a radiation efficiency of $\eta \sim 7 \times 10^{-4}$ for $B_g/B_0 = 0$ and $\eta \sim (1\text{–}2) \times 10^{-3}$ for moderate guide fields ($B_g/B_0 = 0.1\text{–}0.3$). The escaping radiation is 100% linearly polarized X-mode ($\delta\boldsymbol{E} \perp \boldsymbol{B}$), with a polarization angle offset between opposite sides of the sheet of $\Delta\mathrm{PA} \sim 2\arctan(B_g/B_0)$.

🎯 建议精读切入点 (Reading Guide)

Focus on Figure 2 for the 3D wave Poynting flux spatial structure and Section 3 for the efficiency scaling with guide field B_g/B_0.

PAPER 10 ⭐ 必读精选 🧭 综合前沿课题 (General Highlights) arXiv:2610.38673

[arXiv:2610.38673] Black Hole Spin Evolution in Magnetized Accretion Flows

Black Hole Spin Evolution in Magnetized Accretion Flows

作者：Danat Issa, Ore Gottlieb, Jonatan Jacquemin-Ide, Matthew Liska arXiv 原文 | PDF 直达 | SciX (ADS) | 站内交互雷达

RESEARCH HIGHLIGHT & BREAKTHROUGH / 核心突破与推荐理由

【核心突破】： Horizon electromagnetic torques cause black hole equilibrium spin to decrease monotonically with normalized flux $\tilde{\phi}$, tumbling to $a_{\star;\mathrm{eq}} \approx 0.05$–$0.06$ in MAD flows after accreting $\lesssim 50\%$ of their mass.

【推荐理由】： Presents the first continuous dynamical determination of black hole spin evolution across the entire magnetic flux spectrum using self-consistent 3D GRMHD.

🔍 科学问题与背景 (Problem)

Prior GRMHD simulations assumed static black hole spacetimes and explored only binary extremes (pure SANE or saturated MAD), lacking continuous quantitative descriptions of BH spin evolution and equilibrium spins across intermediate magnetic flux levels.

🛠️ 观测与研究方法 (Methodology)

Performed long-duration 3D GRMHD simulations with the H-AMR code tracking horizon energy and angular momentum fluxes ($dM_{\mathrm{BH}}/dt = \int T^r_t \, dA_{\theta\varphi}$, $dJ/dt = -\int T^r_\varphi dA_{\theta\varphi}$), updating the Kerr-Schild metric self-consistently with a moving boundary at $r_{\mathrm{b}} = 0.7 r_{\mathrm{H}}$. Numerical fits for hydrodynamical and electromagnetic fluxes yield the spin rate $s = l_{\mathrm{HD}} + l_{\mathrm{EM}} - 2a_\star(e_{\mathrm{HD}} + e_{\mathrm{EM}})$.

📐 物理模型与关键假设 (Models & Assumptions)

- Coordinate metric changes instantaneously over time steps without backreacting on external gas through non-local metric work terms ($|E_{\mathrm{miss}}|/\Delta M_{\mathrm{BH}} \lesssim 2\%$–$4\%$)

⚠️ 局限性与观测边界 (Limits & Caveats)

- Fitting formulae calibrated for $\tilde{\phi} \lesssim 1.2$ in radiatively inefficient thick disks; radiative cooling (e.g., in collapsars or thin disks) shifts $a_{\star;\mathrm{eq}}$ higher




Fig. 1 [Physical Evolution]： Figure 1: Equilibrium spin as a function of the MAD parameter. The circles are roots of $s(a_{\star;\mathrm{eq}}, \tilde{\phi}) = 0$, and the line marks the fit in Eq. (4).




Fig. 2 [Physical Evolution]： Figure 2: Final BH spin, $a_{\star;\mathrm{f}}$, as a function of (fixed) $\tilde{\phi}$ and of the ratio of the accreted rest mass, $M_{0,\mathrm{acc}}$, to the initial BH mass, shown for initial spins $a_{\star,0} = 0$ (a), $a_{\star,0} = 0.5$ (b), and $a_{\star,0} = 1$ (c).

📊 关键结论与定量测量 (Quantitative Results)

The equilibrium spin declines steeply with normalized horizon magnetic flux $\tilde{\phi} \equiv \phi/\phi_{\mathrm{MAD}}$, dropping from $a_{\star;\mathrm{eq}} \approx 0.9$ in weakly magnetized flows ($\tilde{\phi} \lesssim 0.2$), through $a_{\star;\mathrm{eq}}(\tilde{\phi} = 0.5) \approx 0.5$, down to $a_{\star;\mathrm{eq}} \approx 0.05$–$0.06$ at MAD saturation ($\tilde{\phi} = 1$). In the sustained MAD regime, the BH reaches within 3%–6% of its equilibrium spin by accreting only $\lesssim 50\%$ of its initial mass.

🎯 建议精读切入点 (Reading Guide)

Inspect Figure 3 and Section 3.2 for the equilibrium spin curve $a_{\star;\mathrm{eq}}(\tilde{\phi})$ and the hydrodynamic vs. electromagnetic angular momentum decomposition.

3. 📌 本周重点关注文献 (Worth-Knowing Selected Works · 共 76 篇)

STRUCTURED COMPENDIUM

以下文献针对特定物理机制或子方向提供重要观测线索与实测约束，采用前沿精选卡片与总览表格双轨呈现。

3.1 重点关注 · 特选导读 (Featured Highlights · 共 10 篇)

⭐ 重点关注 **[arXiv:2610.31754] Image Morphology of Misaligned Magnetically Arrested Disks**

原题： Image Morphology of Misaligned Magnetically Arrested Disks | **作者：** Samuel Feyan, Cora Prather, Dominic O. Chang 等 | PDF

【推荐理由】： Quantifies the systematic biases induced in EHT black hole spin parameter estimation when assuming axisymmetric disk-jet geometry for tilted MADs.

【核心要点】： While high-spin prograde MADs align naturally near the horizon via jet torques, intermediate and retrograde tilted systems maintain asymmetric warps that bias spin inferences by over 50% under standard geometric templates.

【阅读切入】： Check Figure 4 for the synthetic 230 GHz ray-traced images across tilt angles and Section 4 for parameter estimation discrepancies.

⭐ 重点关注 **[arXiv:2610.31755] From Inspiral to Outburst: Modeling a Potential Luminous Red Nova from Betelgeuse**

原题： From Inspiral to Outburst: Modeling a Potential Luminous Red Nova from Betelgeuse | **作者：** Naunet Leonhardes-Barboza, Andrea Antoni, Morgan MacLeod 等 | PDF

【推荐理由】： Provides an end-to-end self-consistent 1D radiation-hydrodynamic simulation linking common-envelope companion inspiral to luminous red nova shock breakouts.

【核心要点】： Tidal disruption and orbital energy deposition of a 0.7–$2.0\ M_{\odot}$ companion inside a red supergiant launches a shock breakout yielding peak optical luminosities of $(2.4$–$6.2) \times 10^{41}\ \mathrm{erg\,s^{-1}}$ scaling as $L_{\mathrm{peak}} \propto M_{\mathrm{comp}}^{0.93}$.

【阅读切入】： Review Section 2 for the energy deposition coupling in Athena++ and Figure 3 for the resulting radiation diffusion light curves.

⭐ 重点关注 **[arXiv:2610.31990] Tilt, warp, and dissipation flare in hydrodynamic disks around kicked compact objects**

原题： Tilt, warp, and dissipation flare in hydrodynamic disks around kicked compact objects | **作者：** Yoonsoo Kim | PDF

【推荐理由】： Presents 3D hydrodynamic simulations of how impulsive natal kicks distort bound accretion disks, producing transient dissipation flares and retrograde disk warps.

【核心要点】： Impulsive compact object kicks induce hydrodynamic self-collisions within $r_0 \equiv GM/V_{\mathrm{kick}}^2$ that dissipate 5–30% of the bound kinetic energy, driving an electromagnetic flare and inducing large-angle or retrograde tilts.

【阅读切入】： Examine Section 2 for analytical bound boundaries and Figure 5 for AthenaK 3D disk morphology and energy dissipation profiles.

⭐ 重点关注 **[arXiv:2610.32028] Magnetic field effect on neutral color superconductivity as a mechanism to boost the inner magnetic field of magnetars**

原题： Magnetic field effect on neutral color superconductivity as a mechanism to boost the inner magnetic field of magnetars | **作者：** Bo Feng | PDF

【推荐理由】： Formulates an alternative microscopic origin for magnetar interior fields based on charged-gluon chromomagnetic instability in color-superconducting quark cores.

【核心要点】： In the neutral 2SC phase, Fermi surface mismatch triggers tachyon gluon Meissner masses and paramagnetic anti-screening, naturally boosting internal magnetic fields to $\tilde{B} \sim 10^{17}\ \mathrm{G}$ from modest initial seed fields.

【阅读切入】： Refer to Section 3 for the derivation of the rotated electromagnetic gauge field and the gluon-vortex energy minimization condition.

⭐ 重点关注 **[arXiv:2610.32047] A Unified Model for Radioactive Signatures and Source Identifiability in Neutron Star Mergers**

原题： A Unified Model for Radioactive Signatures and Source Identifiability in Neutron Star Mergers | **作者：** Hisham Anwer | PDF

【推荐理由】： Establishes a rigorous statistical and radiative transfer framework demonstrating the severe degeneracy in identifying specific isotopes (e.g., $^{254}\mathrm{Cf}$) from kilonova light curves.

【核心要点】： Thermalization delays and multi-layer diffusion smooth radioactive decay pulses such that models with and without $^{254}\mathrm{Cf}$ achieve $< 9\%$ statistical discrimination power under realistic multi-band photometric uncertainties.

【阅读切入】： See Section 2 for the causal energy deposition operator and Section 4 for the Neyman-Pearson identifiability bounds applied to AT 2023vfi.

⭐ 重点关注 **[arXiv:2610.32168] SMMAN Studies of AGN Variability. I. Long-term Spectral Evolution and a Candidate ~1000-day Quasi-periodic Modulation in 4C+72.28**

原题： SMMAN Studies of AGN Variability. I. Long-term Spectral Evolution and a Candidate ~1000-day Quasi-periodic Modulation in 4C+72.28 | **作者：** Xin Wang, Timur Mufakharov, Guiping Tan 等 | PDF

【推荐理由】： Applies rigorous red-noise Monte Carlo simulations to radio monitoring of the blazar 4C+72.28, isolating a candidate ~ 1000 d quasi-periodic modulation.

【核心要点】： A 1026.7 d periodic flux modulation with 5.45% fractional amplitude and zero inter-band lag is uncovered, naturally accounted for by a slight geometric Doppler factor variation ($\delta_{\mathrm{max}}/\delta_{\mathrm{min}} \simeq 1.05$) in a precessing jet.

【阅读切入】： Review Section 3 for the broken power-law red-noise significance assessment and Section 5 for the helical jet precession interpretation.

⭐ 重点关注 **[arXiv:2610.32316] Radio Galaxies as UHECR Sources in Light of Recent Microquasar Results**

原题： Radio Galaxies as UHECR Sources in Light of Recent Microquasar Results | **作者：** A. M. Taylor, J. H. Matthews | PDF

【推荐理由】： Re-evaluates radio galaxies as ultrahigh-energy cosmic ray sources using microquasar-calibrated near-Bohm shear acceleration efficiency constraints.

【核心要点】： Microquasar SS 433 synchrotron cutoffs confirm near-Bohm electron acceleration, implying that radio galaxy jet shear layers can accelerate UHECR protons to > 10 EeV during episodic bursts like Cen A's event ~ 30 Myr ago.

【阅读切入】： Inspect Section 2 for the generalized Hillas luminosity criterion and Section 3 for jet shear acceleration timescale comparisons.

⭐ 重点关注 **[arXiv:2610.32323] A Framework for Analysing Nebular-Phase JWST Supernova Spectra with Application to Type Ic SN 2024aecx**

原题： A Framework for Analysing Nebular-Phase JWST Supernova Spectra with Application to Type Ic SN 2024aecx | **作者：** S. Barmentloo, A. Jerkstrand | PDF

【推荐理由】： Establishes a systematic mid-infrared NLTE/LTE line analysis framework for JWST nebular spectroscopy, applied to stripped-envelope core-collapse SN 2024aecx.

【核心要点】： Self-consistent analysis of [Ni II] and [Co II] lines reveals a low iron-zone filling factor ($f_{\rm FeHe} = 0.059$), proving that radioactive ^{56}Ni bubble expansion was negligible, while chlorine yields favor an O-C shell merger.

【阅读切入】： Read Section 2 for the critical density de-excitation formula and Section 4 for the parameter inversion of the JWST MIRI/NIRSpec spectrum.

⭐ 重点关注 **[arXiv:2610.32430] Local Analogs of Little Red Dots: Clues to the X-ray Weakness**

原题： Local Analogs of Little Red Dots: Clues to the X-ray Weakness | **作者：** Shi-Jiang Chen, Jun-Xian Wang, Zhen-Ya Zheng 等 | PDF

【推荐理由】： Uses targeted Einstein Probe FXT observations of local Little Red Dot analogs to resolve the mechanism behind their puzzling X-ray weakness.

【核心要点】： Stacking detection of local LRD analogs at $\log L_{2-10} = 42.55\ {\rm erg\,s^{-1}}$ reveals a relatively hard X-ray spectrum, ruling out unshielded super-Eddington accretion and Compton-thick obscuration in favor of mild intrinsic coronal weakness with Compton-thin absorption.

【阅读切入】： Examine Section 3 for the EP-FXT image stacking methodology and Figure 3 for the optical-to-X-ray spectral slope comparison.

⭐ 重点关注 **[arXiv:2610.32786] Probing the star-forming environment of GRB across cosmic ages with high-resolution X-ray spectroscopy: From XMM-Newton/RGS and XRISM/Resolve to NewAthena/X-IFU**

原题： Probing the star-forming environment of GRB across cosmic ages with high-resolution X-ray spectroscopy: From XMM-Newton/RGS and XRISM/Resolve to NewAthena/X-IFU | **作者：** Luigi Piro, Aishwarya Linesh Thakur, Alfredo Luminari 等 | PDF

【推荐理由】： Demonstrates the power of time-dependent non-equilibrium photoionization models for extracting circumburst abundances and Pop III signatures via future microcalorimeters.

【核心要点】： Non-equilibrium ionization modeling with TEPID reveals that XRISM and NewAthena will measure host gas densities to 0.05 dex and distinguish Pop III from Pop II progenitors via narrow Fe-K absorption line profiles (EW ~ 4–10 eV).

【阅读切入】： Check Section 2 for the updated TEPID code atomic rate implementation and Figure 6 for mock NewAthena X-IFU absorption line spectra.

3.2 重点关注 · 全周论文简表 (Weekly Compendium · 共 66 篇)

arXiv ID / 日期	文献题名与作者	核心要点与观测约束	阅读切入
arXiv:2610.40609 2026-10-29	**From Dispersion to Metallicity: Probing the Milky Way Circumgalactic Medium with Fast Radio Bursts** *From Dispersion to Metallicity: Probing the Milky Way Circumgalactic Medium with Fast Radio Bursts* Manami Roy, Sanskriti Das, Smita Mathur 等	Using 25 localized FRB sightlines at $z < 0.3$, the bootstrap median metallicity of the Milky Way CGM is inferred to be $Z_{\rm MW} = 0.12^{+0.09}_{-0.04}\ Z_\odot$ for diffuse baryon fraction $f_{\rm d} = 0.0$, $Z_{\rm MW} = 0.26^{+0.18}_{-0.12}\ Z_\odot$ for $f_{\rm d} = 0.6$, and $Z_{\rm MW} = 0.39^{+0.26}_{-0.20}\ Z_\odot$ for $f_{\rm d} = 0.8$. Even at the conservative limit $f_{\rm d} = 0.0$, the halo metallicity exceeds pristine IGM accretion ($Z_{\rm IGM} \lesssim 0.1\ Z_\odot$), whereas $f_{\rm d} = 0.8$ produces unphysical super-solar metallicities ($Z \gg 1\ Z_\odot$) along multiple sightlines.	Result
arXiv:2610.40024 2026-10-29	**Inverse Gertsenshteĭn-Zel'dovich Effect in Time-Independent Electromagnetic Field** *Inverse Gertsenshteĭn-Zel'dovich Effect in Time-Independent Electromagnetic Field* Rajendra Prasad Bhatt, Rudraksh Verma, R. Prasad 等	For GR tensor modes traversing a neutron star magnetosphere of radius $R_{\rm LC} = c/\Omega$, the total conversion factor scales as $\alpha^{+\times}_{\rm tot} = \frac{4G(B_y^{(0)})^2 R_{\rm LC}^2}{3c^4} \propto (B_y^{(0)})^2 R_{\rm LC}^2$, generating an induced spectral flux density $\langle S \rangle_{\rm tot}/f_g \sim 0.001 - 1000$ Jy consistent with observed FRB peak flux densities. The induced wave flux scales with the interaction length squared, $(R_1 + R_2)^2$, and yields completely polarized electromagnetic emission (degree of polarization $\Pi = 1$) reflecting the incoming gravitational wave polarization.	Applications
arXiv:2610.38956 2026-10-29	**Probing Scalar Polarization with Next-Generation Gravitational-Wave Detectors: The Roles of Detector Geometry and Source Inclination** *Probing Scalar Polarization with Next-Generation Gravitational-Wave Detectors: The Roles of Detector Geometry and Source Inclination* Sk Md Adil Imam, Macarena Lagos	Next-generation detectors ($2 \times$ CE and ETΔ) improve constraints on the breathing scalar amplitude α_B by a factor of ~ 40--$85\times$ over the O5 HLV network. Due to the intrinsic antenna pattern $Y_b(\iota) = \sqrt{3/2}\,\sin^2\iota$, the constraint width scales as $\sigma(\alpha_B) \propto 1/\sin^2\iota$; incorporating electromagnetic jet inclination priors tightens α_B constraints by $\approx 56\times$ (reducing the 68% uncertainty from 2.3×10^{-2} to 4.0×10^{-4}). The triangular geometry of ETΔ provides consistently superior separation of scalar polarization, yielding $\sim 2\times$ tighter bounds on α_B ($\pm 1.9 \times 10^{-4}$) compared to CE ($\pm 3.9 \times 10^{-4}$) for GW170817.	Results

arXiv ID / 日期	文献题名与作者	核心要点与观测约束	阅读切入
arXiv:2610.38848 2026-10-29	**SuperCrude EoS: A Simple, Analytical, and Reasonably Accurate Equation of State for Astrophysical Fluid Dynamics** *SuperCrude EoS: A Simple, Analytical, and Reasonably Accurate Equation of State for Astrophysical Fluid Dynamics* Paul C. Duffell, Dillon Hasenour	SuperCrude models the fluid via decomposed components $P = P_{\rm deg} + P_{\rm gas} + P_{\rm rad} + P_{\rm pairs}$ and $\mathcal{E} = \mathcal{E}_{\rm deg} + \mathcal{E}_{\rm gas} + \mathcal{E}_{\rm rad} + \mathcal{E}_{\rm pairs}$, recovering sound speed $c_s = \sqrt{(B_{\rm deg} + B_{\rm Th})/(\rho h)}$ with relativistic enthalpy $h = 1 + (P + \mathcal{E})/(\rho c^2)$. In hydrodynamic closures $P(\rho, \mathcal{E})$, errors are $< 5\%$ (mostly $< 10^{-3}$) relative to helmeos; the peak error in $\Gamma_H \equiv 1 + P/\mathcal{E}$ is $1.3\% - 2.5\%$ and Γ_1 error is $\lesssim 5\%$. In 1D hydro tests, density L_1 error is 3.64×10^{-3}, and single-zone execution time decreases from $571\,\mathrm{ns}$ (helmeos) to $89.7\,\mathrm{ns}$.	2. METHOD
arXiv:2610.38766 2026-10-29	**Unifying stochastic and deterministic signals: a practical search for the intermittent binary black-hole background** *Unifying stochastic and deterministic signals: a practical search for the intermittent binary black-hole background* Tore Boybeyi, Vuk Mandic, Joseph Romano	Applying the search to simulated O4 LIGO data allows combining unresolved subthreshold events with resolved catalogs, successfully pinning down the BBH merger peak redshift to $z_p = 1.77^{+0.65}_{-0.74}$ with local rate $R_0 = 19.1\,\mathrm{Gpc^{-3}\,yr^{-1}}$. Phase-randomization tests demonstrate that the background evidence completely vanishes (cumulative log Bayes factor dropping from $\ln \mathcal{B}_{\rm U} \approx 1.3 \times 10^3$ to -1.3) without inter-frequency bin correlations, showing single-bin cross-correlations fail to capture intermittent chirp backgrounds.	II. CALCULATING THE MARGINALIZED LIKELIHOOD RATIO
arXiv:2610.38765 2026-10-29	**Constructing likelihoods for a non-Gaussian gravitational-wave background** *Constructing likelihoods for a non-Gaussian gravitational-wave background* Tore Boybeyi, Vuk Mandic, Joseph Romano	The exact non-Gaussian GWB likelihood is decoupled via Hubbard-Stratonovich transformation into the characteristic function of single mergers $z(\eta; \Lambda)$. For a toy model with segment noise σ^2, event rate $\bar{N}$, and amplitude variance A_0^2, the exact detection floor flattens at $\bar{N} A_0^2 \simeq 10^{-2}$ for $\bar{N} \gtrsim 10^{-2}$ and drops by a factor of ~ 3 toward $\bar{N} = 10^{-3}$. The single-component likelihood $\mathcal{L}_{M=1}$ is sensitive only to the degenerate combination $\bar{N} A_0^2$, whereas multi-component Gaussian mixtures $\mathcal{L}_M$ converge to the exact detection threshold with only 3 components.	Constructing likelihoods for a non-Gaussian gravitational-wave background
arXiv:2610.37646 2026-10-27	**Measuring the magnetization profiles of galaxy groups and clusters** *Measuring the magnetization profiles of galaxy groups and clusters* Kritti Sharma, Vikram Ravi, Elisabeth Krause	For a coherence length $l_c = 10\,\mathrm{kpc}$ within $R < R_{200c}$, the 95% upper limits on the line-of-sight rms magnetic field are $B_{\parallel,\rm rms} < 3.9$–$11.6\,\mu\mathrm{G}$ for DESI BGS groups and < 2.3–$6.4\,\mu\mathrm{G}$ for RASS clusters, scaling as $l_c^{-1/2}$; the 95% lower limits on plasma beta are $\beta > 0.31$ and $\beta > 11$ respectively, derived via $B_{\parallel,\rm rms}(R) = B_{\parallel,\rm eff}(R)\sqrt{L(R)/l_c}$ with $B_{\parallel,\rm eff}(R) = \mathrm{RM_{rest}}(R)/[0.812\,\mathrm{DM}(R)]\,\mu\mathrm{G}$.	III.3 From RM 2 and tau to $Bparallel$
arXiv:2610.37574 2026-10-27	**Effects of Dust in Electron-Capture Supernovae: Difficulty in Explaining Intermediate Luminosity Red Transients** *Effects of Dust in Electron-Capture Supernovae: Difficulty in Explaining Intermediate Luminosity Red Transients* Yuma Takeichi, Daichi Tsuna, Daichi Hiramatsu 等	Standard ECSN models with dusty winds cannot simultaneously reproduce the light curves and colors of ILRTs: the intrinsic drop in input luminosity after the plateau causes $r_{\rm sub} \propto L_{\rm input}^{1/2}$ to shrink, triggering a runaway increase in optical depth τ_V and causing the optical luminosity to plummet by orders of magnitude; attempting to flatten the decline using dense CSM shock interaction ($L_{s,\rm rad} \propto t^{-0.3}$) requires $\dot{M}_{\rm dense} \simeq 9.7 \times 10^{-3}\,M_\odot\,\mathrm{yr^{-1}}$ and produces excessively red post-plateau colors ($B - V > 4$) that contradict observed ILRT colors ($B - V \lesssim 1.2$–1.6).	4.1. Comparison to ILRT light curves
arXiv:2610.37507 2026-10-27	**Faint and Fast Transients from Double Detonations of Low Mass White Dwarfs** *Faint and Fast Transients from Double Detonations of Low Mass White Dwarfs* Miranda Pikus, Abigail Polin, Dillon Hasenour 等	Low-mass WD double detonations ($M_{\rm tot} \sim 0.525$–$0.6\,M_\odot$, $M_{\rm shell} \sim 0.15$–$0.3\,M_\odot$) produce low ^{56}Ni yields (core produces $\lesssim 5 \times 10^{-3}\,M_\odot$ for thin shells, up to $\sim 10^{-2}\,M_\odot$), peak luminosities $L_{\rm peak} \sim 1.4 \times 10^{41}\,\mathrm{erg\,s^{-1}}$ ($M_{\rm peak} \sim -15$ to $-17\,\mathrm{mag}$), rapid rise times (5–10 days), and fast post-peak declines ($\Delta m_{10} \sim 1$–$1.75\,\mathrm{mag}$ in 10 days). Ejecta retain up to 28% unburned carbon with low expansion velocities ($v_{\rm Ca} \sim 5300$–$13300\,\mathrm{km\,s^{-1}}$).	3. RESULTS
arXiv:2610.37464 2026-10-27	**How common are extreme scattering events?: First large sample from the ASKAP VAST survey** *How common are extreme scattering events?: First large sample from the ASKAP VAST survey* Akash Anumarlapudi, Dylan L. Jow, Artem Tuntsov 等	The authors identify 236 ESE candidates (72 gold, 76 silver, 88 bronze) with flux variations ≥ 2. Under a characteristic scale $a \approx 1\,\mathrm{AU}$ and scale height $\approx 1\,\mathrm{kpc}$, the inferred core electron density is $n_0 \gtrsim 100\,\mathrm{cm^{-3}}$, implying an overpressured plasma lens with $P/k > 10^2 T_e\,\mathrm{K\,cm^{-3}}$. The measured ESE event rates at $888\,\mathrm{MHz}$ over a $12,500\,\mathrm{deg^2}$ footprint are $0.003\ (0.006)\ \mathrm{events/source/year}$ and $0.006\ (0.012)\ \mathrm{events/year/deg^2}$ for the gold (gold+silver) samples, with light curves strongly favoring overdense lenses ($\alpha > 0$).	IV Results

arXiv ID / 日期	文献题名与作者	核心要点与观测约束	阅读切入
arXiv:2610.37414 2026-10-27	**Evidence for a nova super-remnant surrounding the recurrent nova V745 Scorpii using eROSITA** *Evidence for a nova super-remnant surrounding the recurrent nova V745 Scorpii using eROSITA* Michael W. Healy-Kalesh, Sara Saeedi, Manami Sasaki 等	eROSITA reveals an elliptical shell with angular size $\sim 19' \times 10'$ (physical dimensions $\sim 90 \times 50$ pc at $d \sim 7.8$–8.2 kpc) with peak emission in the 0.5–1.2 keV range. Annular photometry in the $16'$–$24'$ shell region yields an unabsorbed X-ray flux of $\sim 2 \times 10^{-12}$ erg s^{-1} cm^{-2} and net luminosity $L_{\rm X} \sim 1.6 \times 10^{34}$ erg s^{-1}, nested inside a larger infrared dust cavity peaking at $\sim 46'$ (~ 100 pc).	5.2 X-ray modelling
arXiv:2610.37394 2026-10-27	**Probing the inner disk edge of rotating neutron stars with dipole excitations in finite nuclei** *Probing the inner disk edge of rotating neutron stars with dipole excitations in finite nuclei* P. S. Koliogiannis, N. Paar	Nuclear electric dipole polarizability constraints (CNSP-4 set) reduce the spread in the surface–ISCO transition gravitational mass $M_{\rm tr}(f)$ and equatorial radius $R_{\rm eq}$ across spin frequencies f by factors of 7--10 (85--90% reduction), bringing mass uncertainty below $0.025\,M_\odot$ and radius uncertainty below 0.15 km. Along the transition, the disk inner-edge Keplerian orbital frequency $\nu_{\rm in} = \Omega_{\rm orb}(R_{\rm in})/(2\pi)$ is confined to a band narrower than 20 Hz (down from 38--134 Hz across the full DD–PC family). For the 414 Hz pulsar 4U 0614+09 with highest kHz QPO at $\nu_{\rm QPO} \geq 1267$ Hz, the allowed mass range is narrowed from 0.81--$1.97\,M_\odot$ to $0.91 \lesssim M/M_\odot \lesssim 1.97$, and the lower-mass boundary equatorial radius is constrained to 12.40--12.71 km.	3 Results and discussion
arXiv:2610.37360 2026-10-27	**Impact of Neutrino Trapping on the Equation of State and Structure of Hybrid Stars with 2SC Quark Matter** *Impact of Neutrino Trapping on the Equation of State and Structure of Hybrid Stars with 2SC Quark Matter* Andrea Sabatucci, Armen Sedrakian	Neutrino trapping ($Y_{Le} = 0.4$) robustly shifts the hadron-quark deconfinement transition toward higher baryon densities compared to neutrino-transparent matter. Trapping raises the proton fraction and diminishes the isospin asymmetry $\delta = 1 - 2Y_p$, which sharply reduces the equation of state's sensitivity to the hadronic symmetry energy slope ($L_{\rm sym} \in [30, 70]$ MeV) because symmetry-energy energy density scales as $\propto E_{\rm sym}(\varrho)\,\delta^2$. For the reference DDLS50-NJL model ($\eta_D = \eta_V = 1$), maximum gravitational masses are $M_{\rm max}^{(S=1)} = 2.36\,M_\odot$ and $M_{\rm max}^{(S=2)} = 2.37\,M_\odot$; varying the vector coupling $\eta_V = G_V/G_S \in [0.8, 1.2]$ alters $M_{\rm max}$ by $\pm 1.3\%$, while strong diquark pairing ($\eta_D = 1.25$) reduces $M_{\rm max}$ by up to -8.5%. While Gibbs (global lepton conservation) and Maxwell (local lepton conservation) phase constructions produce nearly indistinguishable macroscopical $M - R$ curves, Gibbs produces a vastly more extended geometric radial mixed-phase core.	VI Summary and Conclusions
arXiv:2610.37308 2026-10-27	**Nonradial oscillation modes of axion condensed neutron stars in the Cowling approximation** *Nonradial oscillation modes of axion condensed neutron stars in the Cowling approximation* Antonio Gómez-Bañón, Pantelis Pnigouras, José A. Pons	The equilibrium background modification shifts f- and p-mode frequencies upward as $\nu_a \equiv m_a/(2\pi)$ increases (for instance, reaching $\nu_f = 2.87\,\mathrm{kHz}$, $\nu_{p_1} = 8.19\,\mathrm{kHz}$, and $\nu_{p_2} = 12.11\,\mathrm{kHz}$ at $\nu_a = 15\,\mathrm{kHz}$), while dynamical fluid-axion coupling partially counteracts this shift. When the mode frequency exceeds the threshold ν_a, outgoing axion emission damps the fluid modes with damping rate $\lvert\mathrm{Im}\,\omega\rvert \simeq L_a/(2E_{\rm mode})$, reaching damping timescales $\tau_a = 1/(2\pi\lvert\mathrm{Im}\,\nu\rvert) \sim 10\,\mathrm{ms}$. In addition, gravitational atom bound states exhibit hydrogenic eigenvalues $\tilde{k}_n^2 \equiv k_n^2/m_a^2 = -\alpha^2/n^2$ with $\omega_n \simeq m_a[1 - \alpha^2/(2n^2)]$ and undergo avoided crossings with fluid p-modes accompanied by an inversion of the axion energy fraction $x_a = E_a/(E_a + E_{\rm fluid})$.	III Results
arXiv:2610.37303 2026-10-27	**$γ$-ray variability of changing-look blazars: FSRQ-like jet behaviour** *$γ$-ray variability of changing-look blazars: FSRQ-like jet behaviour* Zahoor Malik, B. van Soelen, Sikandar Akbar 等	Changing-look blazars (CLBs) exhibit fractional variability $F_{\rm var} = 0.483$ at a 30-day cadence that is statistically indistinguishable from FSRQs (0.437, $p = 0.94$) but significantly higher than BL Lacs (0.262, $p = 1.4 \times 10^{-4}$). CLBs have Eddington-scaled accretion disk luminosities $\log(L_{\rm disk}/L_{\rm Edd}) = -1.42$, which are close to FSRQs (-0.85) and far above BL Lacs (-3.36), confirming that the optical changing-look behavior is driven by jet synchrotron continuum swamping rather than accretion mode transitions.	10 Conclusions

arXiv ID / 日期	文献题名与作者	核心要点与观测约束	阅读切入
arXiv:2610.37277 2026-10-27	**An SRG/eROSITA Study of Be X-ray binaries in Quiescence** *An SRG/eROSITA Study of Be X-ray binaries in Quiescence* Ajith Balu, Kinjal Roy, Biswajit Paul	Detections in the $0.5-10\,\mathrm{keV}$ range reveal quiescent X-ray luminosities spanning $L_{\mathrm{X}}\sim10^{33}-10^{36}\,\mathrm{erg\,s^{-1}}$, with photon indices $\Gamma\sim0.5-2.0$ and blackbody temperatures $kT_{\mathrm{bb}}\sim0.8-1.6\,\mathrm{keV}$ from emitting radii $R\sim0.1-2\,\mathrm{km}$. A power-law correlation connects X-ray luminosity and emission radius as $L\propto R^{\alpha}$ with index $\alpha=2.3\pm0.2$. In addition, upper limits on non-detected sources (e.g. 2S 1553−542 and RX J0812.4−3114) fall below the theoretical propeller luminosity $L_{\mathrm{prop}}\simeq4\times10^{37}k^{7/2}B_{12}^{2}P^{-7/3}M_{1.4}^{-2/3}R_{6}^{5}\,\mathrm{erg\,s^{-1}}$, confirming propeller or cold disc inhibition.	IV.2 Accretion models for BeXRBs in quiescence
arXiv:2610.37198 2026-10-27	**Searches for new particle physics by methods of classical and neutrino astronomy** *Searches for new particle physics by methods of classical and neutrino astronomy* Sergey Troitsky	The review synthesizes constraints on new physics across stellar laboratories, transient photon propagation, and neutrino telescopes. For axion-electron coupling from globular cluster TRGB calibrations, the 95% CL bound reaches $g_{ae}<6.4\times10^{-14}$ via electron bremsstrahlung energy loss $\varepsilon_a\propto g_{ae}^2T^4$. For high-energy photon propagation from GRB 221009A ($z\simeq0.151$), photon-ALP mixing ($m_a\simeq5.2\times10^{-7}\,\mathrm{eV}$, $g_{a\gamma}\simeq6\times10^{-11}\,\mathrm{GeV^{-1}}$) and subluminal quadratic LIV ($E_{\mathrm{LIV},2}\sim4\times10^{12}\,\mathrm{GeV}$) are tested against LHAASO ($>10\,\mathrm{TeV}$) and Carpet-3 ($E_\gamma=300^{+43}_{-38}\,\mathrm{TeV}$) observations, with an allowed host magnetic field window $1.3\,g_{11}^{-1}\lesssim B/\mu\mathrm{G}\lesssim4.2\,g_{11}$. For neutrino astronomy, superheavy dark matter ($m_X\gg10^2\,\mathrm{TeV}$) is bounded to lifetimes $\tau_X\gtrsim10^{26}$–$10^{28}\,\mathrm{s}$, and neutrino flavor ratios evolve from source states (e.g., standard $(1:2:0)_S$ or muon-damped $(0:1:0)_S$) toward $(1:1:1)_\oplus$ across cosmic baselines.	III High-energy photon propagation as a probe of new physics
arXiv:2610.37114 2026-10-27	**X-ray Emission Study of TeV Blazars** *X-ray Emission Study of TeV Blazars* Zhihao Ouyang, Hubing Xiao, Shangchun Xie 等	Luminosity correlations show $\log L_{\mathrm{TeV}}$ correlates positively with soft X-ray luminosity ($k=1.54\pm0.20$, $r=0.63$) and hard X-ray luminosity ($k=1.23\pm0.28$, $r=0.52$), consistent with leptonic scenarios. X-ray luminosity correlates strongly with disk luminosity $\log L_{\mathrm{disk}}$ ($r=0.61$ soft, $r=0.77$ hard) but lacks significant correlation with black hole mass $\log M_{\mathrm{BH}}$ ($p=0.21$ and 0.77). The majority of well-sampled sources exhibit log-normal soft X-ray flux distributions with PSD slopes $\beta_{\mathrm{PSD}}\sim1.0$–$1.6$.	IV Discussion
arXiv:2610.37052 2026-10-27	**Self lensing in BH-NS binaries as progenitors for periodically repeating Fast Radio Bursts** *Self lensing in BH-NS binaries as progenitors for periodically repeating Fast Radio Bursts* L. E. Rodríguez, M. Cruces, G. Voisin 等	Pure Keplerian self-lensing fails with a tiny activity window $\tau_{\mathrm{aw}}^{\mathrm{KP}}\sim\mathcal{O}(0.5\,\mathrm{hours})$, whereas Lense-Thirring precession with period $\tau_{\mathrm{LT}}=\frac{\pi(1-e^2)^{3/2}c^3a^3}{G^2M_{\mathrm{BH}}^2\chi}=2\tau_{\mathrm{FRB}}$ reproduces wide activity windows $\tau_{\mathrm{aw}}^{\mathrm{LT}}\sim0.5\tau_{\mathrm{FRB}}$ for small inclinations $i\sim(5$–$8)\times10^{-2}\,^\circ$. Splitting time delays $\tau_{\mathrm{lens}}^{\mathrm{Spl}}\approx\frac{4GM_{\mathrm{BH}}}{c^3}(1+z)\lambda(r_s/a)^{1/2}\sim10^{-3}$–$10^{-2}\,\mathrm{s}$ match the observed millisecond waiting-time peak in FRB 20121102A, constraining the companion mass to $M_{\mathrm{BH}}\sim10^3$–$10^4\,M_\odot$ with maximum amplification $\mathcal{A}_{\mathrm{max}}\approx\sqrt{8r_sa/R^2}\sim10^4$ and gravitational wave merger timescale $\tau_{\mathrm{GW}}\sim10^3\,\mathrm{yr}$.	3.1 Keplerian orbits
arXiv:2610.37034 2026-10-27	**FLITS: an interactive, reproducible workbench for fast radio burst analysis** *FLITS: an interactive, reproducible workbench for fast radio burst analysis* Dirk Kuiper, Ziggy Pleunis, Jason W. T. Hessels	FLITS unifies FRB data inspection, RFI mitigation, burst property extraction (signal-to-noise ratio, peak flux density, fluence, widths, and arrival times), burst-model fitting, and weighted RM synthesis (Q,U,V polarization analysis) into reproducible JSON session states with checksum validation.	Software design
arXiv:2610.36987 2026-10-27	**History of Turn-Ons - the irregular 35-day clock - of Hercules X-1** *History of Turn-Ons - the irregular 35-day clock - of Hercules X-1* Ruediger Staubert, Konstantin Postnov, Vladislav Revnivtsev 等	Turn-Ons preferentially occur at binary orbital phases $\Phi_{\mathrm{orb}}\sim0.25$ and ~0.75, with quantized cycle durations concentrated at 20.0, 20.5 (accounting for $\sim57\%$), and $21.0\times P_{\mathrm{orb}}$ ($P_{\mathrm{orb}}=1.70016759\,\mathrm{d}$). The long-term $(O-C)$ timing residual curve reveals two distinct parabolic segments separated by Anomalous Low 3 (AL3): pre-AL3 has $P=34.95\,\mathrm{d}$ and $\mathrm{d}P/\mathrm{d}t=(-1.8\pm0.2)\times10^{-5}\,\mathrm{s\,s^{-1}}$, while post-AL3 has $P=35.03\,\mathrm{d}$ and $\mathrm{d}P/\mathrm{d}t=(-5.30\pm0.23)\times10^{-5}\,\mathrm{s\,s^{-1}}$. Anomalous Lows recur with a mean period of $(5.35\pm0.03)\,\mathrm{yr}$.	7 Summary

arXiv ID / 日期	文献题名与作者	核心要点与观测约束	阅读切入
arXiv:2610.36953 2026-10-27	**Quasiparticle Modeling of Strange Quark Stars with In-Medium QCD Effects at Finite Temperature and Strong Magnetic Fields** *Quasiparticle Modeling of Strange Quark Stars with In-Medium QCD Effects at Finite Temperature and Strong Magnetic Fields* Yogesh Kumar, Mihir Arora, Manas Lohani 等	At $B = 10^{15}$ G, increasing temperature from $T = 0$ to 50 MeV stiffens the EoS, boosting the maximum gravitational mass from $M_{\max} = 1.66\,M_\odot$ ($R = 9.48$ km) to $2.19\,M_\odot$ (compatible with PSR J0740+6620 and PSR J0952-0607). Conversely, ultra-strong fields ($B = 10^{19}$ G at $T = 50$ MeV) induce Maxwell stress $\propto B^2/(8\pi)$ that softens the configuration, reducing $M_{\max}$ to $0.99\,M_\odot$ ($R = 5.41$ km). The sound speed satisfies causality and asymptotically approaches the conformal limit $v_s^2 \to 1/3$ at high densities, with low-mass radii scaling as $R \propto M^{1/3}$.	IV Comparative Study and Discussion
arXiv:2610.36867 2026-10-27	**A microscopic analysis of shear acceleration in the relativistic regime** *A microscopic analysis of shear acceleration in the relativistic regime* Sophie Aerdker, Xingwei Gong, Martin Lemoine 等	Particle shear acceleration in inhomogeneous relativistic velocity fields is governed by the effective shear tensor $\sigma(x) = \tilde{\sigma}/\gamma_{\rm J}$, which decreases in highly relativistic regions. When $\sigma\tau' \gtrsim 1$, the return time distribution between crossing events exhibits a heavy-tailed power-law $p(t_{\rm cross}) \propto t_{\rm cross}^{-1.27}$, driving sub-diffusive momentum gain ($\langle p'^2 \rangle \propto t^q, q < 1$) and causing a spectral transition/hardening at $\sigma\tau'(p') \sim 1$ from gradual shear ($n(p') \propto p'^{-2}$ for $\delta = 1$) to an intermittent non-gradual acceleration branch.	III.3 Towards non-gradual shear acceleration: $sigmatau^{p}rimegtrsim1$
arXiv:2610.36732 2026-10-27	**2:1 Harmonics in the Radio Light Curves of Supermassive Black Hole Binary Candidates** *2:1 Harmonics in the Radio Light Curves of Supermassive Black Hole Binary Candidates* A. C. S. Readhead, M. F. Aller, R. D. Blandford 等	Systematic sine-wave fits to radio monitoring light curves demonstrate robust 2:1 period commensurabilities in SMBHB candidates PKS J0805−0111, PKS J1309+1154, and PKS 2131−021, where subtracting fundamental and harmonic modes reduces residual variance by 50%–65% and 20%–44%, respectively. In the gravitational-wave-dominated inspiral regime with orbit shrinkage timescale $t_{\rm GR} \propto r^4$, the detection incidence scales with baseline duration τ as $N_2/N_1 = [\tau_2/\tau_1]^{8/3}$, explaining the scarcity of short-period ($P < 1$ yr) systems. For secondary companion-induced modulation, orbital radius and velocity scale as $r/10^{16}\,{\rm cm} \sim 1.5 P_y^{2/3} M_{\rm P9}^{1/3}$ and $v/10^9\,{\rm cm\,s^{-1}} \sim 3 P_y^{-1/3} M_{\rm P9}^{1/3}$, balancing the disk wind magnetic pressure $B_W \sim L_{j44}^{1/2} r_{16}^{-1}$ G at $r_*/r \sim 0.2 L_{j44}^{1/2} M_{\rm P9}^{1/3} P_y^{-1/3}$.	VI The Incidence of SMBHB Candidates in the OVRO+UMRAO Samples
arXiv:2610.36486 2026-10-27	**Three-flavor neutrino Boltzmann transport and flavor instabilities in long-lived neutron-star merger remnants** *Three-flavor neutrino Boltzmann transport and flavor instabilities in long-lived neutron-star merger remnants* Harry Ho-Yin Ng, Jiabao Liu, Arthur Offermans 等	Muonic weak interactions break the $\nu_\mu - \nu_\tau$ degeneracy and generate widespread collisional flavor instability (CFI) bands reaching growth rates $\sigma_{\rm CFI} \gtrsim 10^5\,{\rm s^{-1}}$ outside the ν_τ energy sphere, with narrow resonance CFI (R-CFI) zones reaching $\sigma_{\rm R-CFI} \sim 10^8$–$10^{10}\,{\rm s^{-1}}$ scaling as $\sigma^{\rm R-CFI} \sim \sqrt{\mu_{ww}\Gamma}$. In contrast to two-flavor models where fast flavor instability (FFI) is pervasive, demuonization drives ν_μ excess and depletes $\bar{\nu}_\mu$, creating single-sided lepton number angular distributions that suppress $e\mu$ and $\mu\tau$ FFI, confining FFI mainly to the $e\tau$ sector in the outer accretion disk ($\rho < 10^{11}\,{\rm g\,cm^{-3}}$). Local quantum kinetic equation (QKE) simulations demonstrate that diagonal collisional repopulation prevents flavor equipartition (deviating from $\tilde{n}_{\nu_\alpha} \simeq 1/3$), while non-linear commutators $[H_{\nu\nu}, \varrho]$ dynamically trigger initially stable flavor sectors.	IV Results of the Global simulations
arXiv:2610.36282 2026-10-27	**Reconstruction of very inclined air showers with radio data and application to the first GRANDProto300 cosmic-ray candidates** *Reconstruction of very inclined air showers with radio data and application to the first GRANDProto300 cosmic-ray candidates* Marion Guelfand, Valentin Decoene, Olivier Martineau 等	On realistic GP300 simulations, the ADF reconstruction achieves an angular resolution better than 0.07° and a primary energy resolution $< 15\%$. Applied to raw ADC data from 65 deployed antennas of GRANDProto300, it successfully identified 51 high-quality cosmic-ray candidate events with an illustrative event showing reconstructed direction $\theta = 75.1^\circ \pm 0.03^\circ$, $\phi = 358.7^\circ \pm 0.02^\circ$, reduced $\chi^2_{\nu,\rm ADF} = 4.7$, and primary energy $E = 2.08 \times 10^{18}$ eV.	2 The ADF model
arXiv:2610.36225 2026-10-27	**NOCTURNE. VI. Hour-timescale radio variability in early-stage active galactic nucleus SDSS J122844.81+501751.2** *NOCTURNE. VI. Hour-timescale radio variability in early-stage active galactic nucleus SDSS J122844.81+501751.2* E. Järvelä, P. Romano, J. Nättilä 等	The target SDSS J122844.81+501751.2 displays extreme hour-timescale variability at 37 GHz with an e-folding timescale as short as $\tau = 1.1$ h between MRO 37 GHz and JVLA 33 GHz measurements. The implied variability brightness temperature reaches $T_{\rm b,var} \sim 10^{17} - 10^{18}$ K, well in excess of the inverse Compton limit, requiring Doppler factors $\delta_{\rm var} \approx 140 - 400$. Swift XRT spectroscopy gives a soft photon index $\Gamma = 2.33^{+0.20}_{-0.18}$ and unabsorbed $0.3 - 10$ keV flux of $7.5^{+0.9}_{-0.7} \times 10^{-13}$ erg cm^{-2} s^{-1}.	5.4 Origin of the flares

arXiv ID / 日期	文献题名与作者	核心要点与观测约束	阅读切入
arXiv:2610.36103 2026-10-27	**Dynamical constraints on the companion mass from pulsar timing and cyclotron line confirmation in the SFXT IGR J16418-4532** *Dynamical constraints on the companion mass from pulsar timing and cyclotron line confirmation in the SFXT IGR J16418-4532* Ashwin Devaraj, Rahul Sharma, Ketan Rikame 等	Coherent pulsations are measured at $P_{\rm spin} = 1208.67 \pm 0.04$ s (NuSTAR) and $P_{\rm spin} = 1209.4178 \pm 0.0084$ s (RXTE/PCA), with projected semi-major axes $a_x \sin i = 55.1 \pm 3.5$ lt-s and 51.3 ± 2.6 lt-s, respectively. Combined with binary geometry and Roche-lobe constraints, the companion star mass is constrained to $M_{\rm c} \simeq 12$–$18\ M_\odot$ and radius $R_{\rm c} \simeq 13$–$16\ R_\odot$. The confirmed CRSF at $E_{\rm cyc} \sim 50$ keV implies a neutron star surface magnetic field of $B \approx 5 \times 10^{12}$ G.	4 Discussion and Conclusion
arXiv:2610.36101 2026-10-27	**Phase-Twist Instability in a Coprecessing Neutrino Plasma** *Phase-Twist Instability in a Coprecessing Neutrino Plasma* Damiano F. G. Fiorillo, Dwaipayan Mukherjee, Basudeb Dasgupta 等	The authors identify a gapless phase-twist hydrodynamic instability for coprecessing neutrino polarization vectors with dispersion $\Upsilon_{\mu\nu}K^\mu K^\nu = 0$ (i.e. $\omega^2\Upsilon_{00} + k^2\Upsilon_{11} + \omega k(\Upsilon_{01} + \Upsilon_{10}) = 0$), yielding imaginary frequencies $\mathrm{Im}(\omega) = \gamma \sim k\Delta v\Delta\theta$ in the hydrodynamic regime and saturating at maximum growth rate $\gamma_{\max} \sim \lvert\vec{H}_{\rm p}\rvert\Delta\theta^2$ around $k \sim \lvert\vec{H}_{\rm p}\rvert\Delta\theta/\Delta v$.	Phase-Twist Instability in a Coprecessing Neutrino Plasma
arXiv:2610.36074 2026-10-27	**CROMA: a GPU-accelerated Fokker--Planck solver for modeling non-thermal emission in post-processing cosmological simulations** *CROMA: a GPU-accelerated Fokker--Planck solver for modeling non-thermal emission in post-processing cosmological simulations* Kosuke Nishiwaki, Claudio Gheller, Gianfranco Brunetti 等	CROMA implements an implicit Chang-Cooper scheme solved via Parallel Cyclic Reduction (PCR) on GPUs, achieving a tridiagonal solve throughput saturating at $R_{\rm solve} \approx 10^3$ tracers s^{-1} per MPI task, an end-to-end wall-clock throughput of $R_{\rm wall} \approx 400$ tracers s^{-1} per task at $N_{\rm trc} \sim 10^6$ ($N_{\rm bin} = 128$), an acceleration factor of $\sim 10\times$ compared to 32 CPU cores, and a synchrotron emissivity computation speedup by a factor of $N_\theta \approx 10^3$ using precomputed tables.	2 Fokker–Planck equation
arXiv:2610.36050 2026-10-27	**An intermediate-mass stripped star with a dark companion in the Small Magellanic Cloud** *An intermediate-mass stripped star with a dark companion in the Small Magellanic Cloud* Lisa Blomberg, Kareem El-Badry, Peter Senchyna 等	The binary has an orbital period of $P_{\rm orb} = 19.95 \pm 0.002$ hr (19.947 ± 0.002 hr), radial velocity semi-amplitude $K = 91.7^{+3.2}_{-3.1}$ km s^{-1}, and eccentricity $e = 0.032^{+0.031}_{-0.022}$, yielding a binary mass function $f_m \approx 0.066\ M_\odot$. The stripped star has an evolutionary mass $M_{\rm evol} = 3.840 \pm 0.535\ M_\odot$ ($M_{\rm stripped} = 3.84 \pm 0.54\ M_\odot$), $T_{\rm eff} = 68.3 \pm 6.4$ kK, $\log(g/[\mathrm{cm\,s^{-2}}]) = 4.76 \pm 0.12$, radius $R = 0.99 \pm 0.09\ R_\odot$, luminosity $\log(L/L_\odot) = 4.28 \pm 0.13$, and an anomalously high surface hydrogen mass fraction $X_{\rm H,s} = 0.625^{+0.012}_{-0.011}\,(\mathrm{stat}) \pm 0.035\,(\mathrm{sys})$. For $M_{\rm stripped} = 3.84\ M_\odot$, the companion's minimum mass is $M_2 = 1.19\ M_\odot$ ($i = 90^\circ$) with a median of $1.41\ M_\odot$. Optical light curve limits rule out main-sequence companions (which would produce reflection variability $\geq 10\%$ vs observed limit $\lesssim 1\%$), and X-ray non-detection sets $L_X \lesssim 9 \times 10^{33}$ erg s^{-1} ($0.2 - 12$ keV), highly consistent with a neutron star companion.	IV.4 Constraining the companion mass
arXiv:2610.35849 2026-10-26	**Interaction of Strong X-mode Waves with Magnetized Pair Plasmas** *Interaction of Strong X-mode Waves with Magnetized Pair Plasmas* Navin Sridhar	The authors derive the perturbation dynamics and coupled dispersion relation for the induced scattering instability, $D_{\rm v}^2 - f_0\Omega_{\rm B}^2 \simeq \frac{2\eta^3}{\omega}k(\Omega + k)\left(\frac{1}{D_+} + \frac{1}{D_-}\right)$, where $D_{\rm v} \equiv \Omega + v_x k$, $D_\pm \equiv (\Omega \pm \omega)^2 - k^2 - \omega_{\rm p}^2$, and the scattering parameter is $\eta \equiv \omega(\mathcal{A}\varepsilon_{\rm p}/2)^{2/3}$. In the strong-coupling limit ($\Omega_{\rm B} \ll \eta$), the maximum growth rate scales as $\Gamma_{\rm gr} = \frac{\sqrt{3}}{2}\eta$, while in the weak-coupling limit ($\Omega_{\rm B} \gg \eta$), the instability growth is modified by the magnetic restoring frequency $\Omega_{\rm B} \equiv \sqrt{1 + \hat{a}^2/2}(\omega_{\rm B}/\gamma_0)$.	Interaction of Strong X-mode Waves with Magnetized Pair Plasmas
arXiv:2610.35648 2026-10-26	**The long-term evolution of a black hole accretion torus: a comparison between GRMHD simulations and relativistic thin disk theory** *The long-term evolution of a black hole accretion torus: a comparison between GRMHD simulations and relativistic thin disk theory* Andrew Mummery, Minghao Guo, Eliot Quataert 等	The long-term evolution of the bound accretion disk and mass accretion rate follows power-law decay behaviors: $\dot{M}(t) \propto t^{-n}$ with $n = (4 - 2\mu)/(3 - 2\mu)$, outer disk radius $r_{\rm out} \sim \sigma_{\rm eff}(t) \sim t^{2/(3-2\mu)} \sim t^{2n-2}$, and peak line mass $M'_{\rm max} \sim t^{1-n}$. For a scale-independent turbulent stress $W^r_\phi \approx w \approx 0.05$ ($\mu = 0, n = 4/3$), General Relativistic Thin Disk Theory (GRTDT) reproduces disk spreading within a factor of ~ 2, though it over-predicts the horizon accretion rate by a factor of ~ 2 at $t \gtrsim 2 \times 10^4\, t_{\rm g}$ because unbound disk winds carry away $\sim 50\%$–80% of the lost mass.	IV.2 General Relativistic Thin Disk Theory

arXiv ID / 日期	文献题名与作者	核心要点与观测约束	阅读切入
arXiv:2610.35602 2026-10-26	**Nonlinear Tides of Binary Neutron Stars** *Nonlinear Tides of Binary Neutron Stars* Guangzhou Guo, Zhen Pan	In the late inspiral stage ($D = 2.5R_*$), nonlinear tidal interactions reduce the neutron star quadrupole fundamental mode (f-mode) frequency by $\sim 10\%$ for equal-mass binaries and impart an extra gravitational-wave phase shift of $\Delta\Phi_X \sim \mathcal{O}(1)\ \mathrm{rad}$ beyond the linear tidal response up to $f_{\rm GW} = 1100\ \mathrm{Hz}$. The second-order perturbative frequency shift scales as $\Delta f^{(2)} \propto q^2$ (where $q = M_c/M_*$), and the radial mode frequency drops by 8.8% to $f_r = 0.3173$ due to nonlinear c_{20}–c_r coupling.	Nonlinear Tides of Binary Neutron Stars
arXiv:2610.35589 2026-10-26	**Gravitationally Induced Polarization-Angle Rotation in Pulsar-Black Hole Binaries** *Gravitationally Induced Polarization-Angle Rotation in Pulsar-Black Hole Binaries* Jyotijwal Debnath, Amodio Carleo, Delphine Perrodin 等	Gravitational deflection induces a geometric polarization position angle (PPA) shift $\Delta\psi_{\rm geo}$ reaching $\sim 7^\circ$ near superior conjunction ($A_T + \omega = 90^\circ$) in a pulsar–IMBH system ($M_c = 10^5\,M_\odot$, $P_b = 0.5$ yr, $i = 85^\circ$), while black hole spin ($\tilde{a} = 0.5$) generates a frame-dragging geometric shift $\Delta\psi_{\rm FD,geo}$ of millidegree order and a parallel-transport rotation $\Delta\psi_{\rm pt}$ of microdegree order. In a close pulsar–SMBH binary ($M_c = 10^6\,M_\odot$, $P_b = 5.5$ days, $i = 70^\circ$), strong gravitational deflection reaches up to 180°, $\Delta\psi_{\rm pt}$ increases by a factor of ~ 500, and the characteristic RVM S-shaped polarization swing is completely reversed at $A_T + \omega = 85^\circ$ and 90°.	VI Numerical Study
arXiv:2610.35573 2026-10-26	**Classification of Fermi Blazars of Uncertain Type Using Positive-Unlabelled Machine Learning** *Classification of Fermi Blazars of Uncertain Type Using Positive-Unlabelled Machine Learning* Sikandar Akbar, Zahir Shah, Zahoor Malik 等	Out of 1623 Fermi 4FGL-DR4 BCUs, the PU ensemble classifies 437 sources as FSRQ-like candidates, 597 as BL Lac-like candidates, flags 12 sources as ambiguous candidates meeting both criteria, and leaves 577 sources unclassified. The BCU FSRQ candidates exhibit a softer median photon index of $\Gamma = 2.60$ compared to known FSRQs ($\Gamma = 2.47$), while unclassified BCUs are systematically fainter (median $\log F_{1000} = -9.77$) with 25.3% located in the Galactic plane ($\lvert b\rvert < 10^\circ$).	VI Summary and Conclusions
arXiv:2610.35379 2026-10-26	**Revisiting supernova constraints on dark gauge bosons** *Revisiting supernova constraints on dark gauge bosons* Zun Wang, Yongchao Zhang	Using the SFHo-18.8 ($18.8\,M_\odot$) core profile, the SN1987A cooling bounds exclude gauge boson masses up to $m_{A'} \lesssim 230\ \mathrm{MeV}$, $m_{Z'} \lesssim 280\ \mathrm{MeV}$ ($B-L$), and $m_{Z'} \lesssim 300\ \mathrm{MeV}$ ($L_\mu - L_\tau$). LESN mantle deposition limits constrain couplings down to $\epsilon \gtrsim 5.8\times10^{-12}$ for dark photons ($m_{A'} \le 450\ \mathrm{MeV}$), $g_{BL} \gtrsim 1.9\times10^{-12}$ for $U(1)_{B-L}$ ($m_{Z'} \le 440\ \mathrm{MeV}$), and $g_{\mu\tau} \gtrsim 6.4\times10^{-12}$ for $U(1)_{L_\mu-L_\tau}$ ($m_{Z'} \le 600\ \mathrm{MeV}$), scaling in the optically thin limit as $E_{\rm dep}^{(A')} \propto \epsilon^2$ and $E_{\rm dep}^{(Z')} \propto g_{BL}^4$.	Revisiting supernova constraints on dark gauge bosons
arXiv:2610.35181 2026-10-26	**Investigating supernova-remnant X-ray maps with dendrogram-based techniques** *Investigating supernova-remnant X-ray maps with dendrogram-based techniques* S. Calderaro, G. Parisi, V. Sapienza 等	The dendrogram and GNN embedding analyses reveal distinct elemental spatial co-locations (e.g. Si with S, Ar with Ca, and Fe XXV with Ni XXVI) and demonstrate that W49B clusters closer to core-collapse (CC) SNRs (Cas A, G292.0+1.8) than to Type Ia SNRs (Tycho, Kepler). Computing runtime scales from $15937.0\ \mathrm{ms}$ for GNN and $16962.7\ \mathrm{ms}$ for cosine-distance HC to $1365064.8\ \mathrm{ms}$ for bootstrap-validated SVHC over a 50-band dataset.	Results
arXiv:2610.35174 2026-10-26	**The biases caused by interpolated table models and their impact on X-ray spectroscopy** *The biases caused by interpolated table models and their impact on X-ray spectroscopy* Matteo Lucchini, Benjamin Ricketts, Phil Uttley 等	Linear interpolation of non-linear models introduces a single-parameter error $e(\theta) = m(\theta) - \tilde{m}(\theta) = -\frac{1}{2}m''(\xi)(\theta-\theta_i)(\theta_{i+1}-\theta)$, which distorts the log-likelihood by adding a quadratic penalty $-\frac{1}{2}\sum_k \frac{e_k^2(\theta)}{\sigma_k^2}$ and a residual cross-term $-\sum_k \frac{e_k(\theta)[d_k-m_k(\theta)]}{\sigma_k^2}$. Its first derivative $\frac{d\ln\tilde{\mathcal{L}}}{d\theta} = \frac{d\ln\mathcal{L}}{d\theta} - \sum_k \frac{e_k e_k' - e_k m_k' + e_k'(d_k-m_k)}{\sigma_k^2}$ introduces discontinuous jumps at grid points via $e_k'(d_k-m_k)$, generating artificial modes. In real NuSTAR fits of MAXI J1820+070, table models induced artificial multimodality (e.g., spin a, photon index $\Gamma = 1.6$), causing affine-invariant MCMC acceptance rates to drop to $\approx 5\%$ and autocorrelation times to blow up to $\tau \approx 4000$–16000.	3. Likelihood distortions

arXiv ID / 日期	文献题名与作者	核心要点与观测约束	阅读切入
arXiv:2610.35156 2026-10-26	**$Δ(1232)$ isobars and neutrino opacities in hot dense matter** *$Δ(1232)$ isobars and neutrino opacities in hot dense matter* Subha Mitra, Bharat Kumar	When the threshold condition $\mu_n + \mu_e > M^*_{\Delta^-} + U_{\Delta^-}$ is met, the three-body absorption reaction $\bar{\nu}_e + e^- + n \to \Delta^-$ dominates the opacity, suppressing the antineutrino Rosseland mean free path λ_R from 3–300 m down to 0.5–16 cm (a reduction factor of 83 to 4.5×10^4) at densities of 2–4n_0. For electron neutrinos, λ_R is shortened by factors of 1.7–6.5 (for $Y_L = 0.2$–0.3) and up to 4–200 (for $Y_L = 0.1$, following $\lambda_R^N / \lambda_R^{N\Delta} \approx \exp(aY_\Delta)$). Consequently, the antineutrino share of the lepton-number diffusion coefficient D_2 drops from up to 69% to below 7%, reducing total lepton diffusion by up to a factor of 3 at $Y_L = 0.1$, while total energy diffusion D_4 changes by $< 2\%$.	VI. DISCUSSION AND CONCLUSIONS
arXiv:2610.35117 2026-10-26	**Torque lags from magnetic unlocking in neutron stars and gravitational-wave smearing** *Torque lags from magnetic unlocking in neutron stars and gravitational-wave smearing* Arthur G. Suvorov, Armen Sedrakian	Magnetic unlocking produces differential rotation with lag $\Delta\nu \sim -\tau_{\rm eff}\dot{\nu}_{\rm R}$ between the core toroidal region and the crust, separating quadrupoles $\epsilon_{\rm pol}$ and $\epsilon_{\rm tor}$ into a distinct or smeared spectrum. The resulting continuous GW signal spans $\Delta f_{\rm GW} \approx 2\Delta\nu$, diluting the coherent Fourier strain amplitude per bin by $\lvert\tilde{h}_{\rm bin}\rvert/\lvert\tilde{h}_{\rm line}\rvert \sim (\Delta f_{\rm GW} T)^{-1}$ and degrading the detection sensitivity by a factor of $(\Delta f_{\rm GW} T)^{1/2}$ for semicoherent searches when $\Delta f_{\rm GW} T \gg 1$.	2 Gravitational-wave spectrum
arXiv:2610.35087 2026-10-26	**Comparison of Hybrid and Standard Reconstruction Methods on IXPE Astrophysical Data** *Comparison of Hybrid and Standard Reconstruction Methods on IXPE Astrophysical Data* Riccardo Ferrazzoli, Nicolò Cibrario Stefano Silvestri, Michela Negro 等	In the accreting X-ray pulsar Her X-1, the hybrid method increases polarimetric significance above 2.83 keV, reaching a significance ratio $R = \sigma_{\rm hyb}/\sigma_{\rm std} = 1.180 \pm 0.029$ (6.2σ above unity) at 5.66–8.00 keV, which corresponds to an equivalent effective area increase of $\Delta A_{\rm eff}/A_{\rm eff} = R^2 - 1 = 39.3\% \pm 6.8\%$. In the unpolarized source XTE J1701-461, no spurious polarization is induced ($R \leq 1.7\sigma$ from 1 across all bands).	2.2 Significance ratio and its uncertainty
arXiv:2610.34848 2026-10-26	**A frequency-resolved measurement of the scattering law in FRB 20210415C** *A frequency-resolved measurement of the scattering law in FRB 20210415C* Victor Pop, Mohammed A. Chamma, Aishwarya Kumar 等	Across 8 sub-bands spanning 400.2–765.6 MHz, the scattering index is measured to be $\alpha = -3.95 \pm 0.07$ (free Gaussian width) and $\alpha = -4.14 \pm 0.08$ (fixed Gaussian width), with a 34-parameter joint MCMC yielding $\alpha = -3.95 \pm 0.06$ and $\tau_{\rm ref} = 7.97 \pm 0.09$ ms at $\nu_{\rm ref} = 583$ MHz (corresponding to $\tau_{\rm s} \simeq 7.0$–7.1 ms at 600 MHz). The intrinsic pulse width exhibits frequency narrowing with $\sigma_{\rm int} \propto \nu^{\beta}$ ($\beta = -0.92 \pm 0.19$ for $\nu > 440$ MHz), with low-frequency excess width (4.8σ at ~ 422 MHz) favoring a thick-screen scattering medium.	3 Results
arXiv:2610.34715 2026-10-26	**Gaia DR3 Compact Binary Candidates Constrained by Radial Velocity Measurements** *Gaia DR3 Compact Binary Candidates Constrained by Radial Velocity Measurements* Ying-jie Li, Xianjin Shen, Zehao Lin 等	Among the 518 binaries with effective radial velocity epochs $n_{\rm eff} \geq 3$, 192 are classified as Good-fit sources, containing 171 Medium-priority or higher compact binary candidates; this includes 4 candidates with fitted secondary mass $M_{2,\rm fit} > 2.1\,M_\odot$ (including a Critical candidate with $M_{2,\rm fit} \sim 2.3\,M_\odot$ and companions up to $\sim 3.2\,M_\odot$), 22 sources with $1.4\,M_\odot \leq M_{2,\rm fit} \leq 2.1\,M_\odot$, and 145 sources with $M_{2,\rm fit} < 1.4\,M_\odot$.	5. CONCLUSIONS
arXiv:2610.34699 2026-10-26	**Binary Progenitors of Electron-capture Supernovae** *Binary Progenitors of Electron-capture Supernovae* Liji Zhong, Yong Shao	In surviving binaries, stripping produces predominantly Type Ib ($\sim 60\%$), Type II ($\sim 30\%$), and Type IIb ($\sim 10\%$) events, while Type Ic is $< 0.4\%$. Accretion spin-up yields near-critical companion rotation ($\omega/\omega_{\rm crit} \geq 0.8$) concentrated at $P_{\rm orb,postSN} \sim 10 - 100$ d and $e \simeq 0.1 - 0.3$, matching BeXRB precursors, with a local cosmic ECSN rate density of $(4-7) \times 10^3$ Gpc^{-3} yr^{-1} representing $f_{\rm ECSN} \approx 1\% - 4\%$ of core-collapse events.	3.2 Pre-supernova structure, classification, and mass loss
arXiv:2610.34515 2026-10-26	**Analytical modeling of the gamma-ray morphology of TeV halos around moving pulsars** *Analytical modeling of the gamma-ray morphology of TeV halos around moving pulsars* Sarah Recchia	The authors derive a closed-form analytical approximation for the projected line-of-sight (LOS) electron surface density $\Sigma(\rho, \phi, E) = \mathcal{S}(E)\mathcal{T}(\rho, \phi, E)$ around a moving pulsar. In the far-field limit $\rho, v\tau, r_d \ll d$, the 3D diffusion-loss convolution reduces to a single integral over injection history, and upon spatial integration $\int_0^{2\pi} d\phi \int_0^{\infty} \rho d\rho\, \Sigma$, the dependence on diffusion coefficient $D(E)$ and pulsar velocity v strictly cancels, yielding an equilibrium lepton spectrum $N_{\rm eq}(E) \simeq q(E)\tau_{\rm max}(E)$.	3 Derivation of the gamma-ray morphology

arXiv ID / 日期	文献题名与作者	核心要点与观测约束	阅读切入
arXiv:2610.34478 2026-10-26	**Resolving a Clumpy Disk Wind in 4U 1957+115 with XRISM** *Resolving a Clumpy Disk Wind in 4U 1957+115 with XRISM* Paul A. Draghis, Erin Kara, Ehud Behar 等	The observation detects Fe XXVI and Fe XXV absorption lines with confidence up to 5.6σ during orbital phase $\phi \sim 0$, revealing an outflow velocity $v_{\rm out} = 140^{+30}_{-20}$ km/s and an intrinsic turbulence limit $v_{\rm turb} \lesssim 70$ km/s (narrowed down from $v_{\rm turb} = 350^{+290}_{-170}$ km/s in time-averaged spectra). Photoionization modeling with pion yields $\log \xi = 5.8^{+0.3}_{-0.5}$ and $N_{\rm H,PION} = 4.8^{+1.7}_{-3.7} \times 10^{23}$ cm^{-2}, placing the maximum wind launching radius at $r_{\rm max} = L_{\rm ion}/(\xi N_{\rm H}) \sim 10^8 - 10^{10} (d_{10})^2$ cm, far inside the Compton radius $R_{\rm IC} = (5 \pm 1) \times 10^{11}$ cm and below the local escape velocity.	4. RESULTS AND DISCUSSION
arXiv:2610.34423 2026-10-26	**Constraining Nuclear Matter from Finite Nuclei: the Ab Initio Perspective** *Constraining Nuclear Matter from Finite Nuclei: the Ab Initio Perspective* Francesca Bonaiti, Sonia Bacca, Ingo Tews	Ab initio chiral effective field theory predictions determine neutron-skin thicknesses $R_{\rm skin}(^{48}{\rm Ca}) = 0.120(14)(12)$ fm and $R_{\rm skin}(^{208}{\rm Pb}) = 0.187(25)(18)$ fm, favoring a soft-to-moderate symmetry energy slope $38.3 \leq L \leq 68.5$ MeV and $J \sim 30$–32 MeV consistent with GW170817 and NICER astrophysical constraints, while isoscalar giant monopole resonance (ISGMR) data constrain symmetric nuclear matter incompressibility to $K_0 \sim 190$–240 MeV.	2.2. Dense Matter in Nature: Neutron Stars
arXiv:2610.34395 2026-10-26	**ePNGCAT: the first eROSITA all-sky data release** *ePNGCAT: the first eROSITA all-sky data release* Victor Doroshenko	The catalogue contains 63,260 sources over 99.54% of the sky with an estimated 50% completeness limit of 3.12×10^{-13} erg cm^{-2} s^{-1} in the 0.2--2.3 keV band, a median positional offset of $72''$, and an overall multiplicative broad-band flux dispersion of 26.5% relative to eRASS1.	6. Conclusions
arXiv:2610.34259 2026-10-26	**Contributions of massive stars and classical novae to Galactic 26Al** *Contributions of massive stars and classical novae to Galactic 26Al* Manja L. Zimmerer, Thomas Siegert, Niklas C. Bauer 等	Fitting the radial 1.809 MeV emission across Galactocentric annuli reveals that classical novae dominate Galactic ^{26}Al with a baseline contribution of $77^{+14}_{-15}\%$, while massive stars account for only $23^{+13}_{-14}\%$. A massive star fraction of $\sim 23\%$ yields a revised Galactic star formation rate of ${\rm SFR} = 1.2$–$1.8\,M_\odot\,{\rm yr}^{-1}$ and increases the massive-star/core-collapse supernova $^{60}{\rm Fe}/^{26}{\rm Al}$ yield ratio to 0.8 ± 0.2, matching theoretical predictions.	5 Relative contribution of classical novae and massive stars
arXiv:2610.34234 2026-10-26	**Probing In-Medium Neutron-to-Dark-Fermion Conversion with Multimessenger Neutron Star Observations** *Probing In-Medium Neutron-to-Dark-Fermion Conversion with Multimessenger Neutron Star Observations* Davood Rafiei Karkevandi, Mahboubeh Shahrbaf, Hamid Reza Moshfegh 等	In-medium neutron-to-dark-fermion conversion softens the nuclear EOS by reducing neutron fraction Y_n and increasing dark fermion abundance $Y_\chi = n_\chi/n_B$. Weakly repulsive regimes ($G_\chi = 1$--11 fm^2, $m_\chi = 939$--1300 MeV) fail to support $M_{\rm max} \geq 2M_\odot$ and are ruled out across the nuclear EOS ensemble. In the Illinois-Maryland analysis, the active conservative boundary is driven by $M_{\rm max} \geq 2M_\odot$, while strict X-PSI/Amsterdam tests yield no common parameter point satisfying all 13 baseline EOSs, with the maximum compatibility ($N_{\rm cons} = 11$) confined to $m_\chi = 939$--950 MeV and $G_\chi = 19$--21 fm^2.	II.4 Composition: $beta$ equilibrium and charge neutrality
arXiv:2610.34071 2026-10-26	**A MAGNHIFFIC cold-gas filament fuelling the nuclear activity of NGC 3100** *A MAGNHIFFIC cold-gas filament fuelling the nuclear activity of NGC 3100* F. M. Maccagni, I. Prandoni, F. Fonzo 等	The authors detect a ~ 340 kpc stellar-free H I filament with mass $M_{\rm HI} = 3.4 \times 10^8\ M_\odot$ feeding the circum-nuclear disc of early-type galaxy NGC 3100 ($D = 38$ Mpc). High-velocity-resolution absorption against the core reveals redshifted infalling cold gas clouds with column density $N_{\rm HI} = 63 \times 10^{16}\ (T_{\rm spin}/c_f)$ cm^{-2} and mass $M_{\rm HI} \sim 2.4 \times 10^4\ M_\odot$, coexisting with jet-driven molecular gas outflows of mass $M_{\rm out} = 5.6 \times 10^4\ M_\odot$ and outflow rate $\dot{M}_{\rm out} \lesssim 0.12\ M_\odot\ {\rm yr}^{-1}$ driven by a low-power jet ($P_{1.4\,{\rm GHz}} = 1 \times 10^{23}$ W Hz^{-1}).	7.3 Jet-ISM interaction in NGC 3100 and its black-hole weather
arXiv:2610.34058 2026-10-26	**Experimental Evidence for Reflected Ions and Ion Temperature Anisotropy in a Forming Supercritical Magnetized Collisionless Shock** *Experimental Evidence for Reflected Ions and Ion Temperature Anisotropy in a Forming Supercritical Magnetized Collisionless Shock* Ryo Yamazaki, T. Takezaki, T. Morita 等	The authors detect a foot-like structure and pronounced ion temperature anisotropy ($T_{i,\perp}/T_{i,\parallel} = T_{i,x}/T_{i,y} \lesssim 4.5$) upstream of the shock ramp in a supercritical collisionless shock with upstream parameters $v_A \approx 65$ km s^{-1}, sound speed $a_s \approx 11$ km s^{-1}, $M_A > 3.8$, $M_s > 23$, $\beta_i \approx 0.02$, and $\beta_e \approx 0.06$. Linear kinetic dispersion analysis for foot plasma ($T_{i,\parallel} \sim 700$ eV, $T_{i,\perp} \sim 2300$ eV) demonstrates susceptibility to the Alfvén ion cyclotron (AIC) instability with growth rate $\gamma_{\rm max} = 0.31\,\omega_{ci}$ at $B = 4$ T (characteristic growth timescale $\tau \sim 20$ ns, wavelength $\lambda_{\rm max} \sim 4$ mm).	Experimental Evidence for Reflected Ions and Ion Temperature Anisotropy in a Forming Supercritical Magnetized Collisionless Shock

arXiv ID / 日期	文献题名与作者	核心要点与观测约束	阅读切入
arXiv:2610.34057 2026-10-26	**Identification of rippling interface between laser-ablated plasma and ambient uniform plasma at rest** *Identification of rippling interface between laser-ablated plasma and ambient uniform plasma at rest* Ryo Yamazaki, S. Tomita, T. Takezaki 等	Using optical self-emission and collective Thomson scattering (CTS), the authors identify a rippling interface between the expanding laser-ablated Al piston and uniform N ambient plasma. Upstream parameters with $\lvert B_{y0}\rvert = 4$ T yield an Alfvén speed $v_{\rm A} = \lvert B_{y0}\rvert/\sqrt{4\pi m_{\rm N} n_{\rm N}} \approx 92\ {\rm km\ s^{-1}}$, which together with expansion velocity $V_{X'} \approx 500\ {\rm km\ s^{-1}}$ confirms a supercritical shock regime. CTS diagnoses electron density $n_e = (1.97 \pm 0.26) \times 10^{18}\ {\rm cm^{-3}}$, $Z_{\rm Al} \approx 11$, and temperatures $T_e = 377 \pm 19$ eV and $T_i = 882 \pm 16$ eV, confirming that the bright emission layer is dominated by piston Al plasma rather than compressed ambient plasma.	IV.3 Simulation Results
arXiv:2610.33954 2026-10-26	**Core-Collapse Supernovae from Single and Binary-Stripped Massive Stars - I. Explodability and Nucleosynthesis** *Core-Collapse Supernovae from Single and Binary-Stripped Massive Stars - I. Explodability and Nucleosynthesis* J. M. Pollin, S. Curtis, C. Fröhlich 等	Progenitors with $M_{\rm ZAMS} \leq 22\,{\rm M_\odot}$ explode reliably across both populations, yielding explosion energies $E_{\rm exp} \sim 0.4$–1.3×10^{51} erg (0.4–1.3 B), baryonic remnant masses $M_{\rm rem} \sim 1.4$–$2.0\,{\rm M_\odot}$, and explosion timescales $t_{\rm exp} \sim 0.30$–0.65 s. The Salpeter IMF-weighted Fe yield for binary-stripped models is $0.0589\,{\rm M_\odot}$ (consistent with empirical observations of $0.058 \pm 0.007\,{\rm M_\odot}$), while single-star progenitors yield $0.0747\,{\rm M_\odot}$. Binary stripping enhances K yields by ~ 3.4 times (~ 0.63 dex) and systematically improves agreement with galactic [S/Mg], [K/Mg], and [Ca/Mg] trends.	4.1 Two Representative Models: Ms18 and Mb18
arXiv:2610.33879 2026-10-26	**Finite-temperature ab-initio-Informed Nuclear Equation-of-State Tables (FINEST): Nuclear matter uncertainties in core-collapse supernovae** *Finite-temperature ab-initio-Informed Nuclear Equation-of-State Tables (FINEST): Nuclear matter uncertainties in core-collapse supernovae* Finia P. Jost, André da Silva Schneider, Almudena Arcones 等	Cold nuclear matter parameters introduce up to ~ 150 ms delay in shock revival for high-compactness progenitors ($s24$, $\xi_{2.5} = 0.44$), while very soft high-density matter ($P_{\rm SNM} = 400\ {\rm MeV\ fm^{-3}}$) triggers black hole collapse within 500 ms. Crucially, the thermal pressure is controlled by the nucleon effective mass m^* via the thermal index $\Gamma_{\rm th} = 1 + P_{\rm th}/\epsilon_{\rm th} \approx \frac{5}{3} - \frac{n}{m^*}\frac{\partial m^*}{\partial n}$. A non-monotonic $m^*(n)$ functional ($m^*_{\rm HWS}$) reduces thermal pressure at densities exceeding saturation density n_0, accelerating protoneutron star (PNS) mantle contraction, which pushes neutrinospheres into hotter layers to enhance neutrino luminosity and successfully trigger delayed explosions, whereas standard monotonically decreasing Skyrme forms ($m^*_{\rm skyrme}$) fail to revive the shock and collapse into a black hole at $t_{\rm pb} \approx 1.65$ s.	IV Impact of the Effective Mass Parametrization
arXiv:2610.33865 2026-10-26	**Magnetoelastic avalanches in magnetar crusts with plastic islands** *Magnetoelastic avalanches in magnetar crusts with plastic islands* Arthur G. Suvorov	Dynamic crustal failure leads to collective avalanches and slow creeping failures that multiply the failed volume fraction by a factor of ≈ 4 (from $f_{\rm fail}(0) \approx 9\%$ to $f_{\rm fail} = 35.4\%$ for global quakes, and 0.6% to 2.6% for localized quakes with $E_{\rm kin,0} = 7.07 \times 10^{41}$ erg). Avalanching develops promptly within ≈ 8 ms , accompanied by discrete QPO modes (e.g., $18.79, 24.01, 49.52$ Hz, and localized high-frequency modes at ~ 113.5 Hz) that drift over beat timescales $T_{\rm beat} \sim \lvert\omega_1 - \omega_2\rvert^{-1} \sim 1$ s. The failure volume scales non-monotonically with the poloidal field B_p peaking around $B_p \sim 10^{14}$ G, and scales with kick efficiency as $\frac{f_{\rm fail}(50\ {\rm ms})}{f_{\rm fail}(0)} = 3.7 + 0.193 \ln \eta$.	5 Results: global quakes
arXiv:2610.33844 2026-10-26	**Far Ultraviolet Variability on Timescales of a Few hours in the Recurrent Nova M31N 2008-12a: Origin and Implications** *Far Ultraviolet Variability on Timescales of a Few hours in the Recurrent Nova M31N 2008-12a: Origin and Implications* Judhajeet Basu, Sudhanshu Barway, G. C. Anupama 等	AstroSat/UVIT observations detect intrinsic FUV variability ($R > 4$) at ~ 4.5–5.1 days post-eruption with a candidate modulation period of $P = 4.60^{+0.25}_{-0.15}$ hr (2.3σ, FAP $= 0.02$). The inferred Keplerian radius satisfies $r_{\rm Kep} < r_{\rm circ} < r_{\rm RL}$, ruling out fresh viscous re-accretion (which requires $t_{\rm visc} \sim 100$ days) and demonstrating accretion disk survival with low ejecta mass $M_{\rm ej} \sim 7.9 \times 10^{-8}\,M_\odot$.	V.4 Disk Survival or Rapid Re-establishment?
arXiv:2610.33696 2026-10-26	**Effects of Synchrotron Self-Absorption on Polarization Diagnostics of MHD Turbulence** *Effects of Synchrotron Self-Absorption on Polarization Diagnostics of MHD Turbulence* Ning-Ning Guo, Jian-Fu Zhang, Hua-Ping Xiao 等	SSA significantly suppresses polarized intensity and weakens spatial anisotropy while preserving the Kolmogorov power-law scaling $k^{-8/3}$ in the inertial range of the polarized power spectrum. The self-absorption turnover frequency scales as $\nu_a \propto L^{0.30}$ and $\nu_a \propto B^{1.30}$ (consistent with theoretical $\nu_a \propto (\kappa B_\perp^{(p+2)/2} L)^{2/(p+4)}$), with total intensity reaching $I(\nu) \propto \nu^{5/2}$ in the optically thick regime and $I(\nu) \propto \nu^{-(p-1)/2} = \nu^{-0.85}$ in the optically thin regime. Furthermore, the variance-wavelength relation $P^2(\lambda^2)$ is modified in the presence of SSA: when $\bar{\phi} > \sigma_\phi$, the scaling shifts from $\lambda^{-2-2m+(p-1)}$ to $P^2 \propto \lambda^{-2-2m-5}$, preserving sensitivity to the magnetic turbulence spectral index m.	4 Results

arXiv ID / 日期	文献题名与作者	核心要点与观测约束	阅读切入
arXiv:2610.33548 2026-10-26	**PeV Diffuse Neutrinos from Magnetically Arrested Disks in Low-Luminosity Active Galactic Nuclei** *PeV Diffuse Neutrinos from Magnetically Arrested Disks in Low-Luminosity Active Galactic Nuclei* Fan Wu, Benzhong Dai	For a baseline MAD fraction estimated from the low-excitation radio galaxy luminosity function $f_{\rm MAD}^{\rm LF} \simeq 1.6 \times 10^{-3}$, the all-flavor diffuse neutrino flux peaks at $E_\nu \simeq 1.5\,{\rm PeV}$ with $E_\nu^2 \Phi_\nu \simeq 2.6 \times 10^{-9}\,{\rm GeV\,cm^{-2}\,s^{-1}\,sr^{-1}}$, primarily driven by $p\gamma$ interactions on inverse-Compton photons from thermal electrons. Comparison with IceCube observations constrains the maximum fraction of MAD-hosting LLAGNs to $f_{\rm MAD}^{\rm max} \simeq 2.3 \times 10^{-3} [0.0495/(\epsilon_{\rm NT}\epsilon_{\rm dis})]$, where $\epsilon_{\rm NT} \simeq 0.33$ and $\epsilon_{\rm dis} \simeq 0.15$.	III Diffuse Neutrino Intensity from the LLAGN Population
arXiv:2610.33443 2026-10-26	**Fast Outflow Driven from the Magnetically Arrested Disc** *Fast Outflow Driven from the Magnetically Arrested Disc* Taili Yi, Jia-Wen Li	Magnetically driven outflows from MADs smoothly traverse slow-magnetosonic, Alfvén, and fast-magnetosonic critical points, attaining mildly relativistic velocities ($v_{\rm p} \gtrsim 0.1c - 0.3c$) from inner disc streamlines ($R \lesssim 20 R_{\rm in}$, where $R_{\rm in} = 6R_{\rm g} = 6GM/c^2$) with distinct radial velocity stratification. Owing to high magnetization and low mass loading, the dimensionless magnetic lever arm reaches $\tilde{l} = (R_{\rm A}/R)^2 \sim 10 - 100$, allowing an outflow mass-loss rate of $\dot{M}_{\rm w} \simeq 0.14 \dot{M}_{\rm acc}$ to remove roughly 55% of the disc angular momentum ($J_{\rm w} \simeq 0.55 J_{\rm acc}$).	4 Results
arXiv:2610.33314 2026-10-26	**Neutron Star--Primordial Black Hole Binaries: Dynamical Tides, Resonance Locking, and Gravitational Waves** *Neutron Star--Primordial Black Hole Binaries: Dynamical Tides, Resonance Locking, and Gravitational Waves* Isaac Legred, Nicolás Yunes	Resonance locking modifies the orbital evolution according to $\frac{\dot{\Omega}_{\rm orb}}{\dot{\Omega}_{\rm orb}^{\rm GW}} = \frac{I_{\rm orb}}{I_{\rm orb} - 3I_{\rm NS}}$. If $I_{\rm orb} > 3I_{\rm NS}$ and $\Omega_{\rm orb} > \Omega_{\rm NS}$, tidal torques accelerate inspiral; conversely, when $I_{\rm orb} < 3I_{\rm NS}$ and $\Omega_{\rm NS} > \Omega_{\rm orb}$, angular momentum extracted from the NS spin drives an outspiral ($\dot{\Omega}_{\rm orb} < 0$), preventing coalescence. During resonance locking, the pulsar braking index shifts to $n = \frac{11 I_{\rm orb} - 21 I_{\rm NS}}{3(I_{\rm orb} - 3I_{\rm NS})} \frac{\Omega_{\rm NS}}{\Omega_{\rm orb}}$, yielding $n \to \frac{11}{3}\frac{\Omega_{\rm NS}}{\Omega_{\rm orb}}$ in the limit $I_{\rm orb} \gg 3I_{\rm NS}$. Gravitational-wave detectability exhibits a weak scaling ${\rm SNR} \propto \eta^{1/2}$, enabling Cosmic Explorer to detect $M_{\rm BH} > 10^{-6}\,M_\odot$ at $10\,{\rm Mpc}$ (${\rm SNR} > 10$).	Neutron Star–Primordial Black Hole Binaries: Dynamical Tides, Resonance Locking, and Gravitational Waves
arXiv:2610.33297 2026-10-26	**Initial Conditions of Magnetized AGN Accretion Disks from Cloud Capture and Disruption** *Initial Conditions of Magnetized AGN Accretion Disks from Cloud Capture and Disruption* Philip F. Hopkins	The initial disk formed at the circularization radius r_t is toroidally magnetically dominated with $\beta \ll 1$ and $|B_\phi| \gg |B_z|, |B_R|$. The disk scale height and accretion rate scale as $H_d/r_t \sim 4\alpha_{\rm cl,B}$ and $\dot{m} \sim 7\alpha_{\rm cl,B}(\Sigma_{\rm cl}/100\,M_\odot\,{\rm pc^{-2}})^{3/4}(M_{\rm cl}/10^7\,M_\odot)^{-1/4}(2/\tilde{\alpha})$ for $M_{\rm BH} \sim M_{\rm cl}$, where $\alpha_{\rm cl,B} \equiv E_{\rm mag}/E_{\rm grav}$. For $M_{\rm BH} \gg M_{\rm cl}$, the tidal disruption gives $r_t \approx \xi (M_{\rm BH}/M_{\rm cl})^{1/3} R_{\rm cl}$ and $H_d/r_t \sim (3\alpha_{\rm cl,B})^{1/4}(M_{\rm BH}/M_{\rm cl})^{-1/4}$. In the post-encounter cutoff phase, the mass fallback follows a tidal-disruption-like power law $\dot{M} \propto t^{-5/3}$.	8 Conclusions
arXiv:2610.33163 2026-10-26	**Wolf-Rayet Stars in M33: The Impact of Stellar Crowding on Spectral Classification and Mass-Loss Diagnostics** *Wolf-Rayet Stars in M33: The Impact of Stellar Crowding on Spectral Classification and Mass-Loss Diagnostics* Rohit Raj, Puragra Guhathakurta, Robert E. Williams 等	In M33, approximately 27% (33/123) of WR stars suffer moderate-to-severe continuum dilution ($F_{\rm WR} < 0.85$, with very crowded sources having $F_{\rm WR} \lesssim 0.50$ and optical equivalent widths diluted by a factor of $\gtrsim 2$). Because optically thin recombination line luminosities scale as $L_{\rm line} \propto \dot{M}^2$, uncorrected dilution severely underestimates or distorts $\dot{M}$. Keck/DEIMOS spectroscopy confirms the predicted dilution ($\mathrm{EW_{Hecto}/EW_{DEIMOS}} = 0.55$–$0.90$) and yields wind velocity diagnostics of $v_{\rm FWHM} \approx 330$–$1025\ {\rm km\ s^{-1}}$ for He II and blue absorption edge velocities $v_{\rm blue} \approx 700$–$1500\ {\rm km\ s^{-1}}$ for He I $\lambda5876$.	V.2 Implications for Wind Diagnostics and Mass-Loss Rates
arXiv:2610.33161 2026-10-26	**SN Ares: A Strongly Lensed Type IIb Supernova at $\mathbf{z=1.276}$ with a Predicted 70-Year Time Delay** *SN Ares: A Strongly Lensed Type IIb Supernova at $\mathbf{z=1.276}$ with a Predicted 70-Year Time Delay* Conor Larison, Justin D. R. Pierel, Matthew Siebert 等	SN Ares is spectroscopically classified as a Type IIb core-collapse supernova at $z_{\rm spec} = 1.276 \pm 0.002$ magnified by $\mu \approx 4.6 - 5.6$. Correcting for magnification (gain ≈ 1.9 mag), its inferred intrinsic peak absolute magnitude is $M_{B,\rm max} \approx -17.1$ to -17.9 mag, consistent with binary progenitor systems like SN 1993J. Strong lensing modeling predicts two trailing images appearing $\sim 70 - 76$ yr and $\sim 73 - 79$ yr later.	VI Discussion & Conclusions
arXiv:2610.32935 2026-10-25	**Three subpopulations of binary black holes in mass, spin magnitudes and tilts following GWTC-5** *Three subpopulations of binary black holes in mass, spin magnitudes and tilts following GWTC-5* Asad Hussain, Maximiliano Isi, Aaron Zimmerman	GWTC-5 样本支持存在三个 BBH 亚群：(1) **低自旋亚群（A）**：在 $m_1 \approx 35\,M_\odot$ 占主导（$\eta_A(35\,M_\odot) = 0.8 \pm 0.1$），且支持零自旋假设（$\log_{10}\mathcal{B} = 6.1^{+0.2}_{-0.3}$）；(2) **中等自旋亚群（B）**：局限于低质量端 $m_1 \lesssim 20\,M_\odot$（在 $10\,M_\odot$ 处 $\eta_B \approx 0.4$），双黑洞具有适度自旋 $\chi_i \approx 0.2$ 且自旋倾向于轨道平行对齐；(3) **高自旋亚群（C）**：在对不稳定隙附近及以上占主导，主黑洞自旋 $\chi_1 = 0.7^{+0.3}_{-0.4}$，次黑洞自旋 $\chi_2 = 0.6^{+0.4}_{-0.4}$，倾角分布兼容完全各向同性（$\zeta^C_{\cos\theta} = 0$），转变质量始于 $m_3^\eta = 40^{+10}_{-10}\,M_\odot$ 并在 $m_4^\eta = 60^{+40}_{-20}\,M_\odot$ 达到高自旋占优，与对不稳定超新星及分级动力学并合物理预期一致。	III.3 High-spin population