

# Simultaneous Conformal Prediction of Missing Outcomes with Propensity Score $\varepsilon$ -Discretization

Edgar Dobriban

Department of Statistics and Data Science, the Wharton School, University of Pennsylvania

August 3, 2024



## Collaborators



Yonghoon Lee



Eric Tchetgen Tchetgen

# Table of Contents

Introduction and Overview

Our Methods

Illustrating Our Methods in a Stylized Problem  
A Stronger Guarantee


Empirical illustration

# Introduction: Conformal prediction

- Major developing area in statistics: distribution-free predictive inference (a.k.a. conformal prediction)

# Introduction: Conformal prediction


- Major developing area in statistics: distribution-free predictive inference (a.k.a. conformal prediction)
- Goal, given  $(X_1, Y_1), \dots, (X_n, Y_n)$ , find a prediction set  $C$  such that for new  $X_{n+1}$ ,  $\mathbb{P}\{Y_{n+1} \in C(X_{n+1})\} \geq 1 - \alpha$  under *minimal assumptions*




**Figure:** Towards DS

# Introduction: Conformal prediction

- Major developing area in statistics: distribution-free predictive inference (a.k.a. conformal prediction)
- Goal, given  $(X_1, Y_1), \dots, (X_n, Y_n)$ , find a prediction set  $C$  such that for new  $X_{n+1}$ ,  $\mathbb{P}\{Y_{n+1} \in C(X_{n+1})\} \geq 1 - \alpha$  under *minimal assumptions*



**Figure:** Towards DS



- Motivated by complex applications, e.g., where a machine learning model  $\hat{\mu}$  is used to predict  $Y_{n+1}$  based on  $X_{n+1}$  (not known how to find distribution of  $Y_{n+1} - \hat{\mu}(X_{n+1})$ )

# Introduction: Conformal prediction

- It is known how to achieve this in many settings, due to extensive work by many, starting in the 90s (Vovk, Wasserman, J. Lei, R. J. Tibshirani, Barber, Candes, ... )
- Ideas date back to work on tolerance regions by Wilks, Wald, Tukey ... starting in the 1940s



Samuel S. Wilks



Abraham Wald



Vladimir Vovk

## Conformal prediction ctd.

- Typical setting: *exchangeable datapoints*.
  - For a given nonconformity score  $s : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbb{R}$ , e.g.,  $s(x, y) := |y - \hat{\mu}(x)|$ ,  $s(X_1, Y_1), \dots, s(X_{n+1}, Y_{n+1})$  are exchangeable (if  $\hat{\mu}$  is pre-trained on an indep. dataset—i.e., in split conformal prediction)



## Conformal prediction ctd.



- Typical setting: *exchangeable datapoints*.
  - For a given nonconformity score  $s : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbb{R}$ , e.g.,  $s(x, y) := |y - \hat{\mu}(x)|$ ,  $s(X_1, Y_1), \dots, s(X_{n+1}, Y_{n+1})$  are exchangeable (if  $\hat{\mu}$  is pre-trained on an indep. dataset—i.e., in split conformal prediction)
  - Hence, the rank of  $s(X_{n+1}, Y_{n+1})$  is uniform over  $1, \dots, n + 1$  (if no ties)

## Conformal prediction ctd.

- Typical setting: *exchangeable datapoints*.
  - For a given nonconformity score  $s : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbb{R}$ , e.g.,  $s(x, y) := |y - \hat{\mu}(x)|$ ,  $s(X_1, Y_1), \dots, s(X_{n+1}, Y_{n+1})$  are exchangeable (if  $\hat{\mu}$  is pre-trained on an indep. dataset—i.e., in split conformal prediction)
  - Hence, the rank of  $s(X_{n+1}, Y_{n+1})$  is uniform over  $1, \dots, n+1$  (if no ties)
  - So  $x \mapsto C(x) = \{y : \text{rank}\{s(x, y) : s_1, \dots, s_n\} \leq \lceil (1 - \alpha)(n + 1) \rceil\}$  satisfies  $\mathbb{P}\{Y_{n+1} \in C(X_{n+1})\} \geq 1 - \alpha$



# Conformal prediction ctd.

- Typical setting: *exchangeable datapoints*.
  - For a given nonconformity score  $s : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbb{R}$ , e.g.,  $s(x, y) := |y - \hat{\mu}(x)|$ ,  $s(X_1, Y_1), \dots, s(X_{n+1}, Y_{n+1})$  are exchangeable (if  $\hat{\mu}$  is pre-trained on an indep. dataset—i.e., in split conformal prediction)
  - Hence, the rank of  $s(X_{n+1}, Y_{n+1})$  is uniform over  $1, \dots, n+1$  (if no ties)
  - So  $x \mapsto C(x) = \{y : \text{rank}\{s(x, y) : s_1, \dots, s_n\} \leq \lceil (1 - \alpha)(n + 1) \rceil\}$  satisfies  $\mathbb{P}\{Y_{n+1} \in C(X_{n+1})\} \geq 1 - \alpha$



# Conformal prediction ctd.

- Typical setting: *exchangeable datapoints*.
  - For a given nonconformity score  $s : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbb{R}$ , e.g.,  $s(x, y) := |y - \hat{\mu}(x)|$ ,  $s(X_1, Y_1), \dots, s(X_{n+1}, Y_{n+1})$  are exchangeable (if  $\hat{\mu}$  is pre-trained on an indep. dataset—i.e., in split conformal prediction)
  - Hence, the rank of  $s(X_{n+1}, Y_{n+1})$  is uniform over  $1, \dots, n+1$  (if no ties)
  - So  $x \mapsto C(x) = \{y : \text{rank}\{s(x, y) : s_1, \dots, s_n\} \leq \lceil (1 - \alpha)(n + 1) \rceil\}$  satisfies  $\mathbb{P}\{Y_{n+1} \in C(X_{n+1})\} \geq 1 - \alpha$



- However, there are scenarios that existing methods do not resolve, e.g., missing data


# Our problem setting

- Given data

$$(X_1, A_1, Y_1 A_1), \dots, (X_n, A_n, Y_n A_n) \stackrel{\text{iid}}{\sim} P_X \times P_{A|X} \times P_{Y|X},$$

with outcomes *missing at random* (MAR). Thus,

$A_i = 1$  :  $Y_i$  is observed,       $A_i = 0$  :  $Y_i$  is unobserved.




## Our problem setting: Missing At Random

- **Goal:** Simultaneous inference on the missing outcomes  $\{Y_i : A_i = 0\}$ .

# Our problem setting: Missing At Random

- **Goal:** Simultaneous inference on the missing outcomes  $\{Y_i : A_i = 0\}$ .
- Specifically, construct prediction sets  $\{\hat{C}(X_i) : A_i = 0\}$  for  $\{Y_i : A_i = 0\}$  with coverage guarantees



## Inferential target

- With i.i.d./exchangeable data  $(X_1, Y_1), \dots, (X_n, Y_n)$  and test input  $X_{n+1}$ , standard conformal prediction gives a prediction set  $\hat{C}_n(X_{n+1})$  with *marginal coverage*

$$\mathbb{P} \left\{ Y_{n+1} \in \hat{C}_n(X_{n+1}) \right\} \geq 1 - \alpha.$$



## Inferential target

- With i.i.d./exchangeable data  $(X_1, Y_1), \dots, (X_n, Y_n)$  and test input  $X_{n+1}$ , standard conformal prediction gives a prediction set  $\hat{C}_n(X_{n+1})$  with *marginal coverage*

$$\mathbb{P} \left\{ Y_{n+1} \in \hat{C}_n(X_{n+1}) \right\} \geq 1 - \alpha.$$

- **Question:** Under MAR:
  - In what sense can we do useful distribution-free inference for multiple unobserved outcomes?

# Inferential target

- With i.i.d./exchangeable data  $(X_1, Y_1), \dots, (X_n, Y_n)$  and test input  $X_{n+1}$ , standard conformal prediction gives a prediction set  $\hat{C}_n(X_{n+1})$  with *marginal coverage*

$$\mathbb{P} \left\{ Y_{n+1} \in \hat{C}_n(X_{n+1}) \right\} \geq 1 - \alpha.$$

- **Question:** Under MAR:
  - In what sense can we do useful distribution-free inference for multiple unobserved outcomes?
  - Is it possible to go beyond marginal coverage? E.g., have coverage conditional on the test inputs/feature observations with missing outcomes?

# Overview of results

- We consider coverage guarantees of the form

$$\mathbb{E} \left[ \frac{1}{N^{(0)}} \sum_{i:A_i=0} \mathbb{1} \left\{ Y_i \in \widehat{C}(X_i) \right\} \mid X_{1:n}, A_{1:n} \right] \geq 1 - \alpha, \quad (1)$$

where  $N^{(0)}$  is the number of unobserved labels, and  $0/0 := 1$ .

- The proportion of covered missing outcomes is on average at least  $1 - \alpha$ , conditional on  $X_{1:n}$  and the missingness pattern  $A_{1:n}$ .

# Overview of results

- We consider coverage guarantees of the form

$$\mathbb{E} \left[ \frac{1}{N^{(0)}} \sum_{i:A_i=0} \mathbb{1} \left\{ Y_i \in \widehat{C}(X_i) \right\} \mid X_{1:n}, A_{1:n} \right] \geq 1 - \alpha, \quad (1)$$

where  $N^{(0)}$  is the number of unobserved labels, and  $0/0 := 1$ .

- The proportion of covered missing outcomes is on average at least  $1 - \alpha$ , conditional on  $X_{1:n}$  and the missingness pattern  $A_{1:n}$ .
  - For discrete features  $X$ , we construct a procedure that achieves (1).

# Overview of results

- We consider coverage guarantees of the form

$$\mathbb{E} \left[ \frac{1}{N^{(0)}} \sum_{i:A_i=0} \mathbb{1} \left\{ Y_i \in \widehat{C}(X_i) \right\} \mid X_{1:n}, A_{1:n} \right] \geq 1 - \alpha, \quad (1)$$

where  $N^{(0)}$  is the number of unobserved labels, and  $0/0 := 1$ .

- The proportion of covered missing outcomes is on average at least  $1 - \alpha$ , conditional on  $X_{1:n}$  and the missingness pattern  $A_{1:n}$ .
  - For discrete features  $X$ , we construct a procedure that achieves (1).
  - For general features  $X$ , we prove an impossibility result for (1); and then relax it.

## Overview of results - continued

- As a relaxation, we consider

$$\mathbb{E} \left[ \frac{1}{N^{(0)}} \sum_{i:A_i=0} \mathbb{1} \left\{ Y_i \in \widehat{C}(X_i) \right\} \mid B_{1:n}, A_{1:n} \right] \geq 1 - \alpha, \quad (2)$$

where  $B_i = B_i(X_i)$  is a discretization of  $X_i$  (defined soon).

## Overview of results - continued

- As a relaxation, we consider

$$\mathbb{E} \left[ \frac{1}{N^{(0)}} \sum_{i:A_i=0} \mathbb{1} \left\{ Y_i \in \widehat{C}(X_i) \right\} \mid B_{1:n}, A_{1:n} \right] \geq 1 - \alpha, \quad (2)$$

where  $B_i = B_i(X_i)$  is a discretization of  $X_i$  (defined soon).

- Challenge:** Even though we have MAR ( $Y \perp\!\!\!\perp A \mid X$ ), this does not need to be preserved after discretization (may have  $Y \not\perp\!\!\!\perp A \mid B$  for  $B = B(X)$ ).

## Overview of results - continued

- As a relaxation, we consider

$$\mathbb{E} \left[ \frac{1}{N^{(0)}} \sum_{i:A_i=0} \mathbb{1} \left\{ Y_i \in \widehat{C}(X_i) \right\} \mid B_{1:n}, A_{1:n} \right] \geq 1 - \alpha, \quad (2)$$

where  $B_i = B_i(X_i)$  is a discretization of  $X_i$  (defined soon).

- Challenge:** Even though we have MAR ( $Y \perp\!\!\!\perp A \mid X$ ), this does not need to be preserved after discretization (may have  $Y \not\perp\!\!\!\perp A \mid B$  for  $B = B(X)$ ).
- We introduce a carefully designed **propensity score partitioning scheme**, and show how it can be used to obtain (2) in a distribution-free sense (for any dist. of  $(X, Y)$ ).



# Table of Contents

Introduction and Overview


Our Methods

Illustrating Our Methods in a Stylized Problem  
A Stronger Guarantee

Empirical illustration


# First case: Discrete features

- Discrete features naturally form groups of outcomes  $\{Y_i : X_i = x\}$ ,  $x \in \mathcal{X}$ .



## First case: Discrete features


- Discrete features naturally form groups of outcomes  $\{Y_i : X_i = x\}$ ,  $x \in \mathcal{X}$ .



- Within each group, the outcomes are *exchangeable* conditional on  $X_i = x$ .


# Procedure for discrete features: Naive approach

- Direct method: run split conformal prediction separately for each  $x$ .



## Procedure for discrete features: Naive approach


- Direct method: run split conformal prediction separately for each  $x$ .



- This method attains  $\mathbb{E} \left[ \frac{1}{N^{(0)}} \sum_{i:A_i=0} \mathbb{1} \left\{ Y_i \in \hat{C}(X_i) \right\} \mid X_{1:n}, A_{1:n} \right] \geq 1 - \alpha.$

## Procedure for discrete features: Naive approach

- Direct method: run split conformal prediction separately for each  $x$ .



- This method attains  $\mathbb{E} \left[ \frac{1}{N^{(0)}} \sum_{i: A_i=0} \mathbb{1} \left\{ Y_i \in \hat{C}(X_i) \right\} \mid X_{1:n}, A_{1:n} \right] \geq 1 - \alpha$ .
- However, it can produce infinite-width prediction sets in small groups with  $\geq \alpha$  missingness.

## Procedure for discrete features: our method

- Alternative method: simultaneous inference across multiple feature values.

## Procedure for discrete features: our method

- Alternative method: simultaneous inference across multiple feature values.
- Let
  1. Nonconformity score  $s : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbb{R}$ , and  $S_i = s(X_i, Y_i)$  if  $A_i = 1$
  2. Distinct  $X$  values observed:  $X'_1, \dots, X'_M$
  3. Indices of datapoints with features equal to  $X'_k$ :  $I_k = \{i \in [n] : X_i = X'_k\}$ ,
  4. Indices partitioned according to unobserved and observed outcomes, resp.:  
 $I_k^0 = \{i \in [n] : X_i = X'_k, A_i = 0\}$ ,  $I_k^1 = \{i \in [n] : X_i = X'_k, A_i = 1\}$ .
  5. Sample sizes  $N_k = |I_k|$ ,  $N_k^0 = |I_k^0|$



## Procedure for discrete features: our method

- Alternative method: simultaneous inference across multiple feature values.
- Let
  1. Nonconformity score  $s : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbb{R}$ , and  $S_i = s(X_i, Y_i)$  if  $A_i = 1$
  2. Distinct  $X$  values observed:  $X'_1, \dots, X'_M$
  3. Indices of datapoints with features equal to  $X'_k$ :  $I_k = \{i \in [n] : X_i = X'_k\}$ ,
  4. Indices partitioned according to unobserved and observed outcomes, resp.:  
 $I_k^0 = \{i \in [n] : X_i = X'_k, A_i = 0\}$ ,  $I_k^1 = \{i \in [n] : X_i = X'_k, A_i = 1\}$ .
  5. Sample sizes  $N_k = |I_k|$ ,  $N_k^0 = |I_k^0|$
- Our prediction set:

$$\hat{C}(x) = \left\{ y \in \mathcal{Y} : s(x, y) \leq Q_{1-\alpha} \left( \sum_{k=1}^M \sum_{i \in I_k^1} \frac{N_k^0}{N_k N^{(0)}} \delta_{S_i} + \sum_{k=1}^M \frac{(N_k^0)^2}{N_k N^{(0)}} \delta_{+\infty} \right) \right\}. \quad (3)$$

## Procedure for discrete features: our method

- Alternative method: simultaneous inference across multiple feature values.
- Let
  1. Nonconformity score  $s : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbb{R}$ , and  $S_i = s(X_i, Y_i)$  if  $A_i = 1$
  2. Distinct  $X$  values observed:  $X'_1, \dots, X'_M$
  3. Indices of datapoints with features equal to  $X'_k$ :  $I_k = \{i \in [n] : X_i = X'_k\}$ ,
  4. Indices partitioned according to unobserved and observed outcomes, resp.:  
 $I_k^0 = \{i \in [n] : X_i = X'_k, A_i = 0\}$ ,  $I_k^1 = \{i \in [n] : X_i = X'_k, A_i = 1\}$ .
  5. Sample sizes  $N_k = |I_k|$ ,  $N_k^0 = |I_k^0|$
- Our prediction set:

$$\hat{C}(x) = \left\{ y \in \mathcal{Y} : s(x, y) \leq Q_{1-\alpha} \left( \sum_{k=1}^M \sum_{i \in I_k^1} \frac{N_k^0}{N_k N^{(0)}} \delta_{S_i} + \sum_{k=1}^M \frac{(N_k^0)^2}{N_k N^{(0)}} \delta_{+\infty} \right) \right\}. \quad (3)$$

- Idea: symmetry of data distribution; see also *SymmPI* (D. & Yu, 2023)
- Provides uniform-width prediction sets for all  $x$  values.

# Procedure for discrete features: guarantee

## Theorem 1

The prediction set (3) satisfies *feature- and missingness-conditional coverage*

$$\mathbb{E} \left[ \frac{1}{N^{(0)}} \sum_{i:A_i=0} \mathbb{1} \left\{ Y_i \in \hat{C}(X_i) \right\} \mid X_{1:n}, A_{1:n} \right] \geq 1 - \alpha.$$

## Discrete features: improvement via partitioning

- If missingness proportion is high, this can still be conservative.

## Discrete features: improvement via partitioning

- If missingness proportion is high, this can still be conservative.
- Idea: *Partition* datapoints. For each partition, use pro-CP on *all datapoints with observed labels* to predict outcomes missing in that partition.

## Discrete features: improvement via partitioning

- If missingness proportion is high, this can still be conservative.
- Idea: *Partition* datapoints. For each partition, use pro-CP on *all datapoints with observed labels* to predict outcomes missing in that partition.
- Since guarantee is feature- and missingness-conditional, this is still valid!

## Discrete features: improvement via partitioning

- If missingness proportion is high, this can still be conservative.
- Idea: *Partition* datapoints. For each partition, use pro-CP on *all datapoints with observed labels* to predict outcomes missing in that partition.
- Since guarantee is feature- and missingness-conditional, this is still valid!
- Previous methods are at two endpoints: partition is all singletons (“naive method”) vs whole set (“our method”).

## Discrete features: improvement via partitioning

- If missingness proportion is high, this can still be conservative.
- Idea: *Partition* datapoints. For each partition, use pro-CP on *all datapoints with observed labels* to predict outcomes missing in that partition.
- Since guarantee is feature- and missingness-conditional, this is still valid!
- Previous methods are at two endpoints: partition is all singletons (“naive method”) vs whole set (“our method”).
- Why practically useful? Partition can depend on  $X_{1:n}, A_{1:n}$ ; can aim to ensure small missingness per group.




## Procedure for general feature distributions

- If the propensity score  $x \mapsto p_{A|X}(x) = \mathbb{P}\{A = 1 \mid X = x\}$  is known,  $\varepsilon$ -**discretize** it
- Let  $\varepsilon$  be a predefined discretization level, and  $z_k = (1 + \varepsilon)^k / [1 + (1 + \varepsilon)^k]$  for all integers  $k$


## Procedure for general feature distributions

- If the propensity score  $x \mapsto p_{A|X}(x) = \mathbb{P}\{A = 1 \mid X = x\}$  is known,  **$\varepsilon$ -discretize** it
- Let  $\varepsilon$  be a predefined discretization level, and  $z_k = (1 + \varepsilon)^k / [1 + (1 + \varepsilon)^k]$  for all integers  $k$
- Partition the feature space into  $D_k = \{x : p_{A|X}(x) \in [z_k, z_{k+1})\}$ ,  $\mathcal{B} = \{D_k : k \in \mathbb{Z}\}$ .



## Procedure for general feature distributions

- If the propensity score  $x \mapsto p_{A|X}(x) = \mathbb{P}\{A = 1 \mid X = x\}$  is known,  **$\varepsilon$ -discretize** it
- Let  $\varepsilon$  be a predefined discretization level, and  $z_k = (1 + \varepsilon)^k / [1 + (1 + \varepsilon)^k]$  for all integers  $k$
- Partition the feature space into  $D_k = \{x : p_{A|X}(x) \in [z_k, z_{k+1})\}$ ,  $\mathcal{B} = \{D_k : k \in \mathbb{Z}\}$ .



## Pro-CP

- Show *approximate within-partition exchangeability* of the scores, enabling inference.

## Pro-CP

- Show *approximate within-partition exchangeability* of the scores, enabling inference.
- **Propensity score discretization-based conformal prediction (pro-CP)**: Procedure (4) applied to the discretized data  $(B_i, A_i, A_i Y_i)_{i \in [n]}$ , i.e.,

$$\hat{C}^{\text{pro-CP}}(x) = \left\{ y \in \mathcal{Y}, \text{ : } s(x, y) \leq Q_{1-\alpha} \left( \sum_{k=1}^M \sum_{i \in I_k^{\mathcal{B},1}} \frac{N_k^{\mathcal{B},0}}{N^{(0)} N_k^{\mathcal{B}}} \cdot \delta_{S_i} + \frac{1}{N^{(0)}} \sum_{k=1}^M \frac{(N_k^{\mathcal{B},0})^2}{N_k^{\mathcal{B}}} \cdot \delta_{+\infty} \right) \right\}. \quad (4)$$

## Pro-CP

- Show *approximate within-partition exchangeability* of the scores, enabling inference.
- **Propensity score discretization-based conformal prediction (pro-CP)**: Procedure (4) applied to the discretized data  $(B_i, A_i, A_i Y_i)_{i \in [n]}$ , i.e.,

$$\hat{C}^{\text{pro-CP}}(x) = \left\{ y \in \mathcal{Y}, :, s(x, y) \leq Q_{1-\alpha} \left( \sum_{k=1}^M \sum_{i \in I_k^{\mathcal{B},1}} \frac{N_k^{\mathcal{B},0}}{N^{(0)} N_k^{\mathcal{B}}} \cdot \delta_{S_i} + \frac{1}{N^{(0)}} \sum_{k=1}^M \frac{(N_k^{\mathcal{B},0})^2}{N_k^{\mathcal{B}}} \cdot \delta_{+\infty} \right) \right\}. \quad (4)$$

### Theorem 2

Suppose  $0 < p_{A|X}(X) < 1$  almost surely. Then  $\hat{C}^{\text{pro-CP}}$  from (4) satisfies *propensity score discretized feature- and missingness-conditional coverage*:

$$\mathbb{E} \left[ \frac{1}{N^{(0)}} \sum_{i: A_i=0} \mathbb{1} \left\{ Y_i \in \hat{C}^{\text{pro-CP}}(X_i) \right\} \mid B_{1:n}, A_{1:n} \right] \geq 1 - \alpha - \varepsilon.$$

## Pro-CP

- Show *approximate within-partition exchangeability* of the scores, enabling inference.
- **Propensity score discretization-based conformal prediction (pro-CP)**: Procedure (4) applied to the discretized data  $(B_i, A_i, A_i Y_i)_{i \in [n]}$ , i.e.,

$$\hat{C}^{\text{pro-CP}}(x) = \left\{ y \in \mathcal{Y}, :, s(x, y) \leq Q_{1-\alpha} \left( \sum_{k=1}^M \sum_{i \in I_k^{\mathcal{B},1}} \frac{N_k^{\mathcal{B},0}}{N^{(0)} N_k^{\mathcal{B}}} \cdot \delta_{S_i} + \frac{1}{N^{(0)}} \sum_{k=1}^M \frac{(N_k^{\mathcal{B},0})^2}{N_k^{\mathcal{B}}} \cdot \delta_{+\infty} \right) \right\}. \quad (4)$$

### Theorem 2

Suppose  $0 < p_{A|X}(X) < 1$  almost surely. Then  $\hat{C}^{\text{pro-CP}}$  from (4) satisfies *propensity score discretized feature- and missingness-conditional coverage*:

$$\mathbb{E} \left[ \frac{1}{N^{(0)}} \sum_{i: A_i=0} \mathbb{1} \left\{ Y_i \in \hat{C}^{\text{pro-CP}}(X_i) \right\} \mid B_{1:n}, A_{1:n} \right] \geq 1 - \alpha - \varepsilon.$$

- The error from discretization is bounded by  $\varepsilon$ , for *any*  $n$  and  $\#$  of missing outcomes.

## Pro-CP with estimated propensity score

- If the propensity score is unknown, we may run pro-CP with an estimator  $\hat{p}_{A|X}$  of  $p_{A|X}$ .



## Pro-CP with estimated propensity score

- If the propensity score is unknown, we may run pro-CP with an estimator  $\hat{p}_{A|X}$  of  $p_{A|X}$ .

### Theorem 3

Suppose  $0 < p_{A|X}(X) < 1$  and  $0 < \hat{p}_{A|X}(X) < 1$  almost surely. Then pro-CP run with  $\hat{p}_{A|X}$  satisfies

$$\mathbb{E} \left[ \frac{1}{N^{(0)}} \sum_{i:A_i=0} \mathbb{1} \left\{ Y_i \in \hat{C}^{\text{pro-CP}}(X_i) \right\} \mid B_{1:n}, A_{1:n} \right] \geq 1 - \alpha - (\varepsilon + \delta_{\hat{p}_{A|X}} + \varepsilon \delta_{\hat{p}_{A|X}}),$$

where

$$\delta_{\hat{p}_{A|X}} = e^{2\|\log f_{p,\hat{p}}\|_\infty} - 1, \quad f_{p,\hat{p}}(x) = \frac{p_{A|X}(x)/(1-p_{A|X}(x))}{\hat{p}_{A|X}(x)/(1-\hat{p}_{A|X}(x))}.$$

## Pro-CP with estimated propensity score

- If the propensity score is unknown, we may run pro-CP with an estimator  $\hat{p}_{A|X}$  of  $p_{A|X}$ .

### Theorem 3

Suppose  $0 < p_{A|X}(X) < 1$  and  $0 < \hat{p}_{A|X}(X) < 1$  almost surely. Then pro-CP run with  $\hat{p}_{A|X}$  satisfies

$$\mathbb{E} \left[ \frac{1}{N^{(0)}} \sum_{i:A_i=0} \mathbb{1} \left\{ Y_i \in \hat{C}^{\text{pro-CP}}(X_i) \right\} \mid B_{1:n}, A_{1:n} \right] \geq 1 - \alpha - (\varepsilon + \delta_{\hat{p}_{A|X}} + \varepsilon \delta_{\hat{p}_{A|X}}),$$

where

$$\delta_{\hat{p}_{A|X}} = e^{2\|\log f_{p,\hat{p}}\|_\infty} - 1, \quad f_{p,\hat{p}}(x) = \frac{p_{A|X}(x)/(1-p_{A|X}(x))}{\hat{p}_{A|X}(x)/(1-\hat{p}_{A|X}(x))}.$$

- The error from estimation does not grow with the number of missing outcomes.

## New result underlying pro-CP guarantee

- Balancing property of the propensity score [Rosenbaum and Rubin (1983)]: the missingness is independent of the outcome conditional on the propensity:  $A \perp\!\!\!\perp Y \mid p_{A|X}$ .

## New result underlying pro-CP guarantee

- Balancing property of the propensity score [Rosenbaum and Rubin (1983)]: the missingness is independent of the outcome conditional on the propensity:  $A \perp\!\!\!\perp Y \mid p_{A|X}$ .
- We show *approximate version*: dist. of  $s(X, Y)$  close for  $A = 0, 1$  given small range of  $p_{A|X}$

Lemma (Bounded prop. score implies closeness of cond. distrib. for obs. and missing)

Suppose that  $(X, Y, A) \sim P_X \times P_{Y|X} \times \text{Bernoulli}(p_{A|X})$  on  $\mathcal{X} \times \mathcal{Y} \times \{0, 1\}$ , and that for a set  $B \subset \mathcal{X}$  and  $t \in (0, 1)$ ,  $\varepsilon \geq 0$ ,

$$t \leq \frac{p_{A|X}(x)}{1 - p_{A|X}(x)} \leq t(1 + \varepsilon), \text{ for all } x \in B.$$

Let  $s : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbb{R}$  be any measurable function and let  $S = s(X, Y)$ . Then

$$d_{TV}(P_{S|A=1, X \in B}, P_{S|A=0, X \in B}) \leq \varepsilon.$$

# Table of Contents

Introduction and Overview

Our Methods

Illustrating Our Methods in a Stylized Problem

A Stronger Guarantee

Empirical illustration

# Application to simultaneous inference on ITEs

- Consider a potential outcomes model

$$(X_i, T_i, Y_i(0), Y_i(1))_{1 \leq i \leq n} \stackrel{\text{iid}}{\sim} P_X \times P_{T|X} \times P_{Y(1)|X} \times P_{Y(0)|X},$$

where we observe  $(X_i, T_i, T_i Y_i(1) + (1 - T_i) Y_i(0))_{1 \leq i \leq n}$ .

## Application to simultaneous inference on ITEs

- Consider a potential outcomes model

$$(X_i, T_i, Y_i(0), Y_i(1))_{1 \leq i \leq n} \stackrel{\text{iid}}{\sim} P_X \times P_{T|X} \times P_{Y(1)|X} \times P_{Y(0)|X},$$

where we observe  $(X_i, T_i, T_i Y_i(1) + (1 - T_i) Y_i(0))_{1 \leq i \leq n}$ .

- Applying pro-CP, we can construct  $\hat{C}^{\text{counterfactual}}$  such that

$$\mathbb{E} \left[ \frac{1}{N^{(0)}} \sum_{i: T_i=0} \mathbb{1} \left\{ Y_i(1) \in \hat{C}^{\text{counterfactual}}(X_i) \right\} \mid B_{1:n}, T_{1:n} \right] \geq 1 - \alpha.$$

## Application to simultaneous inference on ITEs

- Consider a potential outcomes model

$$(X_i, T_i, Y_i(0), Y_i(1))_{1 \leq i \leq n} \stackrel{\text{iid}}{\sim} P_X \times P_{T|X} \times P_{Y(1)|X} \times P_{Y(0)|X},$$

where we observe  $(X_i, T_i, T_i Y_i(1) + (1 - T_i) Y_i(0))_{1 \leq i \leq n}$ .

- Applying pro-CP, we can construct  $\hat{C}^{\text{counterfactual}}$  such that

$$\mathbb{E} \left[ \frac{1}{N^{(0)}} \sum_{i: T_i=0} \mathbb{1} \left\{ Y_i(1) \in \hat{C}^{\text{counterfactual}}(X_i) \right\} \mid B_{1:n}, T_{1:n} \right] \geq 1 - \alpha.$$

- By letting  $\hat{C}_i^{\text{ITE}} = \{y - Y_i(0) : y \in \hat{C}^{\text{counterfactual}}(X_i)\}$ , we obtain prediction sets for individual treatment effects

$$\mathbb{E} \left[ \frac{1}{N^{(0)}} \sum_{i \in I_{T=0}} \mathbb{1} \left\{ (Y_i(1) - Y_i(0)) \in \hat{C}_i^{\text{ITE}} \right\} \mid B_{1:n}, T_{1:n} \right] \geq 1 - \alpha.$$



# Table of Contents

Introduction and Overview

**Our Methods**

Illustrating Our Methods in a Stylized Problem  
**A Stronger Guarantee**

Empirical illustration

# Achieving a stronger guarantee on the coverage proportion

- Can we achieve a stronger guarantee beyond bounding the *mean coverage*?

# Achieving a stronger guarantee on the coverage proportion

- Can we achieve a stronger guarantee beyond bounding the *mean coverage*?
- Possible goal: PAC-type guarantee of the form

$$\mathbb{P} \left\{ \frac{1}{N^{(0)}} \sum_{i:A_i=0} \mathbb{1} \left\{ Y_i \in \hat{C}^{\text{pro-CP}}(X_i) \right\} \geq 1 - \alpha \right\} \geq 1 - \delta.$$

# Achieving a stronger guarantee on the coverage proportion

- Can we achieve a stronger guarantee beyond bounding the *mean coverage*?
- Possible goal: PAC-type guarantee of the form

$$\mathbb{P} \left\{ \frac{1}{N^{(0)}} \sum_{i:A_i=0} \mathbb{1} \left\{ Y_i \in \hat{C}^{\text{pro-CP}}(X_i) \right\} \geq 1 - \alpha \right\} \geq 1 - \delta.$$

- Turns out to be hard to achieve in the distribution-free sense

# Achieving a stronger guarantee on the coverage proportion

- Can we achieve a stronger guarantee beyond bounding the *mean coverage*?
- Possible goal: PAC-type guarantee of the form

$$\mathbb{P} \left\{ \frac{1}{N^{(0)}} \sum_{i:A_i=0} \mathbb{1} \left\{ Y_i \in \hat{C}^{\text{pro-CP}}(X_i) \right\} \geq 1 - \alpha \right\} \geq 1 - \delta.$$

- Turns out to be hard to achieve in the distribution-free sense
- As a surrogate target, we consider bounding the *squared coverage*

$$\mathbb{E} \left[ \left( \frac{1}{N^{(0)}} \sum_{i:A_i=0} \mathbb{1} \left\{ Y_i \notin \hat{C}^{\text{pro-CP}}(X_i) \right\} \right)^2 \right] \leq \alpha^2.$$

# Achieving a stronger guarantee on the coverage proportion

- Can we achieve a stronger guarantee beyond bounding the *mean coverage*?
- Possible goal: PAC-type guarantee of the form

$$\mathbb{P} \left\{ \frac{1}{N^{(0)}} \sum_{i:A_i=0} \mathbb{1} \left\{ Y_i \in \hat{C}^{\text{pro-CP}}(X_i) \right\} \geq 1 - \alpha \right\} \geq 1 - \delta.$$

- Turns out to be hard to achieve in the distribution-free sense
- As a surrogate target, we consider bounding the *squared coverage*

$$\mathbb{E} \left[ \left( \frac{1}{N^{(0)}} \sum_{i:A_i=0} \mathbb{1} \left\{ Y_i \notin \hat{C}^{\text{pro-CP}}(X_i) \right\} \right)^2 \right] \leq \alpha^2.$$

(motivated by Lee et. al. (2023): Hierarchical CP)

## Interpretation of the squared-coverage guarantee

- Let  $\hat{m} = \frac{1}{N^{(0)}} \sum_{i:A_i=0} \mathbb{1} \left\{ Y_i \in \hat{C}^{\text{pro-CP}}(X_i) \right\}$  denote the *miscoverage proportion*.

# Interpretation of the squared-coverage guarantee

- Let  $\hat{m} = \frac{1}{N^{(0)}} \sum_{i:A_i=0} \mathbb{1} \left\{ Y_i \in \hat{C}^{\text{pro-CP}}(X_i) \right\}$  denote the *miscoverage proportion*.
- Conditional on (discretized) features, pro-CP attains  $\mathbb{E}[\hat{m}] \leq \alpha$ .



# Interpretation of the squared-coverage guarantee

- Let  $\hat{m} = \frac{1}{N^{(0)}} \sum_{i:A_i=0} \mathbb{1} \left\{ Y_i \in \hat{C}^{\text{pro-CP}}(X_i) \right\}$  denote the *miscoverage proportion*.
- Conditional on (discretized) features, pro-CP attains  $\mathbb{E}[\hat{m}] \leq \alpha$ .
- The squared-coverage guarantee is  $\mathbb{E}[\hat{m}^2] \leq \alpha^2$ , and provides a stronger control over  $\hat{m}$  being close to unity, preventing e.g.,  $\hat{m} = 0$  w.p.  $1 - \alpha$  and  $1$  w.p.  $\alpha$ .

## Pro-CP2 procedure

- Define

1. For all  $i \in [n]$ ,  $\bar{S}_i = S_i$  if  $A_i = 1$  and  $\bar{S}_i = +\infty$  if  $A_i = 0$ .
2. Pairwise minima:  $\bar{S}_{ij} := \min\{\bar{S}_i, \bar{S}_j\}$  for all  $i, j$ .

## Pro-CP2 procedure

- Define
  - For all  $i \in [n]$ ,  $\bar{S}_i = S_i$  if  $A_i = 1$  and  $\bar{S}_i = +\infty$  if  $A_i = 0$ .
  - Pairwise minima:  $\bar{S}_{ij} := \min\{\bar{S}_i, \bar{S}_j\}$  for all  $i, j$ .
- Pro-CP2 prediction set:

$$\hat{C}^{\text{pro-CP2}}(x) = \left\{ y \in \mathcal{Y} : s(x, y) \leq Q_{1-\alpha^2} \left( \sum_{k=1}^M \sum_{i \in I_k^{\mathcal{B}}} \frac{1}{(N^{(0)})^2} \cdot \frac{N_k^{\mathcal{B},0}}{N_k^{\mathcal{B}}} \cdot \delta_{\bar{S}_i} \right. \right. \\ \left. \left. + \sum_{k=1}^M \sum_{\substack{i, j \in I_k^{\mathcal{B}} \\ i \neq j}} \frac{N_k^{\mathcal{B},0} (N_k^{\mathcal{B},0} - 1)}{(N^{(0)})^2 N_k^{\mathcal{B}} (N_k^{\mathcal{B}} - 1)} \delta_{\bar{S}_{ij}} + \sum_{1 \leq k \neq k' \leq M} \sum_{i \in I_k^{\mathcal{B}}} \sum_{j \in I_{k'}^{\mathcal{B}}} \frac{N_k^{\mathcal{B},0} N_{k'}^{\mathcal{B},0}}{(N^{(0)})^2 N_k^{\mathcal{B}} N_{k'}^{\mathcal{B}}} \delta_{\bar{S}_{ij}} \right) \right\}.$$

## Pro-CP2 procedure

- Define
  - For all  $i \in [n]$ ,  $\bar{S}_i = S_i$  if  $A_i = 1$  and  $\bar{S}_i = +\infty$  if  $A_i = 0$ .
  - Pairwise minima:  $\bar{S}_{ij} := \min\{\bar{S}_i, \bar{S}_j\}$  for all  $i, j$ .
- Pro-CP2 prediction set:

$$\hat{C}^{\text{pro-CP2}}(x) = \left\{ y \in \mathcal{Y} : s(x, y) \leq Q_{1-\alpha^2} \left( \sum_{k=1}^M \sum_{i \in I_k^{\mathcal{B}}} \frac{1}{(N^{(0)})^2} \cdot \frac{N_k^{\mathcal{B},0}}{N_k^{\mathcal{B}}} \cdot \delta_{\bar{S}_i} \right. \right. \\ \left. \left. + \sum_{k=1}^M \sum_{\substack{i, j \in I_k^{\mathcal{B}} \\ i \neq j}} \frac{N_k^{\mathcal{B},0} (N_k^{\mathcal{B},0} - 1)}{(N^{(0)})^2 N_k^{\mathcal{B}} (N_k^{\mathcal{B}} - 1)} \delta_{\bar{S}_{ij}} + \sum_{1 \leq k \neq k' \leq M} \sum_{i \in I_k^{\mathcal{B}}} \sum_{j \in I_{k'}^{\mathcal{B}}} \frac{N_k^{\mathcal{B},0} N_{k'}^{\mathcal{B},0}}{(N^{(0)})^2 N_k^{\mathcal{B}} N_{k'}^{\mathcal{B}}} \delta_{\bar{S}_{ij}} \right) \right\}.$$

- Similar intuition as before; but use invariance to find probability of  $\mathbb{1} \left\{ \min\{S_{i^*}, S_{j^*}\} \leq q_{1-\alpha^2}(\tilde{S}_1, \dots, \tilde{S}_M) \right\}$ , where  $i^*, j^*$  are random data indices with  $A = 0$ .

# Squared coverage error control of Pro-CP2

## Theorem 4

If  $0 < p_{A|X}(X) < 1$  almost surely, then  $\hat{C}^{\text{pro-CP2}}$  satisfies

$$\mathbb{E} \left[ \left( \frac{1}{N^{(0)}} \sum_{i: A_i=0} \mathbb{1} \left\{ Y_i \notin \hat{C}^{\text{pro-CP2}}(X_i) \right\} \right)^2 \middle| B_{1:n}, A_{1:n} \right] \leq \alpha^2 + 2\varepsilon.$$

# Table of Contents

Introduction and Overview

Our Methods

Illustrating Our Methods in a Stylized Problem  
A Stronger Guarantee

Empirical illustration

# Simulation 1

**Weighted conformal (Tibshirani et al., 2019) vs pro-CP:** marginal vs conditional coverage

# Simulation 1

**Weighted conformal (Tibshirani et al., 2019) vs pro-CP:** marginal vs conditional coverage

1.  $X \sim \text{Unif}[0, 10]$ ,  $Y \mid X \sim N(X, (3 + X)^2)$ ,  $A \mid X \sim \text{Bernoulli}(p_{A|X}(X))$



# Simulation 1

**Weighted conformal (Tibshirani et al., 2019) vs pro-CP:** marginal vs conditional coverage

1.  $X \sim \text{Unif}[0, 10]$ ,  $Y | X \sim N(X, (3 + X)^2)$ ,  $A | X \sim \text{Bernoulli}(p_{A|X}(X))$
2. (1) :  $p_{A|X}(x) = 0.9 - 0.02x$ , (2) :  $p_{A|X}(x) = 0.8 - 0.1(1 + 0.1x) \sin 3x$

# Simulation 1

**Weighted conformal (Tibshirani et al., 2019) vs pro-CP:** marginal vs conditional coverage

1.  $X \sim \text{Unif}[0, 10]$ ,  $Y \mid X \sim N(X, (3 + X)^2)$ ,  $A \mid X \sim \text{Bernoulli}(p_{A|X}(X))$
2. (1) :  $p_{A|X}(x) = 0.9 - 0.02x$ , (2) :  $p_{A|X}(x) = 0.8 - 0.1(1 + 0.1x) \sin 3x$
3. Fit OLS with  $n_{\text{train}} = 500$ ,  $s(x, y) = |y - \hat{\mu}(x)|$

# Simulation 1

**Weighted conformal (Tibshirani et al., 2019) vs pro-CP:** marginal vs conditional coverage

1.  $X \sim \text{Unif}[0, 10]$ ,  $Y \mid X \sim N(X, (3 + X)^2)$ ,  $A \mid X \sim \text{Bernoulli}(p_{A|X}(X))$
2. (1) :  $p_{A|X}(x) = 0.9 - 0.02x$ , (2) :  $p_{A|X}(x) = 0.8 - 0.1(1 + 0.1x) \sin 3x$
3. Fit OLS with  $n_{\text{train}} = 500$ ,  $s(x, y) = |y - \hat{\mu}(x)|$
4. 500 trials,  $n = 500$ , Pro-CP  $\varepsilon = 0.1$ ,  $\alpha = 0.2$ , partition of size 10;

# Simulation 1


**Weighted conformal (Tibshirani et al., 2019) vs pro-CP:** marginal vs conditional coverage

1.  $X \sim \text{Unif}[0, 10]$ ,  $Y \mid X \sim N(X, (3 + X)^2)$ ,  $A \mid X \sim \text{Bernoulli}(p_{A|X}(X))$
2. (1) :  $p_{A|X}(x) = 0.9 - 0.02x$ , (2) :  $p_{A|X}(x) = 0.8 - 0.1(1 + 0.1x) \sin 3x$
3. Fit OLS with  $n_{\text{train}} = 500$ ,  $s(x, y) = |y - \hat{\mu}(x)|$
4. 500 trials,  $n = 500$ , Pro-CP  $\varepsilon = 0.1$ ,  $\alpha = 0.2$ , partition of size 10;
5. Given  $X_{1:n}, A_{1:n}$ , 100x gen  $(X'_i, Y'_i)_{1 \leq i \leq n} \mid B_i \sim P_{X|B} \times P_{Y|X}$ ,  $n = 500$

# Simulation 1

## Weighted conformal (Tibshirani et al., 2019) vs pro-CP: marginal vs conditional coverage


1.  $X \sim \text{Unif}[0, 10]$ ,  $Y | X \sim N(X, (3 + X)^2)$ ,  $A | X \sim \text{Bernoulli}(p_{A|X}(X))$
2. (1) :  $p_{A|X}(x) = 0.9 - 0.02x$ , (2) :  $p_{A|X}(x) = 0.8 - 0.1(1 + 0.1x) \sin 3x$
3. Fit OLS with  $n_{\text{train}} = 500$ ,  $s(x, y) = |y - \hat{\mu}(x)|$
4. 500 trials,  $n = 500$ , Pro-CP  $\varepsilon = 0.1$ ,  $\alpha = 0.2$ , partition of size 10;
5. Given  $X_{1:n}, A_{1:n}$ , 100x gen  $(X'_i, Y'_i)_{1 \leq i \leq n} | B_i \sim P_{X|B} \times P_{Y|X}$ ,  $n = 500$



## Simulation 2


### pro-CP vs pro-CP2: controlling mean vs squared miscoverage

- Same setting as Simulation 1, but evaluate marginal coverage & estimate propensity score with kernel regression on training data




# Illustration on diabetes dataset (Efron et al., 2004)

- $X$ : ten features (age, bmi, LDL/HDL, ...) of patients (sample sizes: train: 142; calibration+test: 300)
- $A$ : missingness generated from a known logistic model
- $Y$ : a measure of disease progression one year after baseline




## Illustration on diabetes dataset (Efron et al., 2004): II





# Illustration on JOBS II dataset (Imai et al., 2010)

- $X$ : job seekers:  $n_{\text{train}} = 379$ ,  $n = 500$ ; with 14 demographic features
- $A$ : job skills workshop (to evaluate our methods, simulate via logistic model; estimate via RF)
- $Y(0)$ ,  $Y(1)$ : pre- and post-treatment depression measure



# Discussion

- Introduced Pro-CP, a method for simultaneous prediction of multiple missing outcomes, and provided coverage guarantees
- Pro-CP2: stronger squared error miscoverage error control
- What applications might this have an impact on? Where could it be used?
- Preprint: [arxiv.org/abs/2403.04613](https://arxiv.org/abs/2403.04613). Code: [github.com/yhoon31/pro-CP](https://github.com/yhoon31/pro-CP)
- Thanks!

