

Are you aware you should NOT USE more than one of the SAME TYPE of topical treatments at the SAME TIME? • Zineryt, Skinoren and Dalacin T - all antibiotics • Differin and Istorex - Both Retinoids • Duac - Antibiotic with peeling agent (benzoyl peroxide)	Yes
Are you aware Lymecycline antibiotic tablets should not be taken whilst using Zineryt, Dalacin T and Duac (topical antibiotics)? • Lymecycline can be taken whilst using Differin or Istorex	No
Are you aware it usually takes 6-8 weeks or more of continuous usage for treatments to work? If there is no improvement after 14 weeks, treatment should be reviewed and changed.	Yes
Are you aware, if acne has cleared-up, treatment can sometimes be stopped or reduced after 4 to 6 months?	No
Are you aware you should see a GP face-to-face if acne is scarring or getting worse?	Yes
Are you aware topical treatments containing retinoids and peeling agents (Differin, Isotrex and Duac) can be irritant?	Yes
Are you aware the following treatments should NOT be used if you are pregnant or breast feeding? •Differin •Isotrex •Duac •Lymecycline •Skinoren Details can be found in the 'patient information leaflets'	Yes
Please indicate your consent to receiving a phone call from the pharmacist to arrange for collection of your prescription	No
Please select your medicine	rererer
Would you like to request same-day collection at your local pharmacy? (For urgent prescriptions only)	Yes

Your Name	ye
Your Email	yest@yest.com
Phone	0909090909
Location	teytyutu