Reactive Attachment Disorder (DSM-5-TR): Teaching Overview

Clinical Companion Knowledge Library: for counselors and counseling students

1. What it is

Reactive attachment disorder (RAD) is a disorder of early childhood in which a child **rarely seeks or responds to comfort** from caregivers and shows limited positive emotion, after **severely insufficient care**: social neglect, repeated changes of caregivers (as in multiple foster placements), or rearing in settings such as institutions where close bonds are nearly impossible (American Psychiatric Association [APA], 2022). It belongs to DSM-5-TR's "Trauma- and Stressor-Related Disorders" chapter.

How common: among all 1,646 children aged 6 to 8 in a deprived urban area of the United Kingdom, an estimated 1.4% had RAD; the authors concluded it is not rare in such populations (Minnis et al., 2013). Children with RAD in the general population often have several other disorders, lower IQ, and poorer social skills (Pritchett et al., 2013).

A distinct disorder. Research supports RAD and disinhibited social engagement disorder as two separate disorders, although both arise from inadequate care (Zeanah & Gleason, 2015).

Counselor framing: RAD is a wound from what a child did not receive, not a sign of a "bad" or manipulative child. Most of the work happens through the caregiver, and many of the families who seek help are foster or adoptive parents.

2. Core symptoms to watch for

DSM-5-TR criteria, paraphrased (APA, 2022); use the manual itself for formal diagnosis.

- A consistent pattern of **inhibited, emotionally withdrawn behavior toward adult caregivers**: the child **rarely or minimally seeks comfort** when distressed and **rarely or minimally responds to comfort** when it is offered.
- Persistent social and emotional disturbance with **at least 2** of: minimal social and emotional responsiveness to others; **limited positive emotion**; episodes of unexplained irritability, sadness, or fearfulness, even during calm, nonthreatening interactions with caregivers.
- A history of **extremes of insufficient care**: persistent lack of comfort, stimulation, and affection; repeated changes of primary caregivers; or rearing in unusual settings that severely limit opportunities to form selective attachments.
- The insufficient care is presumed to cause the behavior. The child has a developmental age of **at least 9 months**, signs are evident **before age 5**, and criteria for autism spectrum disorder are not met. Specify **persistent** (more than 12 months) and **severe** (all symptoms at relatively high levels).

Rule out: autism spectrum disorder (social difficulty without a history of neglect, plus restricted, repetitive behavior), intellectual disability, and depression. Note that "attachment disorder" is often used loosely online and in parenting circles for any difficult foster or adoptive child; the diagnosis requires the specific pattern above (APA, 2022; Chaffin et al., 2006).

Assessment and referral (general clinical consensus and practice parameter): directly observe the child with the primary caregiver (Zeanah et al., 2016). Refer for pediatric, developmental, and psychiatric evaluation; screen for co-occurring problems such as developmental delay, ADHD, and trauma symptoms; and report suspected ongoing abuse or neglect.

3. Frequently used assessment instruments

These are among the most widely used measures for screening, rating severity, and tracking progress in reactive attachment disorder. **Screening tools do not diagnose;** a positive screen calls for a clinical interview against DSM-5-TR criteria. Click an instrument's website to find it. No single test diagnoses RAD; direct observation of the child with the caregiver is essential.

Instrument	Best used for	Format	Purchase required?	Training or qualification	Where to find it
DAI (Disturbances of Attachment Interview)	Caregiver interview rating signs of RAD and DSED; used in the major research studies.	Clinician interview with the caregiver	No (free)	★ Clinical interviewing skill; training recommended	Original article

Instrument	Best used for	Format	Purchase required?	Training or qualification	Where to find it
RPQ (Relationship Problems Questionnaire)	Brief parent or teacher screen for inhibited and disinhibited attachment behaviors.	Parent- or teacher-report, 10 items	No (free)	None beyond general clinical training	Original article
SDQ (Strengths and Difficulties Questionnaire)	Screening for co-occurring emotional, conduct, attention, and peer problems.	Parent, teacher, and self-report, 25 items	No (free paper version)	None beyond general clinical training	SDQ website
BASC-3 (Behavior Assessment System for Children)	Broad behavioral and emotional assessment, ages 2 to 21.	Parent, teacher, and self-report forms	Yes	★ Level B	Pearson
TSCYC (Trauma Symptom Checklist for Young Children)	Caregiver report of trauma symptoms in children ages 3 to 12.	Caregiver-report, 90 items (screening form 20)	Yes	★ Level B	PAR
NICHQ Vanderbilt Assessment Scales	Screening for co-occurring ADHD in children ages 6 to 12.	Parent and teacher forms, initial and follow-up	No (free)	None beyond general clinical training	NICHQ (PDF)

★ **= requires professional qualification to purchase or specialized training.** Publishers sell these tests only to qualified users. At Pearson, **Level B** generally means a master's degree in a related field with formal assessment training, or a health care license; **Level C** means a doctorate in psychology or a related field with formal assessment training, or state licensure in a related field (Pearson qualifications policy). MHS, PAR, and WPS use similar levels. Check each publisher's page before ordering. "Original article" links go to the published study that introduced a free scale that has no official website; the scale can usually be obtained there or from its authors.

4. What works: treatment and the evidence

Bottom line for class: the core treatment is a stable, sensitive, consistent caregiver. Caregiver-focused interventions that build sensitivity have the best support. Coercive "attachment therapies" should be avoided.

Evidence overview (honest effect sizes):

- **A family heals:** in a randomized trial with institutionalized Romanian children (68 placed in foster care, 68 left in usual care), inhibited, withdrawn signs of RAD dropped quickly after foster placement and by 30 months were indistinguishable from never-institutionalized children (Smyke et al., 2012).
- **Treatment aims:** the practice parameter calls for direct observation of the child with caregivers and has two aims: help the child form an attachment relationship when none exists, and improve any disturbed relating to non-caregivers (Zeanah et al., 2016).
- **Building caregiver sensitivity:** across 70 studies, interventions improved parental sensitivity (d = 0.33) and infant attachment security (d = 0.20); the most effective were brief, behaviorally focused programs, and gains in sensitivity carried over to attachment (Bakermans-Kranenburg et al., 2003). In a randomized trial with 120 maltreated toddlers, Attachment and Biobehavioral Catch-up (ABC) reduced disorganized attachment (32% vs. 57%) and increased secure attachment (52% vs. 33%) (Bernard et al., 2012). These studies did not diagnose RAD itself.
- **Avoid coercive therapies:** a national task force warned that the benefits and risks of many "attachment therapies" are unproven and that some techniques used with foster and adopted children are potentially harmful (Chaffin et al., 2006). Physical restraint, "holding therapy," and forced eye contact or rebirthing-type methods have no place in care.
- **Medication (referral notes):** no medication treats RAD. Medication may help co-occurring conditions such as ADHD.

Cognitive-behavioral work with RAD: how it is used (Bakermans-Kranenburg et al., 2003; Bernard et al., 2012):

- **Coach the caregiver** to follow the child's lead, respond warmly and predictably to distress, and offer comfort even when the child does not seem to want it.
- **Identify and challenge caregiver beliefs:** "This child rejects me," "If we love enough, he will bond right away," "I am failing as a parent," "She is doing this on purpose."
- **Support the caregiver's own regulation and respite,** since steady, calm care over months is the treatment.

5. Religion, spirituality, and Christian integration: what the evidence shows

Disorder-specific research. We found **no study of religion, faith communities, or faith-integrated treatment in reactive attachment disorder** (literature search, November 1, 2026).

Related evidence. Across 49 studies, positive religious coping went with better adjustment to stress and negative religious coping (feeling punished or abandoned by God) with worse (Ano & Vasconcelles, 2005), which bears on how exhausted

foster and adoptive parents cope. Across 97 outcome studies, therapy adapted to clients' religious beliefs worked about as well as standard therapy for distress and added spiritual benefits (Captari et al., 2018).

Honest summary for teaching:

1. A stable, sensitive caregiver and caregiver-focused interventions are the evidence base. Faith integration supports them; it does not replace them.
2. There is no research on faith and RAD. Faith communities can offer practical support and respite to foster and adoptive families.
3. Unproven and coercive "attachment therapies" have sometimes been offered to foster and adoptive families (Chaffin et al., 2006). Guide families toward evidence-based help.

6. Practical Christian-integration notes

These apply Dr. Campbell's identify, challenge, replace-with-Scripture method mainly with caregivers. Clinical items rest on the research above. Pastoral items are theological application, not research findings.

- **Identify (clinical).** Listen for caregiver beliefs that drive harsh or discouraged parenting, including spiritual forms: "God called us to adopt, so it should not be this hard," "A strong-willed child must be broken," "I must be failing God." With older children, listen for "Adults can't be counted on" (Chaffin et al., 2006).
- **Challenge (clinical).** Set realistic expectations: inhibited signs can improve quickly in a stable home, but trust takes time. Track small gains in seeking comfort (Smyke et al., 2012).
- **Replace (pastoral).** Offer Scripture of God's faithful, adopting love, for parent and child:
  - "God settles the lonely in families…" (Psalm 68:6)
  - "For you did not receive a spirit of slavery that returns you to fear, but you received the Spirit of sonship, by whom we cry, 'Abba! Father!'" (Romans 8:15). Adoption is at the heart of the gospel.
  - "Can a woman forget her nursing child, or lack compassion for the son of her womb? Even if she could forget, I will not forget you!" (Isaiah 49:15)
  - "Let us not grow weary in well-doing, for in due time we will reap a harvest if we do not give up." (Galatians 6:9). Encouragement for the long, patient work of parenting.
- **Faith community (clinical).** Offer meals, trained respite care, and parent support groups. Ask families what helps rather than offering unsolicited parenting advice.
- **Ministers:** take caregiver exhaustion seriously, steer families away from coercive "attachment therapies," and refer to clinicians experienced with foster and adopted children.

7. Quick classroom takeaways

- RAD means a withdrawn child who rarely seeks or accepts comfort, with limited positive emotion, after severe neglect or disrupted care; onset before age 5. Rule out autism.
- Treatment is a stable, sensitive caregiver plus caregiver-coaching interventions; withdrawn signs often improve quickly in a good home. Avoid coercive therapies.
- Replace "this child rejects me" with patient, adopting love: "God settles the lonely in families."
- Diagnose through caregiver interview (such as the DAI) and direct observation; brief screens (RPQ, SDQ) flag concern but do not diagnose.

Sources note

All 11 references were checked against Crossref, OpenAlex, and PubMed/Europe PMC records (Nov 1, 2026), and the figures quoted were checked against each study's abstract. Each DOI is a live link to the publisher record. Assessment instrument links and purchase and qualification levels were checked on the publishers' and distributors' websites, or on Crossref for original articles (Nov 1, 2026).

References

American Psychiatric Association. (2022). *Diagnostic and statistical manual of mental disorders* (5th ed., text rev.). https://doi.org/10.1176/appi.books.9780890425787

Ano, G. G., & Vasconcelles, E. B. (2005). Religious coping and psychological adjustment to stress: A meta-analysis. *Journal of Clinical Psychology*, *61*(4), 461–480. https://doi.org/10.1002/jclp.20049

Bakermans-Kranenburg, M. J., van IJzendoorn, M. H., & Juffer, F. (2003). Less is more: Meta-analyses of sensitivity and attachment interventions in early childhood. *Psychological Bulletin*, *129*(2), 195–215. https://doi.org/10.1037/0033-2909.129.2.195

Bernard, K., Dozier, M., Bick, J., Lewis-Morrarty, E., Lindhiem, O., & Carlson, E. (2012). Enhancing attachment organization among maltreated children: Results of a randomized clinical trial. *Child Development*, *83*(2), 623–636. https://doi.org/10.1111/j.1467-8624.2011.01712.x

Captari, L. E., Hook, J. N., Hoyt, W. T., Davis, D. E., McElroy-Heltzel, S. E., & Worthington, E. L., Jr. (2018). Integrating clients' religion and spirituality within psychotherapy: A comprehensive meta-analysis. *Journal of Clinical Psychology*, *74*(11), 1938–1951. https://doi.org/10.1002/jclp.22681

Chaffin, M., Hanson, R., Saunders, B. E., Nichols, T., Barnett, D., Zeanah, C., Berliner, L., Egeland, B., Newman, E., Lyon, T., Letourneau, E., & Miller-Perrin, C. (2006). Report of the APSAC task force on attachment therapy, reactive attachment disorder, and attachment problems. *Child Maltreatment*, *11*(1), 76–89. https://doi.org/10.1177/1077559505283699

Minnis, H., Macmillan, S., Pritchett, R., Young, D., Wallace, B., Butcher, J., Sim, F., Baynham, K., Davidson, C., & Gillberg, C. (2013). Prevalence of reactive attachment disorder in a deprived population. *The British Journal of Psychiatry*, *202*(5), 342–346. https://doi.org/10.1192/bjp.bp.112.114074

Pritchett, R., Pritchett, J., Marshall, E., Davidson, C., & Minnis, H. (2013). Reactive attachment disorder in the general population: A hidden ESSENCE disorder. *The Scientific World Journal*, *2013*, 818157. https://doi.org/10.1155/2013/818157

Smyke, A. T., Zeanah, C. H., Gleason, M. M., Drury, S. S., Fox, N. A., Nelson, C. A., & Guthrie, D. (2012). A randomized controlled trial comparing foster care and institutional care for children with signs of reactive attachment disorder. *American Journal of Psychiatry*, *169*(5), 508–514. https://doi.org/10.1176/appi.ajp.2011.11050748

Zeanah, C. H., & Gleason, M. M. (2015). Annual research review: Attachment disorders in early childhood—clinical presentation, causes, correlates, and treatment. *Journal of Child Psychology and Psychiatry*, *56*(3), 207–222. https://doi.org/10.1111/jcpp.12347

Zeanah, C. H., Chesher, T., Boris, N. W., & the American Academy of Child and Adolescent Psychiatry Committee on Quality Issues. (2016). Practice parameter for the assessment and treatment of children and adolescents with reactive attachment disorder and disinhibited social engagement disorder. *Journal of the American Academy of Child and Adolescent Psychiatry*, *55*(11), 990–1003. https://doi.org/10.1016/j.jaac.2016.08.004

Prepared for Clinical Companion (Brian Campbell). Teaching overview only. Clinically review before use. Not a substitute for the DSM-5-TR or clinical judgment. All 11 references and DOIs verified via Crossref, OpenAlex, and PubMed/Europe PMC records (Sunday, Nov 1, 2026). Effect sizes as reported in the cited sources.