Schizoid Personality Disorder (DSM-5-TR): Teaching Overview

Clinical Companion Knowledge Library: for counselors and counseling students

1. What it is

Schizoid personality disorder (SzPD) is a long-standing pattern of **detachment from social relationships** and a **restricted range of emotional expression** with others, beginning by early adulthood (American Psychiatric Association [APA], 2022). It belongs to **Cluster A** (odd or eccentric). Unlike avoidant personality disorder, the person is generally not longing for closeness while fearing rejection; closeness simply holds little appeal.

How common: in a large U.S. national survey, 3.13% of adults met criteria for SzPD (Grant et al., 2004). Across Western community studies, estimates vary widely depending on how personality disorders are assessed (Volkert et al., 2018).

A debated diagnosis. A review of the research found comparatively little evidence for SzPD's reliability and validity as a separate disorder. The authors suggested that "schizoid" patients may fall into an emotionally constricted group closer to schizotypal personality disorder and a withdrawn group closer to avoidant personality disorder, and they recommended rating schizoid traits on a continuum instead (Triebwasser et al., 2012). Separating schizoid from avoidant patterns remains a practical challenge (Lampe & Malhi, 2018).

Counselor framing: solitude is not a disorder by itself. Many quiet, private people live well. The diagnosis applies only when the pattern is pervasive, inflexible, and causes real distress or impairment.

2. Core symptoms to watch for

DSM-5-TR criteria, paraphrased (APA, 2022); use the manual itself for formal diagnosis. A pervasive pattern, beginning by early adulthood, with **4 or more** of:

- Neither desires nor enjoys **close relationships**, including being part of a family.
- Almost always **chooses solitary activities**.
- Has little, if any, **interest in sexual experiences** with another person.
- Takes **pleasure in few, if any, activities**.
- Lacks **close friends or confidants** other than first-degree relatives.
- Appears **indifferent to praise or criticism**.
- Shows **emotional coldness, detachment, or flattened affect**.

Rule out: the pattern must not occur only during schizophrenia, a mood disorder with psychotic features, another psychotic disorder, or autism spectrum disorder, and must not be due to a medical condition. Also distinguish depression (loss of interest that is new rather than lifelong) and avoidant personality disorder (desire for closeness blocked by fear) (APA, 2022; Lampe & Malhi, 2018).

When counselors must refer (general clinical consensus): refer for evaluation if there are signs of psychosis, significant depression, or substance use, or if withdrawal is new or worsening, which may point to another disorder.

3. What works: treatment and the evidence

Bottom line for class: there are no controlled treatment trials for SzPD. Help usually targets the problem the person seeks help for (often depression, work stress, or family pressure) while respecting the client's preferred distance.

Evidence overview (honest):

- **No SzPD trials:** we found no randomized trial of any psychotherapy or medication for SzPD (literature search, November 1, 2026). The diagnosis itself is debated, which has slowed research (Triebwasser et al., 2012).
- **Borrowed approaches:** clinicians commonly use supportive and cognitive-behavioral methods for co-occurring depression and anxiety, and social-skills practice when the client wants it. These are clinical practice, not SzPD-specific evidence.

- **Medication (referral notes):** no medication treats SzPD itself. Co-occurring depression or anxiety may be treated in the usual ways.

Cognitive work with SzPD: how it is used (clinical practice):

- **Respect the pace.** Keep sessions practical and low-pressure; a steady, undemanding presence can itself be a new experience of safe connection.
- **Identify and examine core beliefs:** "Relationships aren't worth the trouble," "I'm better off alone," "Feelings are pointless," "Nothing is much fun anyway."
- **Experiment gently:** track pleasure and meaning in small activities, and test whether brief, structured contact with one person adds anything worthwhile.

4. Religion, spirituality, and Christian integration: what the evidence shows

Disorder-specific research. We found **no study of religion or spirituality in schizoid personality disorder** and no faith-integrated treatment trial (literature search, November 1, 2026).

Related evidence. Across 49 studies, positive religious coping went with better adjustment to stress and negative religious coping with worse (Ano & Vasconcelles, 2005). Across 97 outcome studies, therapy adapted to clients' religious beliefs worked about as well as standard therapy for distress and produced added spiritual benefits (Captari et al., 2018).

Honest summary for teaching:

1. No treatment has been tested for SzPD. Treat what brings the client in, and respect the client's comfort with distance.
2. There is no research on faith and SzPD. Some schizoid individuals value private, contemplative faith more than group activities; this need not be pathologized.
3. Pressing for church involvement or emotional expression can drive these clients away. Offer connection, do not demand it.

5. Practical Christian-integration notes

These apply Dr. Campbell's identify, challenge, replace-with-Scripture method in a gentle, client-paced way. Clinical items rest on the research above and on general clinical practice. Pastoral items are theological application, not research findings.

- **Identify (clinical).** Listen for beliefs that close off connection: "I don't need anyone," "Church is just people making noise," "God is distant, and that suits me," "Nothing matters much."
- **Challenge (clinical).** Rather than arguing, explore small costs of total detachment the client already notices (loneliness at times, family strain, missed opportunities) and test one small, structured connection (Ano & Vasconcelles, 2005).
- **Replace (pastoral).** Offer Scripture that shows God's design for connection, gently and without pressure:
  - "The LORD God also said, 'It is not good for the man to be alone. I will make for him a suitable helper.'" (Genesis 2:18)
  - "Two are better than one, because they have a good return for their labor. For if one falls down, his companion can lift him up; but pity the one who falls without another to help him up!" (Ecclesiastes 4:9–10)
  - "God settles the lonely in families…" (Psalm 68:6). God's initiative, not the person's performance.
  - "The LORD your God is among you; He is mighty to save. He will rejoice over you with gladness; He will quiet you with His love; He will rejoice over you with singing." (Zephaniah 3:17). A God who draws near.
- **Faith community (clinical).** Offer quiet, task-based ways to belong (setting up, tech support, prayer lists, one-on-one Bible study) instead of high-emotion small groups.
- **Ministers:** accept reserved members without labeling them cold or unspiritual, and refer if withdrawal is new, deepening, or paired with odd beliefs or depression.

6. Quick classroom takeaways

- SzPD is lifelong detachment from relationships and restricted emotion (4 of 7 criteria). It is distinct from avoidant PD (fear-driven) and depression (new loss of interest); its validity as a category is debated.
- No treatment trials exist; treat co-occurring problems and respect the client's pace.
- Gently replace "I don't need anyone" with the truth that "It is not good for the man to be alone," offered as invitation, not pressure.

Sources note

All 7 references were checked against Crossref, OpenAlex, and PubMed/Europe PMC records (Nov 1, 2026), and the figures quoted were checked against each study's abstract. Each DOI is a live link to the publisher record.

References

American Psychiatric Association. (2022). *Diagnostic and statistical manual of mental disorders (5th ed., text rev.)*. https://doi.org/10.1176/appi.books.9780890425787

Ano, G. G., & Vasconcelles, E. B. (2005). Religious coping and psychological adjustment to stress: A meta-analysis. *Journal of Clinical Psychology*, *61*(4), 461–480. https://doi.org/10.1002/jclp.20049

Captari, L. E., Hook, J. N., Hoyt, W. T., Davis, D. E., McElroy-Heltzel, S. E., & Worthington, E. L. (2018). Integrating clients' religion and spirituality within psychotherapy: A comprehensive meta-analysis. *Journal of Clinical Psychology*, *74*(11), 1938–1951. https://doi.org/10.1002/jclp.22681

Grant, B. F., Hasin, D. S., Stinson, F. S., Dawson, D. A., Chou, S. P., Ruan, W. J., & Pickering, R. P. (2004). Prevalence, correlates, and disability of personality disorders in the United States: Results from the National Epidemiologic Survey on Alcohol and Related Conditions. *The Journal of Clinical Psychiatry*, *65*(7), 948–958. https://doi.org/10.4088/JCP.v65n0711

Lampe, L., & Malhi, G. S. (2018). Avoidant personality disorder: Current insights. *Psychology Research and Behavior Management*, *11*, 55–66. https://doi.org/10.2147/PRBM.S121073

Triebwasser, J., Chemerinski, E., Roussos, P., & Siever, L. J. (2012). Schizoid personality disorder. *Journal of Personality Disorders*, *26*(6), 919–926. https://doi.org/10.1521/pedi.2012.26.6.919

Volkert, J., Gablonski, T.-C., & Rabung, S. (2018). Prevalence of personality disorders in the general adult population in Western countries: Systematic review and meta-analysis. *The British Journal of Psychiatry*, *213*(6), 709–715. https://doi.org/10.1192/bjp.2018.202

Prepared for Clinical Companion (Brian Campbell). Teaching overview only. Clinically review before use. Not a substitute for the DSM-5-TR or clinical judgment. All 7 references and DOIs verified via Crossref, OpenAlex, and PubMed/Europe PMC records (Sunday, Nov 1, 2026). Effect sizes as reported in the cited sources.