Schizotypal Personality Disorder (DSM-5-TR): Teaching Overview

Clinical Companion Knowledge Library: for counselors and counseling students

Brian Campbell, Ph.D.

Prepared with AI research and drafting assistance; content directed, reviewed, and approved by the author. © 2026 Brian Campbell, Ph.D. All rights reserved.

1. What it is

Schizotypal personality disorder (STPD) is a long-standing pattern of **social and interpersonal deficits** (acute discomfort with close relationships) combined with **odd thinking, perceptions, and behavior**, beginning by early adulthood (American Psychiatric Association [APA], 2022). It belongs to **Cluster A** and is also listed in DSM-5-TR's schizophrenia spectrum chapter, because it shares many genetic, biological, and thinking features with schizophrenia (Koenigsberg et al., 2003; Rosell et al., 2014).

How common: in a U.S. national survey, 3.9% of adults met lifetime criteria (4.2% of men, 3.7% of women). STPD was linked to substantial mental disability in both sexes and frequently co-occurred with mood, anxiety, and other personality disorders (Pulay et al., 2009).

Counselor framing: these clients are often lonely, anxious, and misunderstood. Their unusual ideas are usually held with some doubt, not with the fixed certainty of a delusion, and their social anxiety tends not to ease with familiarity.

2. Core symptoms to watch for

DSM-5-TR criteria, paraphrased (APA, 2022); use the manual itself for formal diagnosis. A pervasive pattern, beginning by early adulthood, with **5 or more** of:

- **Ideas of reference** (not delusions): feeling that unrelated events carry special personal meaning.
- **Odd beliefs or magical thinking** out of keeping with the person's culture (such as telepathy, a "sixth sense," or bizarre fantasies).
- **Unusual perceptual experiences,** including bodily illusions.
- **Odd thinking and speech** (vague, overelaborate, or stereotyped).
- **Suspiciousness** or paranoid ideas.
- **Inappropriate or constricted emotion.**
- **Odd, eccentric, or peculiar behavior or appearance.**
- **Lack of close friends** or confidants other than first-degree relatives.
- **Excessive social anxiety** that does not lessen with familiarity and tends to come from paranoid fears rather than negative self-judgment.

Cultural caution: religious experiences, beliefs about spiritual realities, and practices shared within a person's faith community are **not** symptoms. DSM-5-TR asks clinicians to judge beliefs against the person's own cultural and religious context (APA, 2022).

When counselors must refer (general clinical consensus): refer promptly for psychiatric evaluation if beliefs become fixed, if hearing voices or seeing things becomes more frequent or convincing, if functioning drops sharply, or if there is suicidal thinking. These can signal emerging psychosis.

3. Frequently used assessment instruments

These are among the most widely used measures for screening, rating severity, and tracking progress in schizotypal personality disorder. **Screening tools do not diagnose;** a positive screen calls for a clinical interview against DSM-5-TR criteria. Click an instrument's website to find it.

Instrument	Best used for	Format	Purchase required?	Training or qualification	Where to find it
SPQ (Schizotypal Personality Questionnaire)	Schizotypal traits (odd beliefs, unusual experiences, social anxiety).	Self-report, 74 yes/no items	No (free)	None beyond general clinical training	Original article
PQ-B (Prodromal Questionnaire-Brief)	Screening for early psychosis risk.	Self-report, 21 items	No (free)	None beyond general clinical training	Original article
APA Clinician-Rated Dimensions of Psychosis Symptom Severity	Rating psychotic-like symptoms and watching for change toward psychosis.	Clinician-rated, 8 items	No (free from the APA for clinical use)	None beyond general clinical training	APA assessment measures
SCID-5-PD (Structured Clinical Interview for DSM-5 Personality Disorders)	Diagnosing all ten DSM-5 personality disorders, with an optional self-report screener.	Semistructured clinician interview	Yes	★ Clinical interviewing skill; training recommended	APA Publishing
PID-5 (Personality Inventory for DSM-5)	Dimensional personality trait profile (negative affect, detachment, antagonism, disinhibition, psychoticism); brief and informant forms.	Self-report; 220-item full or 25-item brief form	No (free from the APA for clinical use)	None beyond general clinical training	APA assessment measures
MCMI-IV (Millon Clinical Multiaxial Inventory-IV)	Adults in treatment; personality patterns alongside clinical syndromes.	Self-report, 195 true/false items, about 25 to 30 minutes	Yes	★ Level C	Pearson
MMPI-3 (Minnesota Multiphasic Personality Inventory-3)	Broad personality and psychopathology assessment, with validity scales for over- and under-reporting.	Self-report, 335 true/false items, about 25 to 50 minutes	Yes	★ Level C	Pearson

★ **= requires professional qualification to purchase or specialized training.** Publishers sell these tests only to qualified users. At Pearson, **Level B** generally means a master's degree in a related field with formal assessment training, or a health care license; **Level C** means a doctorate in psychology or a related field with formal assessment training, or state licensure in a related field (Pearson qualifications policy). MHS, PAR, and WPS use similar levels. Check each publisher's page before ordering. "Original article" links go to the published study that introduced a free scale that has no official website; the scale can usually be obtained there or from its authors.

4. What works: treatment and the evidence

Bottom line for class: STPD research is small. Low-dose antipsychotic medication has modest trial support; psychological help focuses on social anxiety, odd thinking, and skills, borrowing from cognitive-behavioral work for people at risk of psychosis.

Evidence overview (honest effect sizes):

- **Medication (referral notes):** in a 9-week randomized, placebo-controlled trial of 25 patients, low-dose risperidone (0.25 to 2 mg per day) reduced negative and general symptoms by week 3 and positive symptoms by week 7, and was well tolerated (Koenigsberg et al., 2003). This is one small trial.
- **CBT for people at risk of psychosis:** in randomized trials (six trials and 800 participants at 6 months), CBT roughly halved the rate of transition to psychosis, with risk ratios of 0.41 to 0.47 from 6 months through 18 to 24 months (Hutton & Taylor, 2014). These trials studied at-risk mental states rather than STPD specifically, but the overlap is substantial.
- **Thinking skills:** STPD involves problems with attention and working memory similar to those in schizophrenia (Rosell et al., 2014). Practical, concrete, structured therapy tends to work better than open-ended talk.

Cognitive-behavioral work with STPD: how it is used (Hutton & Taylor, 2014):

- **Normalize and collaborate.** Treat unusual experiences as something to examine together, not something to be argued away or feared.
- **Identify and test beliefs:** "Other people can read my thoughts," "That song was a message for me," "People are laughing at me," "I'm too strange to fit in." Look for other explanations and gather evidence.
- **Reduce social anxiety step by step** and practice concrete social skills in low-pressure settings.

© 2026 Brian Campbell, Ph.D. All rights reserved.

5. Religion, spirituality, and Christian integration: what the evidence shows

Faith is not the same as schizotypy. In a study of 189 adults, the strength of religiosity was **not** linked to unusual experiences or disorganized thinking, and was linked to **lower** impulsive nonconformity. Among members of religious communities, religiosity related negatively to schizotypy (Hanel et al., 2019).

How faith is used matters. Across 49 studies, positive religious coping went with better adjustment to stress and negative religious coping (feeling punished or abandoned by God) with worse (Ano & Vasconcelles, 2005). Across 97 outcome studies, therapy adapted to clients' religious beliefs worked about as well as standard therapy for distress and produced added spiritual benefits (Captari et al., 2018).

Intervention studies. We found **no trial of Christian or religiously integrated treatment for STPD** (literature search, November 1, 2026).

Honest summary for teaching:

1. Low-dose medication and structured, CBT-style help for unusual thinking and social anxiety are the best available evidence. Faith integration supports them; it does not replace them.
2. Ordinary faith is not a symptom. Unusual spiritual ideas that are idiosyncratic, distressing, or not shared by the person's faith community deserve careful, respectful evaluation.
3. No faith-integrated trial exists. Avoid both dismissing the client's spiritual life and reinforcing odd beliefs.

6. Practical Christian-integration notes

These apply Dr. Campbell's identify, challenge, replace-with-Scripture method within structured, CBT-informed care. Clinical items rest on the research above. Pastoral items are theological application, not research findings.

- **Identify (clinical).** Listen for idiosyncratic spiritual ideas held apart from the person's church's teaching: "God sends me secret signs in license plates," "I can sense evil in people by their colors," "The pastor's sermon was really about me" (Hanel et al., 2019). To identify deeper false beliefs, see: Core Beliefs, Parental Messages, or use the Semantic Search Tool.
- **Challenge (clinical).** Explore gently: Does your church or Scripture teach this? What else might explain it? What happens when you test it? Keep the tone curious, never mocking (Hutton & Taylor, 2014). **Possible cognitive errors** to watch for, with their definitions from the Belief Counseling Cognitive Errors Checklist and the symptoms they tend to feed:
  - **Jumping to conclusions** — deciding without the facts. *Feeds:* ideas of reference. *Sounds like:* "The pastor's sermon was really about me."
  - **Confirmatory bias** — looking for proof I'm right, ignoring the rest. *Feeds:* magical thinking. *Sounds like:* "Another license plate sign—more proof."
  - **Selective attention** — noticing only what fits my belief. *Feeds:* odd beliefs and perceptions. *Sounds like:* "I only notice the signs that are meant for me."
  - **Emotional reasoning** — I feel it, so it must be true. *Feeds:* unusual perceptions. *Sounds like:* "I sense evil in him, so it must be there."
  - **Mind reading** — assuming I know what others think. *Feeds:* suspicion and social anxiety. *Sounds like:* "They think I'm strange."
  - **Fortune telling** — predicting a bad outcome. *Feeds:* avoiding people. *Sounds like:* "If I go, they'll laugh at me."
- **Questions to dispute the belief.** Keep the tone curious, never mocking. Have the client choose questions from the Questioning False Beliefs form that "speak to" the belief. To see how, watch Dr. Campbell's video Integrated Christian Counseling: Questioning Your Beliefs. Selected questions that fit the errors above:
  - "Are there other ways of looking at this situation?" *For:* jumping to conclusions.
  - "What are some of the best arguments against my belief or conclusion?" and "How does the Bible speak to this?" *For:* confirmatory bias and private spiritual ideas.
  - "Is it possible that I am distorting the facts?" *For:* selective attention.
  - "What's the actual evidence for my belief?" *For:* emotional reasoning and mind reading.
  - "If I were a betting person, would I bet a million dollars that the thing I dread will happen?" *For:* fortune telling.
- **Replace (pastoral).** Offer Scripture that grounds faith in what is clear, shared, and peace-giving:
  - "Beloved, do not believe every spirit, but test the spirits to see whether they are from God…" (1 John 4:1). Testing experiences is itself biblical.

© 2026 Brian Campbell, Ph.D. All rights reserved.

    - "For God has not given us a spirit of fear, but of power, love, and self-control." (2 Timothy 1:7)
    - "Finally, brothers, whatever is true, whatever is honorable, whatever is right, whatever is pure, whatever is lovely, whatever is admirable—if anything is excellent or praiseworthy—think on these things." (Philippians 4:8). Anchoring thought in what is true.
    - "I praise You, for I am fearfully and wonderfully made…" (Psalm 139:14). For the client who feels too strange to belong.
- **Faith community (clinical).** Offer patient, accepting relationships and simple roles. Members can welcome an eccentric person warmly without affirming unusual ideas.
- **Ministers:** avoid interpreting unusual experiences as demonic or as special revelation, and refer promptly if beliefs become fixed or functioning declines.

7. Quick classroom takeaways

- STPD combines social discomfort with odd beliefs, perceptions, and behavior (5 of 9 criteria). About 4% of U.S. adults; it is part of the schizophrenia spectrum. Watch for signs of psychosis.
- Low-dose risperidone has small-trial support; CBT methods for at-risk states and social-skills work are the main psychological tools.
- Religiosity is not schizotypy. "Test the spirits" gently, and replace fear and odd ideas with clear, shared truth.
- Screen traits with the free SPQ and watch for psychosis risk with the PQ-B.

Sources note

All 8 references were checked against Crossref, OpenAlex, and PubMed/Europe PMC records (Nov 1, 2026), and the figures quoted were checked against each study's abstract. Each DOI is a live link to the publisher record. Assessment instrument links and purchase and qualification levels were checked on the publishers' and distributors' websites, or on Crossref for original articles (Nov 1, 2026).

References

American Psychiatric Association. (2022). *Diagnostic and statistical manual of mental disorders (5th ed., text rev.)*. https://doi.org/10.1176/appi.books.9780890425787

Ano, G. G., & Vasconcelles, E. B. (2005). Religious coping and psychological adjustment to stress: A meta-analysis. *Journal of Clinical Psychology*, *61*(4), 461–480. https://doi.org/10.1002/jclp.20049

Captari, L. E., Hook, J. N., Hoyt, W. T., Davis, D. E., McElroy-Heltzel, S. E., & Worthington, E. L. (2018). Integrating clients' religion and spirituality within psychotherapy: A comprehensive meta-analysis. *Journal of Clinical Psychology*, *74*(11), 1938–1951. https://doi.org/10.1002/jclp.22681

Hanel, P. H. P., Demmrich, S., & Wolfradt, U. (2019). Centrality of religiosity, schizotypy, and human values: The impact of religious affiliation. *Religions*, *10*(5), 297. https://doi.org/10.3390/rel10050297

Hutton, P., & Taylor, P. J. (2014). Cognitive behavioural therapy for psychosis prevention: A systematic review and meta-analysis. *Psychological Medicine*, *44*(3), 449–468. https://doi.org/10.1017/S0033291713000354

Koenigsberg, H. W., Reynolds, D., Goodman, M., New, A. S., Mitropoulou, V., Trestman, R. L., Silverman, J., & Siever, L. J. (2003). Risperidone in the treatment of schizotypal personality disorder. *The Journal of Clinical Psychiatry*, *64*(6), 628–634. https://doi.org/10.4088/JCP.v64n0602

Pulay, A. J., Stinson, F. S., Dawson, D. A., Goldstein, R. B., Chou, S. P., Huang, B., Saha, T. D., Smith, S. M., Pickering, R. P., Ruan, W. J., Hasin, D. S., & Grant, B. F. (2009). Prevalence, correlates, disability, and comorbidity of DSM-IV schizotypal personality disorder: Results from the Wave 2 National Epidemiologic Survey on Alcohol and Related Conditions. *The Primary Care Companion to The Journal of Clinical Psychiatry*, *11*(2), 53–67. https://doi.org/10.4088/PCC.08m00679

Rosell, D. R., Futterman, S. E., McMaster, A., & Siever, L. J. (2014). Schizotypal personality disorder: A current review. *Current Psychiatry Reports*, *16*(7), 452. https://doi.org/10.1007/s11920-014-0452-1

Prepared for Clinical Companion (Brian Campbell). Teaching overview only. Clinically review before use. Not a substitute for the DSM-5-TR or clinical judgment. All 8 references and DOIs verified via Crossref, OpenAlex, and PubMed/Europe PMC records (Sunday, Nov 1, 2026). Effect sizes as reported in the cited sources.

© 2026 Brian Campbell, Ph.D. All rights reserved.