

1 New Mathematics and Natural Computation
 2 (2027) 1–24
 3 © World Scientific Publishing Company
 4 DOI: 10.1142/S1793005727500074



5 Threshold Dynamics of Stochastic SIS Epidemic Models 6 with Logistic Recruitment Rate

7 Soulaïmane Aznague*, Adel Settati[†], Aadil Lahrouz[‡],
 8 Bilal Harchaoui[§], Khalid El Bakkoui[¶] and Abdeladim Nait Brahimi^{||}

9 *Laboratory of Mathematics and Applications*
 10 *Faculty of Science and Technology of Tangier (FSTT)*
 11 *Abdelmalek Essaadi University*
 12 *Tetouan, Morocco*

13 *soulaimane.aznague@etu.uae.ac.ma

14 [†]settati.adel@yahoo.fr

15 [‡]lahrouzadil@gmail.com

16 [§]Bilal.harchaoui@outlook.com

17 [¶]khalid.elbakkoui@etu.uae.ac.ma

18 ^{||}abdeladim.naitbrahim@etu.uae.ac.ma



AQ: Please approve the
 edits in the affiliation.

AQ: Please provide valid
 ORCID ID of all authors. Note
 that the http address should be
 in valid format, leading to the
 correct webpage.

19 Received 30 March 2024

20 Accepted 13 April 2025

21 Published



22 This study explores an SIS epidemic model that integrates logistic growth and a saturated
 23 incidence rate among susceptible individuals. The primary objective is to examine the
 24 long-term behavior of the stochastic system. For this purpose, a key parameter, λ , is
 25 introduced to differentiate the system dynamics into two cases. In scenarios where $\lambda < 0$,
 26 the model predicts the eventual extinction of the disease. In contrast, for $\lambda > 0$, we
 27 show an ergodic stationary distribution. Numerical simulations validate our theoretical
 28 conditions for extinction, persistence, and the existence of a stationary distribution.

29 *Keywords:* SIS epidemic model; logistic population growth; disease extinction; stability
 30 in distribution; ergodic properties.

31 ~~Mathematics Subject Classification 2020:~~

32 1. Introduction

During the early 20th century, epidemiology underwent a significant transformation,
 driven by the groundbreaking contributions of eminent scientists such as Anderson
 Gray McKendrick and Janet Leigh. Their pioneering work introduced the concept
 of mathematical modeling, which has since evolved into an indispensable tool in the
 field. This mathematical modeling has profoundly impacted the management of out-
 breaks and epidemics, playing a pivotal role in guiding evidence-based public health

*Corresponding author.

2 *S. Aznague et al.*

interventions. Epidemiology has evolved significantly, owing to notable physicians such as Quinto Tiberio Angelerio, who demonstrated remarkable proficiency in managing the plague outbreak in Alghero, Sardinia, in 1582. However, the actual emergence of modern epidemiology as a formal scientific discipline occurred during the 19th century. Often referred to as the “father of modern epidemiology,” John Snow made a meaningful breakthrough when he meticulously traced a devastating cholera outbreak in London to water contamination from the Broad Street pump. This groundbreaking investigation is the pivotal moment that laid the cornerstone for contemporary epidemiology, shaping it into the scientific field we recognize today. Epidemiology is a scientific discipline investigating epidemics, diseases, and various health-related conditions, including those unrelated to diseases. Its roots can be traced to ancient Greece, notably through the influential work of Hippocrates of Kos, a renowned physician who made notable contributions by distinguishing between epidemic and endemic diseases. Epidemiology, in its broader scope, also encompasses the study of diseases affecting plants, domestic animals, and livestock. An epidemic is characterized by a significant and abnormal surge in the occurrence of a specific disease within a population, typically manifesting rapidly. The intricate disease transmission process is influenced by many factors, encompassing both the characteristics of the infectious agent and the complex dynamics within the host population. Regarding the infectious agent, its inherent characteristics, such as its mode of transmission (e.g. respiratory droplets, direct contact), the duration of infectivity, and its responsiveness to medical interventions like treatments and vaccines, are crucial factors that determine its ability to spread among individuals. Equally important are the host population elements that influence the dynamics of an epidemic. Factors such as social interactions, demographics (e.g. age, gender), cultural practices, geographic distribution, and economic conditions are pivotal in determining a population’s susceptibility and resilience in the face of the disease. Across the pages of recorded history, human civilization has wrestled with recurrent epidemics and pandemics. These outbreaks of infectious diseases have inflicted significant human suffering, societal upheaval, and economic turbulence. Given the formidable nature of these challenges, the precise prediction of outbreak progression becomes paramount to effectively mitigating their adverse impacts. Epidemiologic modeling is at the heart of this pursuit, a fundamental tool for comprehending the intricacies of disease transmission dynamics and formulating informed strategies for containment and prevention. Using dynamical systems to model infectious diseases is crucial in examining both theoretical and quantitative facets of epidemic spread. Following the foundational contributions of McKendrick and Kermack,¹ various compartmental models have emerged, employing ordinary, fractional, and stochastic differential equations (SDEs). Every model presented in existing literature has its unique characteristics and limitations, and it is essential to recognize that no model is flawless, regardless of the number of variables or parameters included. Additionally, there tends to be an inverse relationship between model complexity and the depth of mathematical analysis. Commonly, researchers focus on either

AQ: please
approve the short
title



Threshold Dynamics of Stochastic SIS Epidemic Models 3

SIS (Susceptible–Infectious–Susceptible) or SIR (Susceptible–Infectious–Recovered) models, where S denotes susceptible individuals, I represents infectious persons, and R refers to those who have recovered. It is also possible to incorporate additional compartments, such as vaccinated individuals or those exposed to the disease. In this paper, our attention is specifically directed toward SIS epidemic models, which are effectively employed in the study of diseases such as sexually transmitted infections and bacterial diseases.^{2–15} These models aptly encapsulate the progression of diseases where lasting immunity is not conferred, resulting in individuals reverting to a susceptible state post-infection. While certain researchers^{16,17} have explored analytical or explicit series solutions for two-dimensional SIS differential systems, other mathematical methodologies such as bifurcation theory, stability theory, or the Lyapunov method are more commonly utilized. This preference arises because most two-dimensional differential equations, particularly non-linear ones, are seldom solvable via integration. Additionally, the pursuit of explicit solutions is often less prioritized by authors, as these alternative mathematical tools yield valuable insights into aspects such as the asymptotic behavior of the solutions, which can be more informative for understanding system dynamics. Epidemic models aim to address various queries, including determining the number of new infections that arise when infectious individuals interact with a susceptible group. The resolution to this query typically hinges on two key assumptions. The first assumption posits that each individual in a population is equally likely to come into contact with any other individual at any given moment, a concept referred to as homogeneous mixing. The second assumption is the mass-action principle, which suggests that a certain proportion of these contacts will result in infection at any given time.^{18,19} The concepts above can be mathematically articulated as follows: Let $S(t)$ and $I(t)$ represent the number of susceptible and infectious individuals at time t , respectively. Then, the term $\beta S(t)I(t)$ signifies the new portion of susceptible individuals who become infectious at time t . In this study, we retain the assumption that transmission occurs almost instantaneously. However, the second assumption is modifiable by adopting a different incidence rate function. Rather than the conventional $\beta S(t)I(t)$, alternative models suggest using $\beta g(S, I)$, where $g(S, I)$ is a function that is not bilinear, introducing a variation in the interaction dynamics between susceptible and infectious individuals. In our study, we have opted for a Holling-type functional response function, defined as $g(S, I) = SI/(1 + \alpha S)$. This formulation was initially introduced into epidemic modeling by Macdonald²⁰ and later adopted by May *et al.*²¹ Notably, when the number of susceptible individuals is considerably large, the incidence rate function $\beta g(S, I)$ tends to approximate $\beta I/\alpha$, with β representing the contact parameter between susceptible and infected individuals. This characteristic is why $g(S, I)$ is a saturated function concerning the susceptibles. Commonly in the literature, the population growth rate is assumed to be constant, leading to exponential demographic changes. However, this assumption usually holds only over short time frames. In practical scenarios, the notion of fluctuating total populations is more applicable, especially in the context of diseases with high mortality rates

4 *S. Aznague et al.*

or those that persist over a considerable duration.^{22–28} Additionally, population growth is naturally constrained by environmental factors, such as the availability of resources. A notable approach for encapsulating this concept is the logistic model, initially proposed by Verhulst in Ref. 29. This model illustrates exponential growth within a population, eventually tempered by the environmental carrying capacity, thus providing a more realistic representation of population dynamics under such constraints. Numerous researchers have explored and developed epidemic models incorporating logistic growth equations.^{14,30–33} Building upon this foundation, we present a deterministic SIS epidemic model that includes logistic growth, structured as follows:

$$\begin{cases} \frac{dS}{dt} = rS \left(1 - \frac{S}{K}\right) - \mu S - \frac{\beta SI}{1 + \alpha S} + \gamma I, \\ \frac{dI}{dt} = -(\mu + \gamma)I + \frac{\beta SI}{1 + \alpha S}. \end{cases} \quad (1.1)$$

At a given moment, $S(t)$ denotes the quantity of susceptible individuals, and $I(t)$ indicates the number of currently infected persons. The parameters in the model, all positive, have specific meanings: r is the intrinsic growth rate of the susceptible population, and K symbolizes the carrying capacity. Additionally, β signifies the rate of disease transmission, α represents the half-saturation constant, and γ refers to the recovery rate of infected individuals. The parameter μ reflects the natural mortality rate within the population. We operate under the assumption that $r > \mu$, as the scenario where $\mu \geq r$ lacks biological relevance. Concurrently, stochastic variations are crucial in industrial production and ecosystem dynamics. Consequently, many researchers have integrated the concept of random noise, often represented as stochastic white noise, into deterministic epidemic models. In certain studies, it is posited that every individual in the population is concurrently influenced by various random environmental factors.^{3,4,34} In a recent study,⁵ researchers introduced an SIS epidemic model that incorporates multiple independent Brownian motions. They identified a stochastic threshold crucial for determining whether the disease persists or becomes extinct within the host population. This threshold is the Lyapunov exponent of $I(t)$, particularly when $I(t)$ remains small over an extended period. Inspired by these findings, our proposed model considers stochastic perturbations as environmental noise impacting individuals directly. These perturbations are assumed to be proportional to the variables in the system, and the model can be expressed as follows:

$$\begin{cases} dS = \left[rS \left(1 - \frac{S}{K}\right) - \mu S - \frac{\beta SI}{1 + \alpha S} + \gamma I \right] dt + \sigma_1 S dB_1, \\ dI = \left[-(\mu + \gamma)I + \frac{\beta SI}{1 + \alpha S} \right] dt + \sigma_2 I dB_2. \end{cases} \quad (1.2)$$

- 1 In our model, we incorporate independent Brownian motions, represented by $B_i(t)$ for $i = 1, 2$, each associated with their respective intensities of white noise, denoted

as σ_i , and all are positive. The organization of this paper follows a systematic approach. Starting with Sec. 2, we present essential results necessary for the subsequent sections. Section 3 is devoted to deriving the required conditions for the extinction of the disease. In Sec. 4, we establish the criteria needed to ensure the system attains an ergodic stationary distribution. Following this, Sec. 5 comprises a series of numerical simulations to corroborate the accuracy of our analytical results. The paper concludes with a summary of our findings and a discussion of potential future research directions.

2. Preliminary Results

We define $(\Omega, \mathcal{T}, \{\mathcal{T}_t\}_{t \geq 0}, \mathbb{P})$ as a complete probability space, complete with a filtration $\{\mathcal{T}_t\}_{t \geq 0}$ that meets the usual conditions. The notations $\mathbb{E}_{(s,i)}$ and $\mathbb{P}_{(s,i)}$ are used to denote the expected value and probability, starting from $S(0) = s, I(0) = i$. In this paper, we adopt the following notations:

$$\mathbb{R}_+^{2,o} = \{(s, i) : s > 0, i > 0\}, \quad \text{for the strictly positive quadrant.}$$

$$\mathbb{R}_+^2 = \{(s, i) : s \geq 0, i \geq 0\}, \quad \text{for the non-negative quadrant.}$$

$$\mathbb{R}_+^{2,*} = \{(s, i) : s \geq 0, i > 0\}, \quad \text{for the semi-positive quadrant.}$$

Our research focuses on examining a d -dimensional Itô process, characterized by the subsequent SDE:

$$dX(t) = f(X)dt + g(X)dB(t) \quad \text{for each } t \geq t_0. \quad (2.1)$$

Here, $B(t)$ represents a d -dimensional white noise and the starting value $X(0) \in \mathbb{R}^d$. In virtue of the Itô formula, the stochastic equation verified by $V(X)$, with V being a function that is continuously twice differentiable and defined on $\mathbb{R}^d \times \mathbb{R}^+$ is

$$dV(X) = \mathcal{L}V(X)dt + \nabla V(X)g(X)dB(t),$$

where $\nabla V = (\frac{\partial V}{\partial x_1}, \dots, \frac{\partial V}{\partial x_d})$ is gradient of V and \mathcal{L} is the differential operator associated to system (2.1) defined by

$$\mathcal{L} = \frac{\partial}{\partial t} + \sum_{i=1}^d f_i(X) \frac{\partial}{\partial X_i} + \frac{1}{2} \sum_{i,j=1}^d [g^T(X) \cdot g(X)]_{ij} \frac{\partial^2}{\partial X_i \partial X_j}.$$

An appropriate beginning would be introducing a theorem that establishes the existence, uniqueness of solutions, and bounded nature.

Theorem 2.1. (i) For any given initial value $X(0) \in \mathbb{R}_+^2$, a unique and positive solution X to the SDE as stated in Eq. (1.2) can be assured for all $t \geq 0$, such

6 *S. Aznague et al.*

that

$$\mathbb{P}_{(s,i)} \{X(t) \in \mathbb{R}_+^2\} = 1.$$

Furthermore

$$\mathbb{P}_{(s,0)} [I(t) = 0; \text{ for any } t \geq 0] = 1,$$

then

$$\mathbb{P}_{(s,i)} \left[(S(t), I(t)) \in \mathbb{R}_+^{(2,o)}; \text{ for all } t > 0 \right] = 1, \quad i > 0.$$

- (ii) For each $\theta > 0$ and sufficiently small, it can be established that there are values θ_1, θ_2 , both exceeding zero, for which the following conditions hold:

$$\mathbb{E}_{(s,i)} [S(t) + I(t) + 1]^{\theta+1} \leq [s + i + 1]^{\theta+1} \exp(-\theta_2 t) + \frac{\theta_1}{\theta_2}. \quad (2.2)$$

Moreover, for all $M, T, \epsilon > 0$, there exists $H_{M,T,\epsilon} > 0$ such that

$$\mathbb{P}_{(s,i)} \left[\sup_{t \in [0,T]} \{S_t + I_t\} \leq H_{M,\epsilon,T} \right] \geq 1 - \epsilon \quad \text{for } (s, i) \in [0, M]^2. \quad (2.3)$$

Proof. The demonstration of solution positivity is based on methods akin to those found in Refs. 14 and 35, Theorem 2.1. However, to maintain conciseness, these details are not included here. Considering (1.2), it becomes evident that $I(t)$ conforms to the following criteria:

$$I(t) = i \exp \left[\int_0^t \frac{\beta S(u)}{1 + \alpha S(u)} du - (\mu + \gamma)t + \sigma_2 B_2 \right]. \quad (2.4)$$

Furthermore, deriving from Eq. (2.4), when setting $i = 0$, it leads to the following result:

$$\mathbb{P}_{(s,0)} [I(t) = 0; \text{ for each } t \geq 0] = 1.$$

For $i > 0$ and $s = 0$, consider $X(t)$ to represent the positive solution to the SDE, which is depicted in the following manner:

$$dX = \left[rX \left(1 - \frac{X}{K} \right) - \mu X - \frac{\beta X I}{1 + \alpha X} \right] dt + \sigma_1 X dB_1(t).$$

Applying the comparison theorem as delineated in Ref. 36, it can be established that for each $t \geq 0$:

$$S(t) > X(t) \quad \text{a.s.},$$

which implies

$$\mathbb{P}_{(0,i)} (S_t > 0; \text{ for each } t > 0) = 1.$$

We now transition to the discussion on bounds. To begin, let us introduce and define the subsequent function:

$$G(S, I) = (1 + S + I)^{1+\theta},$$

wherein θ represents a positive real number that satisfies

$$\mu - \frac{\theta}{2}(\sigma_1^2 \vee \sigma_2^2) > 0.$$

Applying the operator \mathcal{L} on G , gives

$$\begin{aligned} \mathcal{L}G(S, I) &= (\theta + 1)(S + I + 1)^\theta \left[rS \left(1 - \frac{S}{K} \right) - \mu(S + I) \right] \\ &\quad + \frac{\theta}{2}(\theta + 1)(S + I + 1)^{\theta-1} (\sigma_1^2 S^2 + \sigma_2^2 I^2) \\ &\leq (\theta + 1)(S + I + 1)^\theta \left[\sup_{S \in (0, \infty)} \left\{ -\frac{r}{K} S^2 + rS \right\} + \mu - \mu(S + I + 1) \right] \\ &\quad + \frac{\theta}{2}(1 + \theta)(S + I + 1)^{1+\theta} (\sigma_1^2 \vee \sigma_2^2) \\ &\leq (\theta + 1)(S + I + 1)^\theta \left[\frac{r}{4} K + \mu - \left(\mu - (\sigma_1^2 \vee \sigma_2^2) \frac{\theta}{2} \right) (S + I + 1) \right]. \end{aligned}$$

Consequently, following the selection of θ , it can be established that there are positive real numbers $\theta_1, \theta_2 > 0$ such that

$$\mathcal{L}G(S, I) \leq \theta_1 - \theta_2 G(S, I). \quad (2.5)$$

For values of n that are sufficiently large, let us establish the stopping time τ_n in the following manner:

$$\tau_n = \inf\{t \geq 0 : n \leq G(S_t, I_t)\}.$$

Using the Itô formula in conjunction with Eq. (2.5), we get

$$\begin{aligned} &\mathbb{E}_{(s,i)} [\exp(\theta_2(t \wedge \tau_n)) G(S(t \wedge \tau_n), I(t \wedge \tau_n))] \\ &\leq G(s, i) + \mathbb{E}_{(s,i)} \left(\int_0^{t \wedge \tau_n} \theta_1 \exp(\theta_2 u) du \right) \\ &\leq G(s, i) + \frac{\theta_1}{\theta_2} \exp(\theta_2 t). \end{aligned} \quad (2.6)$$

Using the Fatou lemma and letting $n \rightarrow \infty$, we get

$$\mathbb{E}_{(s,i)} [\exp(\theta_2 t) G(S(t), I(t))] \leq G(s, i) + \frac{\theta_1}{\theta_2} \exp(\theta_2 t).$$

Thus

$$\mathbb{E}_{(s,i)} [1 + S(t) + I(t)]^{1+\theta} \leq (s + i + 1)^{1+\theta} \exp(-\theta_2 t) + \frac{\theta_1}{\theta_2}.$$

1 As a result, we achieve the necessary inequality as specified in (2.2) of part (ii).

8 *S. Aznague et al.*

Conversely, based on Eq. (2.6), it is deduced that for any given $\overline{T} > 0$, the following holds:

$$\begin{aligned} n^{\theta+1} \mathbb{P}_{(s,i)} \{ \tau_n \leq \overline{T} \} &= \mathbb{E}_{(s,i)} \left[G(S(\tau_n), I(\tau_n)) \mathbb{I}_{\{ \tau_n \leq \overline{T} \}} \right] \\ &\leq \mathbb{E}_{(s,i)} \left[\exp(\theta_2 \tau_n) G(S(\tau_n), I(\tau_n)) \mathbb{I}_{\{ \tau_n \leq \overline{T} \}} \right] \\ &\leq G(s, i) + \frac{\theta_1}{\theta_2} \exp(\theta_2 \overline{T}). \end{aligned}$$

1 Consequently, we establish the requisite inequality (2.3) of part (ii). This concludes
2 the proof of Theorem 2.1. \square

We now establish the threshold that governs the system model dynamics as outlined in Eq. (1.2). Moving forward, we examine the specific equation obtained by assigning $I(t) = 0$ in the initial equation of model (1.2), and we refer to $\hat{S}(t)$ as the solution to this equation. Hence,

$$d\hat{S}(t) = \left[r\hat{S}(t) \left(1 - \frac{\hat{S}(t)}{K} \right) - \mu\hat{S}(t) \right] dt + \sigma_1 \hat{S}(t) dB_1. \quad (2.7)$$

Beginning with the initial value $\hat{S}(0)$, it is proven in Ref. 33 that the process $\hat{S}(t)$ manifests a singular stationary distribution, denoted as Θ_0 , under the condition $\frac{\sigma_1^2}{2} < r - \mu$. The density of this distribution is articulated through the utilization of the Fokker–Planck equation, as follows:

$$f^*(x) = \mathcal{H} x^{-2 + \frac{2(r-\mu)}{\sigma_1^2}} \exp\left(\frac{-2rx}{K\sigma_1^2}\right), \quad x \in (0, \infty),$$

where

$$\mathcal{H} = \left[\sigma_1^{-2} \left(\frac{K\sigma_1^2}{2r} \right)^{-1 + 2\frac{(r-\mu)}{\sigma_1^2}} \Gamma\left(-1 + 2\frac{(r-\mu)}{\sigma_1^2}\right) \right]^{-1}.$$

By ergodicity of $\hat{S}(t)$, one obtains

$$\lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t \hat{S}(x) dx = \int_0^\infty s f^*(s) ds < \infty \quad \text{a.s.}$$

In the subsequent analysis, we aim to show how the long-term dynamics of $I(t)$ are influenced by

$$\lambda = \int_0^\infty \frac{\beta s}{1 + \alpha s} \Theta_0(ds) - \mu - \gamma - \frac{\sigma_2^2}{2}.$$

Let us examine the modified version of Eq. (2.7), which is presented as follows:

$$d\bar{S}^{(\vartheta_0)}(t) = \left[r\bar{S}^{(\vartheta_0)} \left(1 - \frac{\bar{S}^{(\vartheta_0)}}{K} \right) - \mu\bar{S}^{(\vartheta_0)} + \gamma\vartheta_0 \right] dt + \sigma_1\bar{S}^{(\vartheta_0)} dB_1(t). \quad (2.8)$$

Similar to Eq. (2.7). If $r - \mu > \frac{\sigma_1^2}{2}$, then the function $\bar{S}^{(\vartheta_0)}(t)$ admits an ergodic invariant measure, which is denoted by Θ_{ϑ_0} . Thus, we have

$$\int_0^\infty \frac{\beta s}{1 + \alpha s} \Theta_{\vartheta_0}(ds) < \infty.$$

- 1 Subsequently, we introduce a crucial lemma that elucidates the connection between
2 Eqs. (2.7) and (2.8).

Lemma 2.1. *Let $\vartheta_0 \in [0, 1]$, we have*

$$\lim_{\vartheta_0 \rightarrow 0} \left| \int_0^\infty \frac{\beta s}{1 + \alpha s} \Theta_{\vartheta_0}(ds) - \int_0^\infty \frac{\beta s}{1 + \alpha s} \Theta_0(ds) \right| = 0.$$

Proof. It is easy to verify that

$$\lim_{t \rightarrow \infty} \sup \mathbb{E} \left(\bar{S}^{(\vartheta_0)}(t) \right)^{1+q} < \infty \text{ for } q > 0.$$

Consequently, the process $(\bar{S}^{(\vartheta_0)}, t \geq 0)$ exhibits uniform integrability. This characteristic implies that the family $\{\Theta_{\vartheta_0}, \vartheta_0 \in [0, 1]\}$ demonstrates tightness over the interval $(0, \infty)$. Consider an integer n such that ϑ_n , a sequence within $[0, 1]$, approaches 0 as n tends to infinity. From this, it follows that $\{\Theta_{\vartheta_n}, n \geq 0\}$ is also tight. Therefore, one can extract a subsequence that converges weakly to a probability measure Θ over the interval $(0, \infty)$. For ease of notation, this subsequence will continue to be referred to as $\{\Theta_{\vartheta_n}, n \geq 0\}$. Thus, we have

$$\lim_{n \rightarrow \infty} \int_0^\infty g(s) \Theta_{\vartheta_n}(ds) = \int_0^\infty g(s) \bar{\Theta}(ds).$$

For every smooth function, g possesses compact support. Furthermore, applying the dominated convergence theorem leads to the conclusion that

$$\begin{aligned} \int_0^\infty \mathcal{L}^{\vartheta_n} g(s) \Theta_{\vartheta_n}(ds) &= \lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t \mathbb{E} \mathcal{L}^{\vartheta_n} g \left(\bar{S}^{(\vartheta_n)}(u) \right) du \\ &= \lim_{t \rightarrow \infty} \frac{1}{t} \mathbb{E} \left[g \left(\bar{S}^{(\vartheta_n)}(t) \right) - g \left(\bar{S}^{(\vartheta_n)}(0) \right) \right] \\ &= 0, \end{aligned}$$

where $\bar{S}^{(\vartheta_n)}(t)$ denotes the invariant solution corresponding to Eq. (2.8), with the substitution of ϑ_n for ϑ_0 . It is important to observe that

$$\mathcal{L}^{\vartheta_n} g(s) = \mathcal{L}^0 g(s) + \gamma\vartheta_n,$$

which implies that

$$\int_0^\infty \mathcal{L}^0 g(s) \Theta_{\vartheta_n}(ds) + \gamma\vartheta_n = 0. \quad (2.9)$$

10 *S. Aznague et al.*

By Eq. (2.9) and considering the weak convergence of Θ_{ϑ_n} toward $\bar{\Theta}$, we obtain the following conclusion:

$$\int_0^\infty \mathcal{L}^0 g(s) \bar{\Theta}(ds) = 0.$$

1 This indicates that $\bar{\Theta}$ serves as an invariant measure for Eq. (2.7). Given its unique
 2 nature, it follows that $\bar{\Theta}$ is equivalent to Θ_0 . Consequently, Θ_{ϑ_0} exhibits weak
 3 convergence toward Θ_0 as ϑ_0 approaches zero. \square

4 3. Patterns of Extinction Dynamics

5 In this section, the assumption remains that $(r - \mu) > \frac{\sigma^2}{2}$. The ensuing finding
 6 confirms that $\lambda < 0$ represents the rate at which the count of infectious individuals
 7 approaches zero.

Lemma 3.1. *For any given values of ϵ and $M > 0$, there is a corresponding $\vartheta_1 > 0$ such that*

$$\mathbb{P}_{(s,i)} \left(\lim_{t \rightarrow \infty} I(t) = 0 \right) \geq 1 - \epsilon \text{ for each } (s, i) \in [0, M] \times [0, \vartheta_1].$$

Proof. Assuming that ϑ_0 adheres to the conditions outlined in Lemma 2.1, it then follows that

$$\left| \int_0^\infty \frac{\beta s}{1 + \alpha s} \Theta_{\vartheta_0}(ds) - \int_0^\infty \frac{\beta s}{1 + \alpha s} \Theta_0(ds) \right| \leq \frac{|\lambda|}{4}.$$

Using the ergodicity properties of $\bar{S}^{\vartheta_0}(t)$, it can be established that a certain $T_1 > 0$ exists, ensuring that for every $\epsilon > 0$, the probability $\mathbb{P}(\Omega_1) \geq 1 - \frac{\epsilon}{4}$, wherein

$$\Omega_1 = \left\{ w : \frac{1}{t} \int_0^t \frac{\beta \bar{S}^{\vartheta_0}(u)}{1 + \alpha \bar{S}^{\vartheta_0}(u)} du \leq \int_0^\infty \frac{\beta s}{1 + \alpha s} \Theta_{\vartheta_0}(ds) + \frac{|\lambda|}{4}, \ t \geq T_1 \right\}. \quad (3.1)$$

On the other hand, by the strong law of large numbers applicable to local martingale (see, e.g. Ref. 37 for more information and the references cited therein), we derive the following conclusion:

$$\lim_{t \rightarrow \infty} \frac{1}{t} B_2(t) = 0, \text{ a.s.} \quad (3.2)$$

Therefore, there is a T_2 such that the probability $\mathbb{P}(\Omega_2) \geq 1 - \frac{\epsilon}{4}$, wherein

$$\Omega_2 = \left\{ w : \frac{1}{t} |\sigma_2 B_2(t)| \leq \frac{|\lambda|}{4}, \text{ for all } t \geq T_2 \right\}. \quad (3.3)$$

Define $T = \max\{T_1, T_2\}$. Additionally, we choose a sufficiently large $C > \frac{\beta}{\alpha}T$ to ensure that the probability $\mathbb{P}(\Omega_3) \geq 1 - \frac{\epsilon}{4}$, where

$$\Omega_3 = \left\{ w : |\sigma_2 B_2(t)| \leq C - \frac{\beta}{\alpha}T \text{ for all } t \in [0, T] \right\}. \quad (3.4)$$

By analyzing the I -equation in conjunction with Eq. (3.4), it leads us to the following result:

$$\begin{aligned} I(t) &= i \exp \left\{ \int_0^t \frac{\beta S(u)}{1 + \alpha S(u)} du - \left(\mu + \gamma + \frac{\sigma_2^2}{2} \right) t + \sigma_2 B_2(t) \right\} \\ &\leq i \exp \left\{ \frac{\beta}{\alpha} t + \sigma_2 B_2(t) \right\} \\ &\leq \vartheta_1 \exp(C). \end{aligned} \quad (3.5)$$

Assume $\vartheta_1 \in [0, \vartheta_0 \exp(-C)]$. Under the condition that $I(0) = i \leq \vartheta_1$, it can be deduced that

$$I(t) < \vartheta_0 \quad \text{for all } t \in [0, T], \quad w \in \Omega_3.$$

Subsequently, let us establish the stopping time as follows:

$$\bar{\nu} = \inf\{t \geq 0 : \vartheta_0 \leq I(t)\}.$$

At this juncture, it can be shown that $\bar{\nu} = \infty$. For any $\omega \in \Omega_3$, it follows that $\bar{\nu} > T$. Consequently, considering $S(t) \leq \bar{S}^{\vartheta_0}$, it results in the following for $t \leq \bar{\nu}$:

$$\begin{aligned} I(t) &= i \exp \left\{ \int_0^t \frac{\beta S(u)}{1 + \alpha S(u)} du - \left(\mu + \gamma + \frac{\sigma_2^2}{2} \right) t + \sigma_2 B_2(t) \right\} \\ &\leq i \exp \left\{ \int_0^t \frac{\beta \bar{S}^{\vartheta_0}}{1 + \alpha \bar{S}^{\vartheta_0}} du - \left(\mu + \gamma + \frac{\sigma_2^2}{2} \right) t + \sigma_2 B_2(t) \right\}. \end{aligned} \quad (3.6)$$

Using Eqs. (3.1), (3.3), and (3.5) into Eq. (3.6), it follows that for any $\omega \in (\cap_{j=1}^3 \Omega_j)$, and for every $t \in [T, \bar{\nu})$, we deduce the following:

$$\begin{aligned} I(t) &\leq i \exp \left\{ t \int_0^\infty \frac{\beta s}{1 + \alpha s} \Theta_{\vartheta_0}(ds) + \frac{|\lambda|}{4} t - \left(\mu + \gamma + \frac{\sigma_2^2}{2} \right) t + \frac{|\lambda|}{4} t \right\} \\ &\leq i \exp \left\{ \left(\int_0^\infty \frac{\beta s}{1 + \alpha s} \Theta_0(ds) + \frac{|\lambda|}{4} \right) t + \frac{|\lambda|}{4} t - \left(\mu + \gamma + \frac{\sigma_2^2}{2} \right) t + \frac{|\lambda|}{4} t \right\} \\ &\leq i \exp \left\{ t\lambda + \frac{|\lambda|}{4} t + \frac{|\lambda|}{4} t + \frac{|\lambda|}{4} t \right\} \end{aligned}$$

12 *S. Aznague et al.*

$$\begin{aligned} &\leq i \exp \left\{ t\lambda - \frac{3\lambda}{4}t \right\} \\ &= i \exp \left\{ \frac{\lambda}{4}t \right\} < i < \vartheta_0. \end{aligned}$$

As a result, we must have $\bar{\nu} = \infty$ for $\omega \in (\cap_{j=1}^3 \Omega_j)$, when $I(0) \leq \vartheta_1$. Under these conditions, we can establish that for all $t \geq T$, and $\omega \in (\cap_{j=1}^3 \Omega_j)$, with $I(0) = i \leq \vartheta_1$, the following holds:

$$I(t) \leq i \exp \left\{ \frac{\lambda}{4}t \right\}.$$

1 Consequently, this leads to the conclusion that $\lim_{t \rightarrow \infty} I(t) = 0$. \square

Lemma 3.2. *For every ϵ and $M > 0$, there exists $\vartheta_1 > 0$ such that*

$$\mathbb{P}_{(s,i)} \left(\lim_{t \rightarrow \infty} \frac{1}{t} \ln(I(t)) = \lambda < 0 \right) \geq 1 - \epsilon, \quad \text{for each } (s,i) \in [0, M] \times [0, \vartheta_1].$$

Proof. Given that $I(t) \leq \vartheta_0$ holds for all $t \geq 0$ and for the majority of $\omega \in (\cap_{j=1}^3 \Omega_j)$, one can deduce that $S(t) \leq \bar{S}^{\vartheta_0}$. Moreover, for almost all $\omega \in (\cap_{j=1}^3 \Omega_j)$ and considering the ergodic properties of $S^{\bar{\vartheta}_0}(u)$, it is established that for a certain small $q > 0$:

$$\begin{aligned} \lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t (S(u))^{1+q} du &\leq \lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t (S^{\bar{\vartheta}_0}(u))^{1+q} du \\ &\leq \int_0^\infty s^{1+q} \Theta_{\vartheta_0}(ds) < \infty. \end{aligned} \quad (3.7)$$

Applying Lemma 3.1 and considering the uniform integrability specified in Eq. (3.7), the assortment of random occupation measures is presented in the following manner:

$$\bar{\mathcal{O}}^t(\cdot) = \frac{1}{t} \int_0^t \mathbb{I}_{\{S(u), I(u) \in (\cdot)\}} du.$$

For the majority of $\omega \in (\cap_{j=1}^3 \Omega_j)$, the set is notably tight. Moreover, one can easily prove that the process $X(t)$ possesses a distinctive invariant probability measure, $(\Theta_0 \times \delta)$, over the interval $[0, \infty] \times \{0\}$, with δ representing the Dirac measure concentrated at 0. Furthermore, it has been shown (refer to Refs. 38, 39) that as t approaches infinity, any weak limit of $\bar{\mathcal{O}}^t$ a.s. becomes an invariant probability measure, predominantly supported on $[0, \infty] \times \{0\}$. Consequently, $\bar{\mathcal{O}}^t(\cdot)$ exhibits weak convergence toward $(\Theta_0 \times \delta)$ a.s. in $(\cap_{j=1}^3 \Omega_j)$ when t tends to infinity. Following this line of weak convergence (as discussed in Ref. 38), it is concluded that

$$\lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t \frac{\beta S(u)}{1 + \alpha S(u)} du = \int_0^\infty \frac{\beta s}{1 + \alpha s} \Theta_0(ds). \quad (3.8)$$

Injecting Eq. (3.8) into (3.5), for almost every $\omega \in (\cap_{j=1}^3 \Omega_j)$, and for each pair $(s, i) \in [0, M] \times [0, \vartheta_1]$, yields

$$\begin{aligned} \lim_{t \rightarrow \infty} \frac{1}{t} \ln(I(t)) &= \int_0^\infty \frac{\beta s}{1 + \alpha s} \Theta_0(ds) - \mu - \gamma - \frac{\sigma_2^2}{2} \\ &= \lambda < 0. \end{aligned}$$

- 1 Thus, given that the probability $\mathbb{P}(\omega \in \cap_{j=1}^3 \Omega_j) > 1 - \epsilon$, we obtain the desired
2 assertion. The proof is complete. \square

Theorem 3.1. *If the threshold is $\lambda < 0$, then*

$$\mathbb{P}_{(s,i)} \left(\lim_{t \rightarrow \infty} \frac{1}{t} \ln(I(t)) = \lambda < 0 \right) = 1. \quad (3.9)$$

Proof. Consider any $\epsilon > 0$ as given. By Lemma 3.2, $X(t)$ exhibits transience within $\mathbb{R}^{(2,o)}$, leading to the absence of an invariant probability measure for $X(t)$ in $\mathbb{R}^{(2,o)}$. Consequently, $(\Theta_0 \times \delta)$ stands as the sole invariant probability measure for $X(t)$ in \mathbb{R}_+^2 . We then select M to be sufficiently large to satisfy the subsequent condition:

$$1 - \frac{\epsilon}{2} < \Theta_0 \quad (\{s \in (0, M)\}).$$

By Eq. (2.2), the process $X(t)$ exhibits tightness. Then, we can conclude that the occupation measure is characterized as follows:

$$\bar{O}_{(s,i)}^t(\cdot) = \frac{1}{t} \int_0^t \mathbb{P}_{(s,i)} [X(u) \in (\cdot)] du,$$

is tightness within the state space \mathbb{R}_+^2 . Additionally, the process $X(t)$ is characterized by an invariant measure, identified as the weak limit of $\bar{O}_{(s,i)}^t$ with the progression of t toward infinity. Therefore, it is evident that $\bar{O}_{(s,i)}^t$ converges to $(\Theta_0 \times \delta)$ as t approaches infinity. Consequently, there exists $\bar{T} > 0$ and for any given $\vartheta > 0$ such that

$$\bar{O}_{(s,i)}^{\bar{T}} [(0, H) \times (0, \vartheta)] > 1 - \epsilon.$$

In other words, we have

$$\frac{1}{\bar{T}} \int_0^{\bar{T}} \mathbb{P}_{(s,i)} \{(S_t, I_t) \in (0, M) \times (0, \vartheta)\} dt > 1 - \epsilon.$$

Therefore, upon establishing the following setting:

$$\hat{\nu} = \inf \{t \geq 0 : (S_t, I_t) \in (0, M) \times (0, \vartheta)\},$$

hence

$$\mathbb{P}_{(s,i)} \{\hat{\nu} \leq \bar{T}\} > 1 - \epsilon.$$

14 *S. Aznague et al.*

Using Lemma 3.2 in conjunction with the strong Markov property leads to the subsequent conclusion:

$$\mathbb{P}_{(s,i)} \left(\lim_{t \rightarrow \infty} \frac{1}{t} \ln(I(t)) = \lambda < 0 \right) \geq 1 - \epsilon \text{ for each } (s,i) \in \mathbb{R}_+^{(2,*)}.$$

1 Consequently, as ϵ approaches zero, we get the necessary outcome outlined in Eq.
2 (3.9). The proof is complete. \square

3 4. Ergodic Stationary Distribution

4 In this part of the study, we validate the existence of an ergodic stationary distribu-
5 tion for our epidemic model (1.2), underscoring the ongoing and significant impact
6 of disease on the host population.

7 **Theorem 4.1.** *Assuming $\lambda > 0$ and $\mu > \frac{1}{2}(\sigma_1^2 \vee \sigma_2^2)$, the system defined by Eq.*
8 *(1.2) not only has a unique stationary distribution, denoted as $v^*(\cdot)$, but is also*
9 *ergodic. Furthermore,*

(i) *For every function $f : \mathbb{R}_+^{(2,o)} \mapsto \mathbb{R}$ that is integrable with respect to v^* , it follows that*

$$\mathbb{P} \left\{ \lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t f(X(\tau)) d\tau = \int_{\mathbb{R}_+^{(2,o)}} f(x) v^*(dx) \right\} = 1.$$

(ii) *For each $(s,i) \in \mathbb{R}_+^{(2,*)}$, we have*

$$\lim_{t \rightarrow \infty} \|\mathbb{P}[t, (s,i), (\cdot)] - v^*(\cdot)\| = 0,$$

10 where $\mathbb{P}[t, (s,i), (\cdot)]$ is the transition probability.

Proof. (i) Considering the framework of system (1.2), let us define

$$V_1 = -\ln(I) + c_1[\ln(\hat{S}) - \ln(S)].$$

Applying the Itô formula, we get

$$\begin{aligned} \mathcal{L}V_1 &= \left(\mu + \gamma + \frac{\sigma_2^2}{2} \right) - \frac{\beta S}{1 + \alpha S} + c_1 \left[r \left(1 - \frac{\hat{S}}{K} \right) - r \left(1 - \frac{S}{K} \right) + \frac{\beta I}{1 + \alpha S} - \gamma \frac{I}{S} \right] \\ &\leq -\lambda + \int_0^\infty \frac{\beta s}{1 + \alpha s} \Theta_0(ds) - \frac{\beta \hat{S}}{1 + \alpha \hat{S}} + \frac{\beta \hat{S}}{1 + \alpha \hat{S}} - \frac{c_1 r}{K} \hat{S} + \frac{c_1 r}{K} S + \frac{c_1 \beta I}{1 + \alpha S} \\ &\quad - c_1 \gamma \frac{I}{S} \\ &\leq -\lambda + \int_0^\infty \frac{\beta s}{1 + \alpha s} \Theta_0(ds) - \frac{\beta \hat{S}}{1 + \alpha \hat{S}} + \left(\beta - \frac{c_1 r}{K} \right) \hat{S} + \frac{c_1 r}{K} S + c_1 \beta I - c_1 \gamma \frac{I}{S}. \end{aligned}$$

Choosing $c_1 = \frac{\beta K}{r}$, then

$$\mathcal{L}V_1 \leq -\lambda + \int_0^\infty \frac{\beta s}{1 + \alpha s} \Theta_0(ds) - \frac{\beta \hat{S}}{1 + \alpha \hat{S}} - c_1 \gamma \frac{I}{S} + \frac{c_1 r}{K} S + c_1 \beta I. \quad (4.1)$$

Moreover, we have

$$\begin{aligned} & \mathcal{L} \left(\frac{1}{m+2} (S+I)^{m+2} \right) \\ &= (S+I)^{m+1} \left[rS \left(1 - \frac{S}{K} \right) - \mu S - \mu I \right] + \frac{m+1}{2} (S+I)^m (\sigma_1^2 S^2 + \sigma_2^2 I^2) \\ &\leq rS(S+I)^{m+1} - \frac{r}{K} S^{m+3} - \mu I^{m+2} + \frac{m+1}{2} (S+I)^{m+2} (\sigma_1^2 \vee \sigma_2^2) \\ &\leq -\frac{r}{2K} S^{m+3} - m\mu I^{m+2} - \frac{r}{2K} S^{m+3} - \mu(1-m)I^{m+2} + rS(S+I)^{m+1} \\ &\quad + \frac{m+1}{2} (S+I)^{m+2} (\sigma_1^2 \vee \sigma_2^2) \\ &\leq -\frac{r}{2K} S^{m+3} - m\mu I^{m+2} + B, \end{aligned} \quad (4.2)$$

where m is a constant satisfying

$$0 < m < \frac{2\mu - \sigma_1^2 \vee \sigma_2^2}{2\mu + \sigma_1^2 \vee \sigma_2^2} \quad \text{🗨️}$$

and

$$\begin{aligned} B = \sup_{X \in \mathbb{R}_+^{(2,o)}} & \left\{ -\frac{r}{2K} S^{m+3} - \mu(1-m)I^{m+2} \right. \\ & \left. + \frac{1}{2}(m+1)(S+I)^{m+2}(\sigma_1^2 \vee \sigma_2^2) + rS(S+I)^{m+1} \right\}. \end{aligned}$$

Let $\tilde{V} \in \mathcal{C}^2(\mathbb{R}_+^{(2,o)}; \mathbb{R})$, we define

$$\tilde{V}(S, I) = \overline{M}V_1(S, I, R) + \frac{1}{m+2}(S+I)^{m+2},$$

where $\overline{M} > 0$ is chosen to be sufficiently large to satisfy the subsequent condition:

$$-\overline{M}\lambda + E \leq -2, \quad (4.3)$$

where

$$E = \sup_{S \in (0, \infty)} \left\{ -\frac{r}{4K} S^{m+3} + \frac{rc_1 M}{K} S + B \right\}.$$

Showing that $\tilde{V}(S, I)$ attains a minimum at the point (S^*, I^*) is straightforward. Subsequently, we establish a non-negative \mathcal{C}^2 -class function $W(S, I)$ in the following manner:

$$W(S, I) = \tilde{V}(S, I) - \tilde{V}(S^*, I^*).$$

16 *S. Aznague et al.*

Using Eqs. (4.1) and (4.2), we derive the following result:

$$\begin{aligned}\mathcal{LW} &\leq -\overline{M}\lambda + \frac{rc_1\overline{M}}{K}S + c_1\overline{M}\beta I - \frac{r}{2K}S^{m+3} - \mu mI^{m+2} + B - c_1\gamma\overline{M}\frac{I}{S} \\ &\quad + \overline{M}\left(\int_0^\infty \frac{\beta s}{1+\alpha s}\Theta_0(ds) - \frac{\beta\hat{S}}{1+\alpha\hat{S}}\right), \\ &= F(S, I) + \overline{M}\left(\int_0^\infty \frac{\beta s}{1+\alpha s}\Theta_0(ds) - \frac{\beta\hat{S}}{1+\alpha\hat{S}}\right),\end{aligned}$$

where

$$F(S, I) = -\overline{M}\lambda + \frac{rc_1\overline{M}}{K}S + c_1\overline{M}\beta I - \frac{r}{2K}S^{m+3} - \mu mI^{m+2} + B - c_1\gamma\overline{M}\frac{I}{S}.$$

1 Moreover, by applying the specific expression of $F(S, I)$, it leads us to the following:

Case 1. Under the conditions where either $S \rightsquigarrow 0^+$, $S \rightsquigarrow \infty$, or $I \rightsquigarrow \infty$, it follows that

$$\begin{aligned}F(S, I) &= -\overline{M}\lambda + \frac{rc_1\overline{M}}{K}S + c_1\overline{M}\beta I - \frac{r}{2K}S^{m+3} - \mu mI^{m+2} + B - c_1\gamma\overline{M}\frac{I}{S} \\ &\leq -\infty.\end{aligned}$$

Case 2. If $I \rightsquigarrow 0^+$, then it implies that

$$\begin{aligned}F(S, I) &= -\overline{M}\lambda + \frac{rc_1\overline{M}}{K}S + c_1\overline{M}\beta I - \frac{r}{2K}S^{m+3} - \mu mI^{m+2} + B - c_1\gamma\overline{M}\frac{I}{S} \\ &\leq -\overline{M}\lambda + c_1\overline{M}\beta I + E \rightsquigarrow -\overline{M}\lambda + E \\ &\leq -2.\end{aligned}$$

This conclusion is derived from Eq. (4.3). Consequently, with a sufficiently small $\epsilon > 0$, it is possible to deduce that

$$F(X) \leq -1 \text{ for each } X = (S, I) \in (\mathbb{R}_+^{(2,o)} \setminus \mathcal{D}), \quad (4.4)$$

where

$$\mathcal{D} = \left\{ (S, I) \in \mathbb{R}_+^{(2,o)} : \epsilon < S < \frac{1}{\epsilon}, \epsilon < I < \frac{1}{\epsilon} \right\}.$$

Furthermore, there exists a constant $\bar{C} > 0$ such that

$$F(X) \leq \bar{C} \text{ for all } X \in \mathbb{R}_+^{(2,o)}. \quad (4.5)$$

Therefore, it results in the following conclusion:

$$\begin{aligned}
-\mathbb{E}[W(X_0)] &\leq -\mathbb{E}[W(X_0)] + \mathbb{E}[W(X(t))] \\
&= \int_0^t \mathbb{E}[\mathcal{L}W(X(\tau))] d\tau \\
&\leq \int_0^t \mathbb{E}[F(X(\tau))] d\tau + \overline{M}\mathbb{E}\left[\int_0^t \int_0^\infty \frac{\beta s}{1+\alpha s} \Theta_0(ds) d\tau\right. \\
&\quad \left.- \int_0^t \frac{\beta \widehat{S}(\tau)}{1+\alpha \widehat{S}(\tau)} d\tau\right].
\end{aligned}$$

Consequently, by applying Eqs. (4.4), (4.5) and considering the ergodic nature of \widehat{S} , we get the following result:

$$\begin{aligned}
0 &\leq \liminf_{t \rightarrow \infty} \frac{1}{t} \int_0^t \mathbb{E}[F(X(\tau))] d\tau \\
&= \liminf_{t \rightarrow \infty} \frac{1}{t} \int_0^t [\mathbb{E}(F(X(\tau))\mathbb{I}_{\{X(\tau) \in \mathcal{D}\}}) + \mathbb{E}(F(X(\tau))\mathbb{I}_{\{X(\tau) \in \mathcal{D}^c\}})] d\tau \\
&\leq \liminf_{t \rightarrow \infty} \frac{1}{t} \int_0^t [\bar{C}\mathbb{P}(X(\tau) \in \mathcal{D}) - \mathbb{P}(X(\tau) \in \mathcal{D}^c)] d\tau \\
&= -1 + (\bar{C} + 1) \liminf_{t \rightarrow \infty} \frac{1}{t} \int_0^t \mathbb{P}(X(\tau) \in \mathcal{D}) d\tau.
\end{aligned}$$

This indicates that for every initial condition $X_0 = (s, i)$ within space $\mathbb{R}_+^{(2,o)}$,

$$\liminf_{t \rightarrow \infty} \frac{1}{t} \int_0^t \mathbb{P}(\tau, (s, i), \mathcal{D}) d\tau \geq \frac{1}{\bar{C} + 1}. \quad (4.6)$$

- 1 Given the invariance of the space $\mathbb{R}_+^{(2,o)}$ as outlined in Eq. (1.2), it is appropriate to
- 2 analyze $X(t)$ within this state space. Moreover, the invariance of $\mathbb{R}_+^{(2,o)}$, coupled with
- 3 the implications of inequality (4.6), indicates the existence of a stable probability
- 4 measure ν^* on $\mathbb{R}_+^{(2,o)}$ (refer to Ref. 40 for more information and the references cited
- 5 therein). Furthermore, the fact that $B_i(t)$ for $i = 1, 2$ are independent suggests
- 6 that $\mathbb{R}_+^{(2,o)}$ serves as the support for the measure ν^* , as detailed in Lemma 4.1 of
- 7 Khasminskii.⁴¹ In conclusion, with the insights provided in Refs. 42–44, we can
- 8 substantiate assertions (i) and (ii), thereby completing the proof. \square

9 5. Numerical Simulations

In this section, we will conduct a range of numerical simulations using the Euler-Maruyama method, as further elaborated in the referenced work (refer to Ref. 45 for further information and the references cited therein). These simulations aim

18 *S. Aznague et al.*

to represent and validate our research conclusions visually. We will focus on the discretized version of Eq. (1.2) as follows:

$$\begin{cases} S_{k+1} = S_k + \left[rS_k \left(1 - \frac{S_k}{K} \right) - \mu S_k - \frac{\beta S_k I_k}{1 + \alpha S_k} + \gamma I_k \right] \Delta t + \sigma_1 S_k \sqrt{\Delta t} \tau_{(1,k)}, \\ I_{k+1} = I_k + \left[\frac{\beta S_k I_k}{1 + \alpha S_k} - (\mu + \gamma) I_k \right] \Delta t + \sigma_2 I_k \sqrt{\Delta t} \tau_{(2,k)}. \end{cases}$$

In this formulation, $\tau_{(i,k)}$ for $i = 1, 2$ are independent Gaussian random variables, each adhering to a standard normal distribution, symbolized as $\mathcal{N}(0, 1)$. These simulations aim to offer an extensive visual depiction of the real-world applications of our key findings. Additionally, we consistently have

$$\mathbb{E}_{\Theta_0}(\hat{S}) = \int_0^\infty s \Theta_0(ds),$$

where Θ_0 exhibits absolute continuity compared to the Lebesgue measure over the $(0, \infty)$ range. Moreover, this can be further elaborated as follows:

$$\lambda = \mathbb{E}_{\Theta_0}(f(\hat{S})) - \mu - \gamma - \frac{\sigma_2^2}{2}.$$

- 1 Here, $f(\hat{S}) = \beta \hat{S} / (1 + \alpha \hat{S})$. Using the Monte-Carlo method enables the approxima-
- 2 tion of the expected value $\mathbb{E}_{\Theta_0}(f(\hat{S}))$ at a specific time t . With this methodology,
- 3 the following examples are presented:

Example 1. In system (1.2), we choose $K = 1.5$, $r = 1.0$, $\beta = 0.3$, $\mu = 0.2$, $\gamma = 0.2$, $\alpha = 0.2$, $\sigma_1 = 0.4$, $\sigma_2 = 0.45$, $(S_0, I_0) = (0.8, 0.6)$. These values make the assumption in Theorem 3.1 satisfied. Namely,

$$r - \mu = 0.8 > \frac{\sigma_1^2}{2} \quad \text{and} \quad \lambda \approx -0.13 < 0.$$

- 4 Consequently, the condition for extinction outlined in Theorem 3.1 is met. This
- 5 outcome is validated through computer simulations illustrated in Fig. 1.




Fig. 1. Dynamics of the solution $(S(t), I(t))$ with initial condition $(S_0, I_0) = (0.8, 0.6)$, indicating that the disease dies out when $\lambda < 0$.

Example 2. Choosing $K = 1.5$, $r = 1.0$, $\beta = 0.6$, $\mu = 0.2$, $\gamma = 0.3$, $\alpha = 0.2$, $\sigma_1 = 0.1$, $\sigma_2 = 0.15$. By direct calculation, we have

$$\mu = 0.2 > \frac{1}{2}(\sigma_1^2 \vee \sigma_2^2) \quad \text{and} \quad \lambda \approx 0.22 > 0.$$

According to Theorem 4.1, it is established that a unique ergodic stationary distribution, represented as $v^*(\cdot)$, exists for the system detailed in Eq. (1.2). This is demonstrated in Fig. 3. Furthermore, the persistence of the solution for the stochastic model (1.2) is evident in its mean behavior, as depicted in Fig. 2.

The numerical simulations presented in this section exhibit strong concordance with the theoretical predictions. In Example 1, where $\lambda < 0$, the conditions for disease extinction are satisfied, and the results shown in Fig. 1 confirm that the epidemic eventually vanishes. In contrast, in Example 2, where $\lambda > 0$, the persistence condition is met, and the simulation results illustrated in Figs. 2 and 3

AQ: Figures 2 and 3 are out of sequence in the text citation. Please check.




Fig. 2. Dynamics of the solution $(S(t), I(t))$ with initial condition $(S_0, I_0) = (0.8, 0.6)$, indicating that the disease persists when $\lambda > 0$.




Fig. 3. Kernel density functions of $S(t)$ and $I(t)$ for the SDE described in (1.2).

20 *S. Aznague et al.*

1 indicate that the system converges asymptotically to a unique ergodic stationary
2 distribution. These numerical findings provide strong empirical support for the the-
3 oretical analysis, reinforcing the validity of the extinction and persistence conditions
4 established in Theorems 3.1 and 4.1.

5 **6. Concluding Remarks**

Our study thoroughly explored a stochastic SIS epidemic model, integrating logistic growth and saturation incidence elements. Our comprehensive research has yielded a significant understanding of the disease transmission dynamics. We have delineated crucial criteria that govern whether the disease persists or is eradicated within this sophisticated model. Our findings indicate that a negative value of λ leads to disease extinction, while a positive value implies persistence. To concisely represent this pivotal discovery, we introduce the parameter \mathcal{R}_0^s , which is defined in the following manner:

$$\mathcal{R}_0^s := \left(\mu + \gamma + \frac{\sigma_2^2}{2} \right)^{-1} \int_0^\infty \frac{\beta s}{1 + \alpha s} \Theta_0(ds) = \lambda \left(\mu + \gamma + \frac{\sigma_2^2}{2} \right)^{-1} + 1.$$

6 Our research has established that \mathcal{R}_0^s acts as the critical reproduction number in the
7 stochastic system (1.2), providing a coherent and meaningful interpretation. When
8 \mathcal{R}_0^s falls below one ($\mathcal{R}_0^s < 1$), it indicates a trend toward disease extinction, whereas
9 a value exceeding one ($\mathcal{R}_0^s > 1$) denotes the likelihood of disease persistence. This
10 parameter, \mathcal{R}_0^s , is essential for analyzing and comprehending the key processes that
11 dictate the disease trajectory within the studied framework.

12 **Acknowledgments**

13 The authors express sincere gratitude to the anonymous reviewers for their invaluable
14 input and constructive recommendations, which have significantly enhanced
15 the quality and rigor of our research endeavor.

16 **Funding**

17 This investigation was conducted independently, without receiving any external
18 funding or financial assistance.

19 **Appendix A**

Proof of Theorem 2.1. Note that the coefficients in system (1.2) exhibit local Lipschitz continuity. Hence, given any initial value $X(0) \in \mathbb{R}_+^2$, there exists a single, unique local solution for $t \in [0, \nu_e]$ almost surely, where ν_e represents the explosion time (for additional details, refer to Ref. 37). Let $m_0 \geq 1$ be such that S_0 and I_0 all lie within the interval $[\frac{1}{m_0}, m_0]$. We define a stopping time as per Ref. 37 for each

integer $m \geq m_0$

$$\nu_m = \inf \left\{ 0 \leq t < \nu_e : \frac{1}{m} \geq \min\{S_t, I_t\} \text{ or } m \leq \max\{S_t, I_t\} \right\}.$$

Evidently, ν_m is increasing as $m \rightarrow \infty$. Denote $\nu_\infty = \lim_{m \rightarrow \infty} \nu_m$, from which we deduce $\nu_\infty \leq \nu_e$ a.s. We need to prove that $\nu_\infty = \infty$ a.s. Suppose instead that

$$\mathbb{P}\{\nu_\infty < \infty\} > 0.$$

Then, there exists $T > 0$ and $\kappa \in (0, 1)$ such that

$$\mathbb{P}\{\nu_\infty \leq T\} \geq \kappa.$$

Thus, there exists $m_1 \geq m_0$, where $\mathbb{P}\{\nu_m \leq T\} \geq \kappa$. There is an integer $m \geq m_1$, denote $\Omega_m = \{\nu_m \leq T\}$, then

$$\mathbb{P}(\Omega_m) \geq \kappa. \quad (6.7)$$

Now, consider the Lyapunov function $\hat{V} \in C^2(\mathbb{R}_+^2; \mathbb{R}_+)$ given by

$$\hat{V}(X) = \left[S - a \ln \left(\frac{S}{a} \right) - a \right] + [I - \ln(I) - 1],$$

where a is a positive constant to be determined later. Applying Itô's formula, we obtain

$$d\hat{V}(X) = \mathcal{L}\hat{V}dt + \sigma_1(S - a)dB_1 + \sigma_2(I - 1)dB_2,$$

where

$$\mathcal{L}\hat{V} \leq \frac{-rS^2}{K} + \frac{(K+a)r}{K}S + \frac{a\beta I}{1+\alpha S} + a\mu - \mu I + \gamma + 2\mu + \frac{a\sigma_1^2}{2} + \frac{\sigma_2^2}{2}.$$

Choosing $a = \frac{\mu}{\beta}$, we obtain

$$\mathcal{L}\hat{V} \leq \hat{K},$$

where \hat{K} is a positive constant. Consequently,

$$d\hat{V}(X) \leq \hat{K}dt + \sigma_1(S - a)dB_1 + \sigma_2(I - 1)dB_2. \quad (6.8)$$

Integrating inequality (6.8) over $(0, \nu_m \wedge T)$ and taking expectations, we get

$$\mathbb{E}\hat{V}(X(\nu_m \wedge T)) \leq \hat{V}(X(0)) + \hat{K}T.$$

For each $\omega \in \Omega_m$, either $S(\nu_m, \omega)$ or $I(\nu_m, \omega)$ equals either m or $\frac{1}{m}$. Hence

$$\begin{aligned} \hat{V}(X(\nu_m, \omega)) &\geq \kappa \left(\left[m - a - a \ln \frac{m}{a} \right] \wedge \left[\frac{1}{m} - a + a \ln(am) \right] \wedge [m - 1 - \ln(m)] \right. \\ &\quad \left. \wedge \left[\frac{1}{m} - 1 + \ln(m) \right] \right). \end{aligned}$$

22 *S. Aznague et al.*

By letting $m \rightarrow \infty$, we get

$$\infty > \hat{V}(X(0)) + \hat{K}T = \infty.$$

This is a contradiction. Hence

$$\nu_{\infty} = \infty \quad \text{a.s.}$$

1 The proof is completed. □

2 References

- 3 1. W. O. Kermack and A. G. McKendrick, A contribution to the mathematical theory of
4 epidemics, *Proceedings of the Royal Society of London, Series A, Containing Papers*
5 *of a Mathematical and Physical Character* **115**(772) (1927) 700–721.
- 6 2. A. Gray, D. Greenhalgh, L. Hu, X. Mao and J. Pan, A stochastic differential equation
7 SIS epidemic model, *SIAM Journal on Applied Mathematics* **71** (2011) 876–902.
- 8 3. S. Cai, Y. Cai and X. Mao, A stochastic differential equation SIS epidemic model with
9 two independent Brownian motions, *Journal of Mathematical Analysis and Applica-*
10 *tions* **474** (2019) 1536–1550.
- 11 4. Y. Zhao and D. Jiang, The threshold of a stochastic SIS epidemic model with vacci-
12 nation, *Applied Mathematics and Computation* **243** (2014) 718–727.
- 13 5. S. Li and S. Guo, Permanence and extinction of a stochastic SIS epidemic model with
14 three independent Brownian motions, *Discrete & Continuous Dynamical Systems-B*
15 **26**(5) (2021) 2693–2719.
- 16 6. A. Settati, A. Lahrouz, M. El Fatini, A. El Haitami, M. El Jarroudi and M. Erri-
17 ani, A Markovian switching diffusion for an SIS model incorporating Lévy processes,
18 *Discrete and Continuous Dynamical Systems-B* **28**(1) (2022) 209–229.
- 19 7. A. Settati, A. Lahrouz, M. Zahri, A. Tridane, M. El Fatini, H. El Mahjour and
20 M. Seaid, A stochastic threshold to predict extinction and persistence of an epidemic
21 SIRS system with a general incidence rate, *Chaos, Solitons & Fractals* **144** (2021)
22 110690.
- 23 8. A. Settati, A. Lahrouz, A. Assadouq, M. El Fatini, M. El Jarroudi and K. Wang, The
24 impact of nonlinear relapse and reinfection to derive a stochastic threshold for SIRI
25 epidemic model, *Chaos, Solitons & Fractals* **137** (2020) 109897.
- 26 9. A. El Haitami, A. Settati, A. Lahrouz, M. El Idrissi and M. El Merzguioui, A stochastic
27 switched SIRI epidemic model integrating nonlinear relapse phenomena, *International*
28 *Journal of Dynamics and Control* (2023) 1–15.
- 29 10. I. Bouzalmat, M. El Idrissi, A. Settati and A. Lahrouz, Stochastic SIRS epidemic
30 model with perturbation on immunity decay rate, *Journal of Applied Mathematics*
31 *and Computing* (2023) 1–26.
- 32 11. M. El Idrissi, B. Harchaoui, A. N. Brahim, I. Bouzalmat, A. Settati and A. Lahrouz, A
33 sufficient condition for extinction and stability of a stochastic SIS model with random
34 perturbation, *WSEAS Transactions on Systems* **21** (2022) 367–371.
- 35 12. T. Caraballo, A. Settati, M. El Fatini, A. Lahrouz and A. Imlahi, Global stability and
36 positive recurrence of a stochastic SIS model with Lévy noise perturbation, *Physica*
37 *A: Statistical Mechanics and Its Applications* **523** (2019) 677–690.
- 38 13. Q. Liu, D. Jiang, T. Hayat and A. Alsaedi, Threshold behavior in a stochastic delayed
39 SIS epidemic model with vaccination and double diseases, *Journal of the Franklin*
40 *Institute* **356**(13) (2019) 7466–7485.
- 41 14. J. Liu, L. Chen and F. Wei, The persistence and extinction of a stochastic SIS epidemic
42 model with logistic growth, *Advances in Difference Equations* **2018** (2018) 68.



AQ: Please provide the volume number for Refs. 9, 10, 26, 34 and 35.

- 1 15. N. D. Phu, D. O'Regan and T. D. Tuong, Longtime characterization for the general
2 stochastic epidemic SIS model under regime switching, *Nonlinear Analysis: Hybrid*
3 *Systems* **38** (2020) 100951.
- 4 16. H. Khan, R. N. Mohapatra, K. Vajravelu and S. J. Liao, The explicit series solution of
5 SIR and SIS epidemic models, *Applied Mathematics and Computation* **215**(2) (2009)
6 653–669.
- 7 17. T. Harko, F. S. Lobo and M. Mak, Exact analytical solutions of the Susceptible-
8 Infected-Recovered (SIR) epidemic model and the SIR model with equal death and
9 birth rates, *Applied Mathematics and Computation* **236** (2014) 184–194.
- 10 18. V. Capasso, *Mathematical Structures of Epidemic Systems* (Springer, Berlin, 1993).
- 11 19. A. M. Fofana and A. Hurford, Mechanistic movement models to understand epidemic
12 spread, *Philosophical Transactions of the Royal Society, Biological Sciences* **372**(1719)
13 (2017) 20160086.
- 14 20. G. Macdonald, Epidemiologic models in studies of vector-borne diseases: The RE Dyer
15 lecture, *Public Health Reports* **76**(9) (1961) 753.
- 16 21. R. M. Anderson and R. M. May, Regulation and stability of host-parasite population
17 interactions: I. Regulatory processes, *The Journal of Animal Ecology* **47**(1) (1978)
18 219–247.
- 19 22. R. M. Anderson, H. C. Jackson, R. M. May and A. D. M. Smith, Population dynamics
20 of fox rabies in Europe, *Nature* **289** (1981) 765–771.
- 21 23. A. M. Spagnuolo, M. Shillor, L. Kingsland, A. Thatcher, M. Toeniskoetter and
22 B. Wood, A logistic delay differential equation model for Chagas disease with inter-
23 rupted spraying schedules, *Journal of Biological Dynamics* **6** (2012) 377–394.
- 24 24. N. T. Dieu, D. H. Nguyen, N. H. Du and G. Yin, Classification of asymptotic behavior
25 in a stochastic SIR model, *SIAM Journal on Applied Dynamical Systems* **15** (2016)
26 1062–1084.
- 27 25. A. N. Brahim, B. Harchaoui, S. Boutouil, M. El Idrissi, S. Aznague, A. Settati,
28 A. Lahrouz and M. El Jarroudi, Investigating stochastic dynamics of the Gilpin-Ayala
29 model in dispersed polluted environments, *WSEAS Transactions on Mathematics* **22**
30 (2023) 607–620.
- 31 26. S. Aznague, B. Harchaoui, A. N. Brahim, A. Settati, A. Lahrouz, T. Amtout and
32 M. El Idrissi, A stochastic SIR epidemic with non-linear power functions and logistic
33 growth, *New Mathematics and Natural Computation* (2024) 1–20.
- 34 27. X. Zhang and Q. Yang, Threshold behavior in a stochastic SVIR model with general
35 incidence rates, *Applied Mathematics Letters* **121** (2021) 107403.
- 36 28. D. H. Nguyen, G. Yin and C. Zhu, Long-term analysis of a stochastic SIRS model with
37 general incidence rates, *SIAM Journal on Applied Mathematics* **80** (2020) 814–838.
- 38 29. P. F. Verhulst, Notice sur la loi que la population suit dans son accroissement, *Cor-*
39 *respondance Mathématique et Physique* **10** (1838) 113–126.
- 40 30. L. D. Frank, An analysis of the effect of the economic situation on modeling and fore-
41 casting the diffusion of wireless communications in Finland, *Technological Forecasting*
42 *and Social Change* **71**(4) (2004) 391–403.
- 43 31. S. Aznague, M. El Idrissi, A. N. Brahim, B. Harchaoui, S. Boutouil, A. Settati,
44 A. Lahrouz, M. El Merzguioui and J. El Amrani, A probabilistic SIRI epidemic model
45 incorporating incidence capping and logistic population expansion, *Applied Mathe-*
46 *matics and Information Sciences* **17**(5) (2023) 773–789.
- 47 32. K. Roosa, Y. Lee, R. Luo, A. Kirpich, R. Rothenberg, J. M. Hyman and G. Chowell,
48 Real-time forecasts of the COVID-19 epidemic in China from February 5th to February
49 24th, *Infectious Disease Modelling* **5** (2020) 256–263.



24 *S. Aznague et al.*

- 1 33. Z. Cao, W. Feng, X. Wen and L. Zu, Dynamical behavior of a stochastic SEI epidemic
2 model with saturation incidence and logistic growth, *Physica A: Statistical Mechanics*
3 *and Its Applications* **523** (2019) 894–907.
- 4 34. W. Zhao, J. Liu, M. Chi and F. Bian, Dynamics analysis of stochastic epidemic models
5 with standard incidence, *Advances in Difference Equations* (2019) 1–16.
- 6 35. D. Nguyen, N. Nguyen and G. Yin, General nonlinear stochastic systems motivated by
7 chemostat models: Complete characterization of long-time behavior, optimal controls,
8 and applications to wastewater treatment, *Stochastic Processes and Applications* **130**
9 (2020) 4608–4642.
- 10 36. S. Peng and X. Zhu, Necessary and sufficient condition for comparison theorem of
11 1-dimensional stochastic differential equations, *Stochastic Processes and Their Appli-*
12 *cations* **116** (2006) 370–380.
- 13 37. C. Vargas-De-León, On the global stability of infectious disease models with relapse,
14 *Abstraction and Application* **9** (2013) 50–61.
- 15 38. A. Hening and D. Nguyen, Coexistence and extinction for stochastic Kolmogorov
16 systems, *Annals of Applied Probability* **28** (2018) 1893–1942.
- 17 39. S. J. Schreiber, M. Benaïm and K. A. S. Atchadé, Persistence in fluctuating environ-
18 ments, *Journal of Mathematical Biology* **62**(5) (2011) 655–683.
- 19 40. S. P. Meyn and R. L. Tweedie, Stability of Markovian processes III: Foster-Lyapunov
20 criteria for continuous-time processes, *Advances in Applied Probability* **25** (1993) 518–
21 548.
- 22 41. R. Khasminski, *Stochastic Stability of Differential Equations* (Springer, Berlin, 2012).
- 23 42. N. H. Du, N. H. Dang and G. Yin, Conditions for permanence and ergodicity of certain
24 stochastic predator-prey models, *Journal of Applied Probability* **53** (2016) 187–202.
- 25 43. K. Ichihara and H. Kunita, A classification of the second order degenerate elliptic
26 operators and its probabilistic characterization, *Zeitschrift für Wahrscheinlichkeits-*
27 *theorie und Verwandte Gebiete* **39** (1977) 81–84.
- 28 44. R. Khasminski, Ergodic properties of recurrent diffusion processes and stabilization of
29 the Cauchy problem for parabolic equations, *Theory of Probability and Its Applications*
30 **5** (1960) 179–196.
- 31 45. D. J. Higham, An algorithmic introduction to numerical simulation of stochastic dif-
32 ferential equations, *SIAM Review* **43**(3) (2001) 525–546.

