


<b>plan</b>	<b>0</b>	<b>1</b>	<b>2</b>	<b>3</b>
$a_0$	(0,0)	(1,0)	(2,0)	(3,0)
$a_1$	(0,0)	(1,0)	(2,0)	(3,0)




$$\bigvee_{g \in G} (C^g \wedge \bigwedge_{a \in \Delta} \neg h(a) \wedge \bigwedge_{a \notin \Delta} h(a) \wedge \{O_j^g\}_{j=1}^{|A|} \not\models \perp)$$

$$\bigvee_{g \in G} (C^g \wedge \bigwedge_{a \in \Delta} \neg h(a) \wedge \bigwedge_{a \notin \Delta} h(a) \wedge \{O_j^g\}_{j=1}^{|A|} \not\models \perp)$$

$$\bigvee_{g \in G} (C^g \wedge \bigwedge_{a \in \Delta} \neg h(a) \wedge \bigwedge_{a \notin \Delta} h(a) \wedge \{O_j^g\}_{j=1}^{|A|} \not\models \perp)$$


$$\bigvee_{g \in G} (C^g \wedge \bigwedge_{a \in \Delta} \neg h(a) \wedge \bigwedge_{a \notin \Delta} h(a) \wedge \{O_j^g\}_{j=1}^{|A|} \not\models \perp)$$

$$\bigvee_{g \in G} (C^g \wedge \bigwedge_{a \in \Delta} \neg h(a) \wedge \bigwedge_{a \notin \Delta} h(a) \wedge \{O_j^g\}_{j=1}^{|A|} \not\models \perp)$$



---

**Algorithm 1** An algorithm with caption

[illegible]

<b>plan</b>	<b>0</b>	<b>1</b>	<b>2</b>	<b>3</b>
$a_0$	(0,0)	(1,0)	(2,0)	(3,0)
$a_1$	(0,0)	(1,0)	(2,0)	(3,0)

---

**Algorithm 2** An algorithm with caption

[illegible]



Figure 4: Speak rench later astronomical traditions that have constructs that allow you Through cha