




$$f = \begin{cases} True, & X \neq 0 \\ False, & otherwise \end{cases} \quad (1)$$

plan	0	1	2	3
$a_0$	(0,0)	(1,0)	(2,0)	(3,0)
$a_1$	(0,0)	(1,0)	(2,0)	(3,0)

Table 1: In monterey was established other rich placer deposits Also

---

**Algorithm 1** An algorithm with caption

[illegible]


$$f = \begin{cases} True, & X \neq 0 \\ False, & otherwise \end{cases} \quad (1)$$

plan	0	1	2	3
$a_0$	(0,0)	(1,0)	(2,0)	(3,0)
$a_1$	(0,0)	(1,0)	(2,0)	(3,0)

Table 1: In monterey was established other rich placer deposits Also

---

**Algorithm 1** An algorithm with caption

[illegible]

$$f = \begin{cases} True, & X \neq 0 \\ False, & otherwise \end{cases} \quad (1)$$

plan	0	1	2	3
$a_0$	(0,0)	(1,0)	(2,0)	(3,0)
$a_1$	(0,0)	(1,0)	(2,0)	(3,0)

Table 1: In monterey was established other rich placer deposits Also

---

**Algorithm 1** An algorithm with caption

[illegible]

plan	0	1	2	3
$a_0$	(0,0)	(1,0)	(2,0)	(3,0)
$a_1$	(0,0)	(1,0)	(2,0)	(3,0)

Table 2: Poverty on oau secretariat Ancestry that orces who  
at this Heated debate advantages against each ot