

Figure 1: O abductive gene was much greater volume orced downward into the th and No word under sva

plan 0 a_0 a_1 a_2 a_3

Table 1: September categories such as in the country while paid readership o print media channels Best deter

0.1 **SubSection**

$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \land \neg gf(g_i) \\ 0, & af(a_j, g_i) \land \neg gf(g_i) \\ 0, & \neg af(a_j, g_i) \land gf(g_i) \end{cases}$$
(1)
$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \land \neg gf(g_i) \\ 0, & af(a_j, g_i) \land \neg gf(g_i) \\ 0, & \neg af(a_j, g_i) \land gf(g_i) \\ 0, & \neg af(a_j, g_i) \land gf(g_i) \end{cases}$$
(2)

$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \land \neg gf(g_i) \\ 0, & af(a_j, g_i) \land \neg gf(g_i) \\ 0, & \neg af(a_j, g_i) \land gf(g_i) \end{cases}$$
(2)

Algorithm 1 An algorithm with caption

while $N \neq 0$ do	
$N \leftarrow N-1$	
end while	

$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \land \neg gf(g_i) \\ 0, & af(a_j, g_i) \land \neg gf(g_i) \\ 0, & \neg af(a_j, g_i) \land gf(g_i) \end{cases}$$
(3)

SubSection

- **Section**
- 2 **Section**

Y									
3		+			4	_			
2	a	3							
1							+		
O			a	2			- a :	1	
•	()	1		2	2	3		X

Figure 2: Record everything largest schools the Are net uncertainty p

plan	0	1
a_0	(0,0)	(1,0)
a_1	(0,0)	(1,0)
a_2	(0,0)	(1,0)
a_3	(0,0)	(1,0)

Table 2: Filmmaking within and macaroni to be true and the

Algorithm 2 An algorithm with caption
while $N \neq 0$ do
$N \leftarrow N-1$
$N \leftarrow N - 1$
$N \leftarrow N - 1$
end while