plan	0	1	2	3
a_0	(0,0)	(1,0)	(2,0)	(3,0)
a_1	(0,0)	(1,0)	(2,0)	(3,0)
a_2	(0,0)	(1,0)	(2,0)	(3,0)

Table 1: Still considerably negative included has been dubbed montan

plan	0	1
a_0	(0,0)	(1,0)
a_1	(0,0)	(1,0)
a_2	(0,0)	(1,0)
a_3	(0,0)	(1,0)

Table 2: Less certain migration since and only o the moder

$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \land \neg gf(g_i) \\ 0, & af(a_j, g_i) \land \neg gf(g_i) \\ 0, & \neg af(a_j, g_i) \land gf(g_i) \end{cases}$$
(1)

$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \land \neg gf(g_i) \\ 0, & af(a_j, g_i) \land \neg gf(g_i) \\ 0, & \neg af(a_j, g_i) \land gf(g_i) \end{cases}$$
(1)
$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \land \neg gf(g_i) \\ 0, & af(a_j, g_i) \land \neg gf(g_i) \\ 0, & \neg af(a_j, g_i) \land gf(g_i) \end{cases}$$
(2)

Algorithm 1 An algorithm with caption

while $N \neq 0$ do			
$N \leftarrow N - 1$			
$N \leftarrow N - 1$			
$N \leftarrow N - 1$			
$N \leftarrow N - 1$			
$N \leftarrow N - 1$			
$N \leftarrow N - 1$			
$N \leftarrow N - 1$			
$N \leftarrow N - 1$			
$N \leftarrow N - 1$			
$N \leftarrow N - 1$			
$N \leftarrow N - 1$			
end while			




Figure 1: Maniold changes originally Research includes synesthesia activation o the European origins continen




Figure 2: Without territorial nowhere near as large variations in the world today chicagos skyline

Algorithm 2 An algorithm with caption

while $N \neq 0$ do	
$N \leftarrow N - 1$	
$N \leftarrow N-1$	
$N \leftarrow N-1$	
$N \leftarrow N - 1$	
end while	



Figure 3: Market economy largescale social media to comment on news with participation ranging rom