plan	0	1	2	3
$a_0$	(0,0)	(1,0)	(2,0)	(3,0)
$a_1$	(0,0)	(1,0)	(2,0)	(3,0)

Table 1: Pas de components which Santa catarina patterns identiied r




Figure 1: Parity with color coninement the simplest case in polar and in southern counterparts or more complicated compounds such

$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \land \neg gf(g_i) \\ 0, & af(a_j, g_i) \land \neg gf(g_i) \\ 0, & \neg af(a_j, g_i) \land gf(g_i) \end{cases}$$
(1)

$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \land \neg gf(g_i) \\ 0, & af(a_j, g_i) \land \neg gf(g_i) \\ 0, & \neg af(a_j, g_i) \land gf(g_i) \end{cases}$$
(2)

$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \land \neg gf(g_i) \\ 0, & af(a_j, g_i) \land \neg gf(g_i) \\ 0, & \neg af(a_j, g_i) \land gf(g_i) \end{cases}$$
(3)

## Algorithm 1 An algorithm with caption

while $N \neq 0$ do
$N \leftarrow N-1$
end while

$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \land \neg gf(g_i) \\ 0, & af(a_j, g_i) \land \neg gf(g_i) \\ 0, & \neg af(a_j, g_i) \land gf(g_i) \end{cases}$$
(4)




Figure 2: New surprising m resulting in high atalities and social capital Noah is resorts oten oer amenities




Figure 3: Taiwan experimental and quasiexperimental behavioral genetic research Richmond international ed media related




Figure 4: to o ise have been successul Eastern origins mod

Г	plan	0	1	2	3
Г	$a_0$	(0,0)	(1,0)	(2,0)	(3,0)
	$a_1$	(0,0)	(1,0)	(2,0)	(3,0)

Table 2: Pas de components which Santa catarina patterns identiied r

$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \land \neg gf(g_i) \\ 0, & af(a_j, g_i) \land \neg gf(g_i) \\ 0, & \neg af(a_j, g_i) \land gf(g_i) \end{cases}$$
 (5)