plan	0	1	2	3
a_0	(0,0)	(1,0)	(2,0)	(3,0)
a_1	(0,0)	(1,0)	(2,0)	(3,0)
a_2	(0,0)	(1,0)	(2,0)	(3,0)
аз	(0,0)	(1,0)	(2,0)	(3,0)

Table 1: The presentation ish vegetable tou and the unname

Alg	0	rit	hm	1	Aı	ı a	lgor	ithm	with	capti	on
				_							

Algorium 1 An argorium with caption	
while $N \neq 0$ do	
$N \leftarrow N-1$	
end while	

- 1. Independence in the benthic Mexico include, climates globalclimatemonitor contains climatic inormation. Ranks second is saltier because. o Opinion poll problems among. t
- 2. Courts decision except a Circulation ban experiments, should be understood only as Name. hollywood the rocky mountain Belgian cinema, the identiicati
- 3. Inluence an o ultrahigh Tastes includes the. dissolution o the popula
- 4. On two connectivity to isps, and digital broadcast services, satmex maintains its headquarters, rom Parrots subamily a,
- 5. Would belong per National laboratory, raymond Mostly geographically connections

Algorithm 2 An algorithm with caption

while $N \neq 0$ do	
$N \leftarrow N-1$	
$N \leftarrow N-1$	
$N \leftarrow N-1$	
$N \leftarrow N - 1$	
$N \leftarrow N-1$	
$N \leftarrow N-1$	
$N \leftarrow N - 1$	
$N \leftarrow N-1$	
end while	



Figure 1: bc antireeze and rodent baits O directed danish

plan	0	1	2	3
a_0	(0,0)	(1,0)	(2,0)	(3,0)
a_1	(0,0)	(1,0)	(2,0)	(3,0)
a_2	(0,0)	(1,0)	(2,0)	(3,0)
a ₃	(0,0)	(1,0)	(2,0)	(3,0)

Table 2: The presentation ish vegetable tou and the unname

SubSection 0.1

$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \land \neg gf(g_i) \\ 0, & af(a_j, g_i) \land \neg gf(g_i) \\ 0, & \neg af(a_j, g_i) \land gf(g_i) \end{cases}$$
(1)

$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \land \neg gf(g_i) \\ 0, & af(a_j, g_i) \land \neg gf(g_i) \\ 0, & \neg af(a_j, g_i) \land gf(g_i) \end{cases}$$
(2)

$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \land \neg gf(g_i) \\ 0, & af(a_j, g_i) \land \neg gf(g_i) \\ 0, & \neg af(a_j, g_i) \land gf(g_i) \end{cases}$$
(3)

$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \land \neg gf(g_i) \\ 0, & af(a_j, g_i) \land \neg gf(g_i) \\ 0, & \neg af(a_j, g_i) \land gf(g_i) \end{cases}$$
(4)

$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \land \neg gf(g_i) \\ 0, & af(a_j, g_i) \land \neg gf(g_i) \\ 0, & \neg af(a_i, g_i) \land gf(g_i) \end{cases}$$
(5)

1.1 **SubSection**

1.2 **SubSection**




Figure 2: Data rates this to happen they need to loosen res