

Figure 1: ulltext provides lower division coursework as Th

plan	0	1	2	3
a_0	(0,0)	(1,0)	(2,0)	(3,0)
a_1	(0,0)	(1,0)	(2,0)	(3,0)

Table 1: As showed york institute o Greatly improved appro

0.1 SubSection

$$\frac{1+\frac{c}{b}}{1+\frac{1}{1+\frac{1}{a}}}$$

$$spct_{i,j} = \begin{cases}
1, & \neg af(a_j, g_i) \land \neg gf(g_i) \\
0, & af(a_j, g_i) \land \neg gf(g_i) \\
0, & \neg af(a_j, g_i) \land gf(g_i)
\end{cases} \tag{1}$$

0.2 SubSection

$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \land \neg gf(g_i) \\ 0, & af(a_j, g_i) \land \neg gf(g_i) \\ 0, & \neg af(a_j, g_i) \land gf(g_i) \end{cases}$$

$$\frac{1 + \frac{a}{b}}{1 + \frac{1}{1 + \frac{1}{a}}}$$

$$\frac{1 + \frac{a}{b}}{1 + \frac{1}{1 + \frac{1}{1}}}$$

1 Section

2 Section

2.1 SubSection

plan	0	1	2	3
a_0	(0,0)	(1,0)	(2,0)	(3,0)
a_1	(0,0)	(1,0)	(2,0)	(3,0)

Table 2: As showed york institute o Greatly improved appro




Figure 2: ulltext provides lower division coursework as Th



Figure 3: Arab inluences them an approach which was th in \ensuremath{s}



Figure 4: Arab inluences them an approach which was th in

Algorithm 1 An algorithm with caption	
while $N \neq 0$ do	
$N \leftarrow N-1$	
$N \leftarrow N - 1$	
$N \leftarrow N-1$	
end while	