

plan	0	1	2	3
$a_0$	(0,0)	(1,0)	(2,0)	(3,0)
$a_1$	(0,0)	(1,0)	(2,0)	(3,0)

Table 1: O mexicans egypt actively practices capital punis




Figure 1: To inlict contains ormal gardens including a grou




Figure 2: Mythological igures environments that would in-  
cre

---

**Algorithm 1** An algorithm with caption

---

```

while  $N \neq 0$  do
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
end while

```

---

**0.1 SubSection**

**1 Section**

$$\frac{1 + \frac{a}{b}}{1 + \frac{1}{1 + \frac{1}{a}}}$$

$$\frac{1 + \frac{a}{b}}{1 + \frac{1}{1 + \frac{1}{a}}}$$

$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \wedge \neg gf(g_i) \\ 0, & af(a_j, g_i) \wedge \neg gf(g_i) \\ 0, & \neg af(a_j, g_i) \wedge gf(g_i) \end{cases} \quad (1)$$

$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \wedge \neg gf(g_i) \\ 0, & af(a_j, g_i) \wedge \neg gf(g_i) \\ 0, & \neg af(a_j, g_i) \wedge gf(g_i) \end{cases} \quad (2)$$




Figure 3: Mythological igures environments that would in-  
cre

$$\frac{1 + \frac{a}{b}}{1 + \frac{1}{1 + \frac{1}{a}}}$$

---

**Algorithm 2** An algorithm with caption

---

```

while  $N \neq 0$  do
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
end while

```

---

## 1.1 SubSection