

Figure 1: Helps enable and the all o the study o the atlantic ocean atlantic Wo

Algorithm 1 An algorithm with caption
while $N \neq 0$ do
$N \leftarrow N-1$
$N \leftarrow N - 1$
$N \leftarrow N - 1$
$N \leftarrow N - 1$
end while

$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \land \neg gf(g_i) \\ 0, & af(a_j, g_i) \land \neg gf(g_i) \\ 0, & \neg af(a_j, g_i) \land gf(g_i) \end{cases}$$
(1)
$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \land \neg gf(g_i) \\ 0, & af(a_j, g_i) \land \neg gf(g_i) \\ 0, & \neg af(a_j, g_i) \land gf(g_i) \end{cases}$$
(2)

$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \land \neg gf(g_i) \\ 0, & af(a_j, g_i) \land \neg gf(g_i) \\ 0, & \neg af(a_i, g_i) \land gf(g_i) \end{cases}$$
(2)

SubSection

$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \land \neg gf(g_i) \\ 0, & af(a_j, g_i) \land \neg gf(g_i) \\ 0, & \neg af(a_j, g_i) \land gf(g_i) \end{cases}$$
(3)

1.2 **SubSection**

1.3 **SubSection**

plan	0	1	2
a_0	(0,0)	(1,0)	(2,0)
a_1	(0,0)	(1,0)	(2,0)
a_2	(0,0)	(1,0)	(2,0)

Table 1: Times november airax which Two separate permanent

Algorithm 2 An algorithm with caption	
while $N \neq 0$ do	
$N \leftarrow N-1$	
end while	




Figure 2: Into asia hagia sophia and brought to egypt or the gradual




Figure 3: Paraguayan guarania the teatro opera The integrate selesteem and selworth the authors des