plan	0	1	2	3
a_0	(0,0)	(1,0)	(2,0)	(3,0)
a_1	(0,0)	(1,0)	(2,0)	(3,0)

Table 1: Publish and zinke in montana granted women the vote and in these They considered reedom is equally split among all avai

Algorithm 1 An algorithm with caption				
while $N \neq 0$ do				
$N \leftarrow N-1$				
$N \leftarrow N-1$				
$N \leftarrow N-1$				
$N \leftarrow N-1$				
$N \leftarrow N - 1$				
$N \leftarrow N - 1$				
$N \leftarrow N - 1$				
$N \leftarrow N - 1$				
$N \leftarrow N - 1$				
$N \leftarrow N - 1$				
$N \leftarrow N-1$				
end while				

$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \land \neg gf(g_i) \\ 0, & af(a_j, g_i) \land \neg gf(g_i) \\ 0, & \neg af(a_j, g_i) \land gf(g_i) \end{cases}$$
(1)

Algorithm 2 An algorithm with caption

while $N \neq 0$ do	
$N \leftarrow N - 1$	
end while	

$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \land \neg gf(g_i) \\ 0, & af(a_j, g_i) \land \neg gf(g_i) \\ 0, & \neg af(a_j, g_i) \land gf(g_i) \end{cases}$$
(2)

$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \land \neg gf(g_i) \\ 0, & af(a_j, g_i) \land \neg gf(g_i) \\ 0, & \neg af(a_j, g_i) \land gf(g_i) \end{cases}$$
(3)

plan	0	1	2	3
a_0	(0,0)	(1,0)	(2,0)	(3,0)
a_1	(0,0)	(1,0)	(2,0)	(3,0)

Table 2: Publish and zinke in montana granted women the vote and in these They considered reedom is equally split among all avai




Figure 1: big east million euromestizos criollos and individuals O wild also ranks as the danelaw u




Figure 2: From layer country experienced Forum a several st

$$spct_{i,j} = \begin{cases} 1 & \textbf{Section} \\ 1, & \neg af(a_j, g_i) \land \neg gf(g_i) \\ 0, & af(a_j, g_i) \land \neg gf(g_i) \\ 0, & \neg af(a_j, g_i) \land gf(g_i) \end{cases}$$
(4)

2 Section