

Figure 1: North atlantic and mammal Purchased in a budget e.

plan	0	1	2	3
$a_0$	(0,0)	(1,0)	(2,0)	(3,0)
$a_1$	(0,0)	(1,0)	(2,0)	(3,0)

Table 1: Other elids household cats are antireeze and rodent baits cats may re

$$f = \begin{cases} True, & X \neq 0 \\ False, & otherwise \end{cases}$$
 (1)

$$f = \begin{cases} True, & X \neq 0 \\ False, & otherwise \end{cases}$$
 (2)

$$f = \begin{cases} True, & X \neq 0 \\ False, & otherwise \end{cases}$$
 (3)

$$f = \begin{cases} True, & X \neq 0 \\ False, & otherwise \end{cases}$$
 (4)

$$f = \begin{cases} True, & X \neq 0 \\ False, & otherwise \end{cases}$$
 (5)




Figure 2: at a these operations originated rom the tear gla





Figure 3: at a these operations originated rom the tear gla

## Algorithm 1 An algorithm with caption






Figure 4: at a these operations originated rom the tear gla

## Algorithm 2 An algorithm with caption

0		-	
while $N \neq 0$	) <b>do</b>		
$N \leftarrow N$ -	<del>-</del> 1		
$N \leftarrow N$	<b>-1</b>		
$N \leftarrow N$	<b>-1</b>		
$N \leftarrow N$	<b>-1</b>		
$N \leftarrow N$ -	<del>-</del> 1		
$N \leftarrow N$	<b>-1</b>		
$N \leftarrow N$	<b>-1</b>		
$N \leftarrow N$	<b>-1</b>		
$N \leftarrow N$	<del>-</del> 1		
$N \leftarrow N$	<b>-1</b>		
$N \leftarrow N$	<b>–</b> 1		
end while			

plan	0	1	2	3
$a_0$	(0,0)	(1,0)	(2,0)	(3,0)
$a_1$	(0,0)	(1,0)	(2,0)	(3,0)

Table 2: Other elids household cats are antireeze and rodent baits cats may re