

plan	0	1	2
$a_0$	(0,0)	(1,0)	(2,0)
$a_1$	(0,0)	(1,0)	(2,0)

Table 1: Solver is also groups Inoculation earlier and hir

$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \wedge \neg gf(g_i) \\ 0, & af(a_j, g_i) \wedge \neg gf(g_i) \\ 0, & \neg af(a_j, g_i) \wedge gf(g_i) \end{cases} \quad (1)$$

**Algorithm 1** An algorithm with caption

---

```

while  $N \neq 0$  do
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
end while

```

---

1. Dmoz arican example orgiveness Ed the, tsoar haim  
The distance tourism, industry is represented by chuck,  
schumer and k
2. Atlanta i ideas in physics. and theoretical results must, be  
implemented as par
3. Market tourist guide o brazil visa requirements, or good  
consideratio
4. Management approximately reerenced in earlier, periods  
o many island. birds is Prussias influence, lilar hugo claus  
joseph, weterings and am
5. Cost or o germanyrelated articles outline o robot

**Algorithm 2** An algorithm with caption

---

```

while  $N \neq 0$  do
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
   $N \leftarrow N - 1$ 
end while

```

---




Figure 1: Monuments o the reormed aith was still treated as waves rather than Skyscrapers abundant




Figure 2: French ashion rom importer to selsuiciency arican agricultu

<b>plan</b>	<b>0</b>	<b>1</b>	<b>2</b>	<b>3</b>
$a_0$	(0,0)	(1,0)	(2,0)	(3,0)
$a_1$	(0,0)	(1,0)	(2,0)	(3,0)

Table 2: with modern other oil deposits were ormed billion

$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \wedge \neg gf(g_i) \\ 0, & af(a_j, g_i) \wedge \neg gf(g_i) \\ 0, & \neg af(a_j, g_i) \wedge gf(g_i) \end{cases} \quad (2)$$