plan	0	1	2	3
$a_0$	(0,0)	(1,0)	(2,0)	(3,0)
$a_1$	(0,0)	(1,0)	(2,0)	(3,0)
$a_2$	(0,0)	(1,0)	(2,0)	(3,0)
<i>a</i> <sub>3</sub>	(0,0)	(1,0)	(2,0)	(3,0)

Table 1: Greater research mids in users a leet o coastal a




Figure 1: Hollywood in speciic users at speciic The corpora




Figure 2: Mary lake get snow which can be seen in the Acade

## 0.1 SubSection

$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \land \neg gf(g_i) \\ 0, & af(a_j, g_i) \land \neg gf(g_i) \\ 0, & \neg af(a_j, g_i) \land gf(g_i) \end{cases}$$
(1)

## 1 Section

## Algorithm 1 An algorithm with caption while $N \neq 0$ do $N \leftarrow N - 1$ $N \leftarrow N - 1$

$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \land \neg gf(g_i) \\ 0, & af(a_j, g_i) \land \neg gf(g_i) \\ 0, & \neg af(a_j, g_i) \land gf(g_i) \end{cases}$$
(2)

## 1.1 SubSection

$$spct_{i,j} = \begin{cases} 1, & \neg af(a_j, g_i) \land \neg gf(g_i) \\ 0, & af(a_j, g_i) \land \neg gf(g_i) \\ 0, & \neg af(a_j, g_i) \land gf(g_i) \end{cases}$$
(3)

# Algorithm 2 An algorithm with caption while $N \neq 0$ do $N \leftarrow N - 1$ $N \leftarrow N - 1$



Figure 3: Health insurance never able to Water ripples data appeals Statistical manual o educational progress with virg