

Case Studies: Dementia Care as a Stress Test

Case Studies: Dementia Care as a Stress Test I

We illustrate the framework with one worked case study in dementia care. It is presented not as evidence of efficacy but as a boundary test: a setting chosen because it strains several of DigiPPPiP's core assumptions rather than confirming them. Other applications (early-childhood dyads, long-distance couples, stroke rehabilitation) belong to the empirical program in [sec:agenda]; here we develop the single case that most sharply exposes what the kernel does and does not guarantee.

Dementia Care: Asymmetric Dyads, Biographical Scaffolding, and Fluctuating Consent I

Dementia Care: Asymmetric Dyads, Biographical Scaffolding, and Fluctuating Consent II

DigiPPP's minimum viable kernel (two partners, a shared mark field, perceptible traces of agency, and consensual control over persistence) maps onto dementia care dyads in ways that productively stress-test several of the framework's core assumptions. The most important asymmetry is cognitive: where the framework presupposes two agents with roughly stable capacity to govern the shared archive, dementia introduces radical temporal asymmetry. The adult child experiences each session as one point in a longitudinal caregiving relationship; the parent may experience each session as the first. This does not invalidate the practice.

There is substantial evidence that creative engagement can produce genuine moments of co-presence and recognition even when sustained narrative continuity is lost; however, it requires the consent and governance model to be substantially extended.

Archive control cannot rest symmetrically with both partners when one partner's capacity fluctuates within and across sessions. The framework's default rule (when partners disagree, the