

$$\mathbb{R}^n$$

$$\mathbb{R}^{n_n}$$

$$q=(a,b,c,d)$$

$$A,B,C,D$$

$$k$$

$$q'$$

$$P_0,\ldots,P_3$$

$$d_{ij}$$

$$(\cdot)$$

$$|\cdot|$$

$$V_{ivm}$$

$$V_{xyz}$$

$$S3$$

$$k=(a,b,c,d)q'=q-(k,k,k,k)$$

$$P_iP_j$$

$$V_{ivm}=1$$

$$S3=\sqrt{9/8}V_{ivm}=S3V_{xyz}$$

$$\begin{array}{l} \alpha\\ \gamma\\ \rho\\ \sigma\\ V_{ivm}\end{array}$$

$$\begin{array}{l} \alpha,\gamma,\rho,\sigma\\ V_{ivm}\end{array}$$

$$\begin{array}{l} \mathbb{E}[\cdot] \\ F_{ij} \\ \mathcal{F} \\ [Q\|P] \\ \nabla_{\theta}L \\ \eta \\ \theta \\ ds^2 \\ c\end{array}$$

$$e$$

$$\begin{array}{l} \mathbb{E}[\partial_{\theta_i} \; p \cdot \partial_{\theta_j} \; p] \\ -\,P(o \mid s) + [Q(s) \parallel P(s)] \\ \sum Q\left(Q/P\right) \\ F(\theta)^{-1}\nabla_{\theta}L(\theta)\end{array}$$

$$\begin{array}{l} \theta_i \\ -c^2\,dt^2+dx^2+dy^2+dz^2\end{array}$$

$$\begin{array}{l} M \\ \|\cdot\|_2 \\ R,D\end{array}$$

$$\begin{array}{l} \mathbb{R}^3 \\ \sqrt{x_1^2+\cdots+x_n^2} \\ RDD=2R\end{array}$$

$$\begin{array}{l} \alpha,\gamma,\rho,\sigma\\ \theta\\ \mu\\ \psi\\ \eta\end{array}$$