# Results: Field Overview

## Results: Field Overview I

The de-duplicated retrieval corpus for **Modafinil** contains $N = 2334$ records. The analyzed subset contains 2234 records; 2234 have publication years spanning 2000–2026 (26 years). Publication volume within this dated subset grows at a compound annual rate of 5.48% (mean year-over-year growth 7.8%, doubling time 9.2 years), peaking in 2025 with 147 records that year. The growth curve is a first-order descriptive summary of this retained analyzed subset. A field-wide estimate of research activity would require a separate live coverage audit.

# Results: Field Overview I




Figure 2: Publication growth curve for Modafinil. Annual publication counts (bars) and cumulative total (line) summarize the retained dated subset from 2000 through 2026, with its largest annual count in 2025.

## RQ1: Field Size and Growth I

The temporal analysis describes a retained analyzed literature slice spanning 26 years. The compound annual growth rate of 5.48% implies a corpus doubling time of approximately 9.2 years under the configured retrieval, filtering, and indexing conditions. The peak year 2025 contains 147 retained records. That count may reflect both publication activity and source-indexing delays; a causal trend would require a separate live coverage audit.

**Table 1. Top publication years.**

# RQ1: Field Size and Growth I

| Year | Publications |
|---|---|
| 2016 | 103 |
| 2017 | 106 |
| 2018 | 96 |
| 2019 | 107 |
| 2020 | 112 |
| 2021 | 103 |
| 2022 | 117 |
| 2023 | 106 |

# RQ1: Field Size and Growth I

| Year | Publications |
|---|---|
| 2024 | 123 |
| 2025 | 147 |

Records distribute across the 6 configured subfields as shown in Table 2, with **Clinical Sleep** the largest bucket at 63.0% of the classified corpus. The dominance of Clinical Sleep reflects the clinical primacy of modafinil as a wakefulness-promoting agent: the largest body of literature addresses its use in narcolepsy, shift-work disorder, and obstructive sleep apnea.

**Table 2. Subfield distribution.**

# RQ2: Subfield Composition I

| Subfield | Papers | Share |
|---|---|---|
| Clinical Sleep | 1407 | 63.0% |
| Cognition | 249 | 11.1% |
| Pharmacology | 76 | 3.4% |
| Psychiatry | 359 | 16.1% |
| Safety | 94 | 4.2% |
| Neuroscience | 49 | 2.2% |

# RQ2: Subfield Composition I




Figure 3: Field summary dashboard for Modafinil. The dashboard combines corpus size, temporal range, subfield distribution, and key bibliometric indicators in a single overview panel.

# RQ2: Subfield Composition I





Figure 4: Subfield distribution for Modafinil. The 6-bucket taxonomy shows the relative weight of each configured sub-area, with Clinical Sleep dominant at 63.0%.

# RQ2: Subfield Composition I



## RQ2: Subfield Composition I

Figure 5: Subfield timeline for Modafinil. Stacked annual publication counts by subfield show how each sub-area has evolved over time, revealing emerging and declining research threads.

## Identifier and Full-Text Coverage I

The retrieved corpus carries DOIs for 2260 of 2334 records, enabling robust cross-engine de-duplication. OpenAlex IDs are present for 923 records. Abstract coverage stands at 61.6% (1437 records), which limits the text analytics to that subset. Open access is confirmed for 24.6% of records, and 54.6% have a direct PDF link.

## Descriptive Bibliometrics I

The analyzed subset spans 8152 unique authors across 2234 papers, yielding a mean of 1.28 papers per author. Citation counts range from zero to 1353 (mean 30.3, median 0.0), with a total of 67,646 citations across the analyzed subset. The Gini coefficient of citation concentration is 0.817, indicating a highly skewed distribution characteristic of scientific literature.

# Descriptive Bibliometrics II

**Table 3. Citation count distribution.**

# Descriptive Bibliometrics I

| Citations | Papers |
|---|---|
| 0 | 1253 |
| 1-9 | 165 |
| 10-49 | 416 |
| 50-99 | 213 |
| 100-499 | 176 |
| 500+ | 11 |

# Descriptive Bibliometrics I




Figure 6: Citation distribution for Modafinil. The histogram shows the number of papers in each citation-count bucket, with the Gini coefficient annotated. The heavy-tailed distribution is characteristic of scientific citation networks.

# Descriptive Bibliometrics I

# Descriptive Bibliometrics II

**Table 4. Top publication venues.**

# Descriptive Bibliometrics I

| Venue | Papers |
|---|---|
| Reactions Weekly | 61 |
| Psychopharmacology | 41 |
| SLEEP | 34 |
| Sleep Medicine | 33 |
| The Journal of Clinical Psychiatry | 31 |
| European Neuropsychopharmacology | 26 |
| Neuropharmacology | 26 |
| Inpharma Weekly | 25 |

# Descriptive Bibliometrics I

| Venue | Papers |
|---|---|
| Sleep | 24 |
| American Journal of Psychiatry | 23 |

# Descriptive Bibliometrics I




Figure 7: Top publication venues for Modafinil. The horizontal bar chart shows the leading venues by paper count in the analyzed subset, revealing where the modafinil literature is published.

# Descriptive Bibliometrics I

# Descriptive Bibliometrics II

**Table 5. Top authors by publication count.**

# Descriptive Bibliometrics I

| Rank | Author | Papers |
|---|---|---|
| 1 | &NA; | 41 |
| 2 | Ronghua Yang | 25 |
| 3 | Yves Dauvilliers | 22 |
| 4 | Barbara J. Sahakian | 19 |
| 5 | Edward T. Hellriegel | 17 |
| 6 | Philmore Robertson | 16 |
| 7 | Mona Darwish | 15 |
| 8 | Sanjay Arora | 15 |

# Descriptive Bibliometrics I

| Rank | Author | Papers |
|---|---|---|
| 9 | Amy Hauck Newman | 14 |
| 10 | Jed Black | 14 |

# Descriptive Bibliometrics I




Figure 8: Author productivity for Modafinil. The horizontal bar chart shows the 20 authors with the most papers in the corpus, revealing the most prolific contributors to the modafinil literature.