

CENG 384 - Signals and Systems for Computer Engineers
Spring 2024
Homework 4

Boloğur, Baran
e2521292@ceng.metu.edu.tr

Doğın, Alkım
e2521482@ceng.metu.edu.tr

May 19, 2024

1. (a) The inverse of the integrator operator is the derivative operator. By using this fact, we can get the differential equation as the following.

$$\frac{d^2y(t)}{dt^2} + 5\frac{dy(t)}{dt} + 6y(t) = 4\frac{dx(t)}{dt} + x(t)$$

- (b) We know the following formula.

$$H(jw) = \frac{\sum^M b_k(jw)^k}{\sum^N a_k(jw)^k}$$

By applying this formula, we find $H(jw) = \frac{1+4jw}{(jw)^2+5jw+6}$. We can write this as the following.

$$\frac{1+4jw}{(jw)^2+5jw+6} = \frac{A}{jw+2} + \frac{B}{jw+3}$$

$$A+B=4$$

$$3A+2B=1$$

We get $A = -7$ and $B = 1$.

So,

$$H(jw) = \frac{1+4jw}{(jw)^2+5jw+6} = \frac{-7}{jw+2} + \frac{11}{jw+3}$$

is the answer.

- (c) Fourier transform of Impulse response is equal to frequency response. We have already found the frequency response. We can find impulse response from the look up table. From the look up table, we know that

$$e^{-at}u(t) \Leftrightarrow \frac{1}{a+jw}$$

We know $H(jw) = \frac{-7}{jw+2} + \frac{11}{jw+3}$. So, by using linearity property and the above equation, we find $h(t)$.

$$h(t) = \left[-7e^{-2t} + 11e^{-3t} \right] u(t)$$

- (d) For this part, we should make use of the formula

$$Y(jw) = H(jw)X(jw)$$

We already know the $H(jw)$. Lets find $X(jw)$. From the same formula in the look up table, we find

$$X(jw) = \frac{1}{4\frac{1}{4}+jw}$$

By applying $Y(jw) = H(jw)X(jw)$.

$$Y(jw) = \frac{1}{4} \left[\frac{1}{\frac{1}{4} + jw} \right] \left[\frac{-7}{jw + 2} + \frac{11}{jw + 3} \right]$$

$$Y(jw) = \frac{1}{4} \left[\frac{-7}{(\frac{1}{4} + jw)(jw + 2)} + \frac{11}{(\frac{1}{4} + jw)(jw + 3)} \right]$$

$$Y(jw) = \frac{1}{4} \left[\frac{A}{\frac{1}{4} + jw} + \frac{B}{jw + 2} + \frac{C}{\frac{1}{4} + jw} + \frac{D}{jw + 3} \right]$$

We find $A = -4, B = 4, C = 4, D = -4$. Thus, we get.

$$Y(jw) = \frac{1}{jw + 2} - \frac{1}{jw + 3}$$

2. (a)

$$H(jw) = \frac{jw + 4}{-w^2 + 5jw + 6} = \frac{\sum b_k(jw)^k}{\sum a_k(jw)^k}$$

$$b_0 = 4$$

$$b_1 = 1$$

$$a_0 = 6$$

$$a_1 = 5$$

$$a_2 = 1$$

So, the difference equation can be written as the following.

$$\frac{d^2y(t)}{dt^2} + 5\frac{dy(t)}{dt} + 6y(t) = \frac{dx(t)}{dt} + 4x(t)$$

(b)

$$H(jw) = \frac{jw + 4}{-w^2 + 5jw + 6} = \frac{jw + 4}{(jw)^2 + 5jw + 6} = \frac{A}{jw + 3} + \frac{B}{jw + 2}$$

$$A + B = 1$$

$$2A + 3B = 4$$

$$A = -1$$

$$B = 2$$

We can use the look up table to convert $H(jw)$ into the time domain.

We basically use the formula $e^{-at}u(t) \Leftrightarrow \frac{1}{a+jw}$

$$H(jw) = \frac{-1}{jw + 3} + \frac{2}{jw + 2} \Leftrightarrow (-e^{-3t} + 2e^{-2t})u(t) = h(t)$$

So, the result is

$$h(t) = (-e^{-3t} + 2e^{-2t})u(t)$$

(c)

$$x(t) \Leftrightarrow \frac{1}{jw + 4} - \frac{1}{(jw + 4)^2}$$

$Y(jw) = X(jw)H(jw)$ and we already know the formula for $H(jw)$.

$$Y(jw) = \left[\frac{1}{jw + 4} - \frac{1}{(jw + 4)^2} \right] \left[\frac{jw + 4}{(jw + 3)(jw + 2)} \right]$$

$$Y(jw) = \frac{jw + 3}{(jw + 4)^2} + \frac{jw + 4}{(jw + 3)(jw + 2)}$$

$$Y(jw) = \frac{1}{(jw + 4)(jw + 2)} = \frac{A}{jw + 4} + \frac{B}{jw + 2}$$

$$\begin{aligned}
A + B &= 0 \\
2A + 4B &= 1 \\
A &= -\frac{1}{2} \\
B &= \frac{1}{2}
\end{aligned}$$

So, we get

$$Y(jw) = \frac{1}{2} \left[\frac{-1}{jw + 4} + \frac{1}{jw + 2} \right]$$

(d) Once again, we make use of the formula from the look up table. $e^{-at}u(t) \Leftrightarrow \frac{1}{a+jw}$. We find the answer as

$$y(t) = \frac{1}{2} \left[-e^{-4t} + e^{-2t} \right] u(t)$$

3. (a)

From lookup table;

$$\begin{aligned}
1) \quad a^n u[n] &\Leftrightarrow \frac{1}{1 - ae^{-jw}} \\
2) \quad (n+1)a^n u[n] &\Leftrightarrow \frac{1}{(1 - ae^{-jw})^2}
\end{aligned}$$

$$\text{Using property 1, } x[n] = \left(\frac{2}{3}\right)^n u[n] \Leftrightarrow \frac{1}{1 - \frac{2}{3}e^{-jw}} = X(e^{jw})$$

$$\text{Using property 2, } x[n] + \frac{3}{2}y[n] = (n+1)\left(\frac{2}{3}\right)^n u[n] \Leftrightarrow \frac{1}{(1 - \frac{2}{3}e^{-jw})^2} = X(e^{jw}) + \frac{3}{2}Y(e^{jw})$$

$$\text{Hence } Y(e^{jw}) = \frac{2}{3} \left[\frac{1}{(1 - \frac{2}{3}e^{-jw})^2} - \frac{1}{1 - \frac{2}{3}e^{-jw}} \right]$$

$$H(e^{jw}) = \frac{Y(e^{jw})}{X(e^{jw})} = \frac{2 \left[\frac{1}{(1 - \frac{2}{3}e^{-jw})^2} - \frac{1}{1 - \frac{2}{3}e^{-jw}} \right]}{\frac{1}{1 - \frac{2}{3}e^{-jw}}}$$

$$\text{After simplification, } H(e^{jw}) = \frac{2}{3} \left[\frac{1}{1 - \frac{2}{3}e^{-jw}} - 1 \right]$$

(b)

From lookup table;

$$h[n] = \frac{2}{3} \left[\left(\frac{2}{3}\right)^n u[n] - \delta[n] \right]$$

(c)

$$\text{From part (a), } H(e^{jw}) = \frac{2}{3} \left[\frac{1}{1 - \frac{2}{3}e^{-jw}} - 1 \right] = \frac{2}{3} \left[\frac{1 - 1 + \frac{2}{3}e^{-jw}}{1 - \frac{2}{3}e^{-jw}} \right] = \frac{2}{3} \left[\frac{\frac{2}{3}e^{-jw}}{1 - \frac{2}{3}e^{-jw}} \right]$$

$$H(e^{jw}) = \frac{\frac{4}{3}e^{-jw}}{3 - 2e^{-jw}}$$

$$\text{We also know, } H(e^{jw}) = \frac{\sum b_k e^{-jwk}}{\sum a_k e^{-jwk}}$$

$$b_1 = \frac{4}{3}, \quad a_0 = 3, \quad a_1 = -2, \quad \text{and all other coefficients are equal to 0.}$$

$$\text{The final result is, } 3y[n] - 2y[n-1] = \frac{4}{3}x[n-1]$$

(d)

Block diagram of the equation $3y[n] - 2y[n-1] = \frac{4}{3}x[n-1]$

$$y[n] = \frac{4}{9}x[n-1] + \frac{2}{3}y[n-1]$$




Figure 1: 3 - d Block Diagram

4. (a)

The difference equation that represents this system is:

$$2x[n] - \frac{1}{8}y[n-2] + \frac{3}{4}y[n-1] = y[n]$$

or

$$y[n] - \frac{3}{4}y[n-1] + \frac{1}{8}y[n-2] = 2x[n]$$

(b)

$$Y(e^{jw}) = H(e^{jw})X(e^{jw})$$

$$H(e^{jw})X(e^{jw}) - \frac{3}{4}e^{-jw}H(e^{jw})X(e^{jw}) + \frac{1}{8}e^{-2jw}H(e^{jw})X(e^{jw}) = 2X(e^{jw})$$

$$H(e^{jw}) \left[1 - \frac{3}{4}e^{-jw} + \frac{1}{8}e^{-2jw} \right] = 2 \quad \longrightarrow \quad H(e^{jw}) = \frac{2}{1 - \frac{3}{4}e^{-jw} + \frac{1}{8}e^{-2jw}}$$

$$H(e^{jw}) = \frac{16}{8 - 6e^{-jw} + e^{-2jw}}$$

(c)

$$H(e^{jw}) = \frac{16}{8 - 6e^{-jw} + e^{-2jw}} = \frac{A}{e^{-jw} - 4} + \frac{B}{e^{-jw} - 2}$$

$$A + B = 0 \quad \text{and} \quad -2A - 4B = 16 \quad \longrightarrow \quad A = 8 \quad \text{and} \quad B = -8$$

$$H(e^{jw}) = \frac{8}{-4(1 - \frac{1}{4}e^{-jw})} + \frac{-8}{-2(1 - \frac{1}{2}e^{-jw})} = \frac{-2}{1 - \frac{1}{4}e^{-jw}} + \frac{4}{1 - \frac{1}{2}e^{-jw}}$$

From lookup table;

$$h[n] = \left(-2 \left(\frac{1}{4} \right)^n + 4 \left(\frac{1}{2} \right)^n \right) u[n]$$

(d)

$$x[n] = \left(\frac{1}{4}\right)^n u[n] \Leftrightarrow \frac{1}{1 - \frac{1}{4}e^{-jw}} = X(e^{jw})$$

$$Y(e^{jw}) = H(e^{jw})X(e^{jw})$$

$$Y(e^{jw}) = \frac{-2}{(1 - \frac{1}{4}e^{-jw})^2} + \frac{4}{(1 - \frac{1}{2}e^{-jw})(1 - \frac{1}{4}e^{-jw})}$$

$$Y(e^{jw}) = \frac{-2}{(1 - \frac{1}{4}e^{-jw})^2} + \frac{A}{(1 - \frac{1}{2}e^{-jw})} + \frac{B}{(1 - \frac{1}{4}e^{-jw})}$$

After the calculations, we find that: $A = 8, B = -4$

$$Y(e^{jw}) = \frac{-2}{(1 - \frac{1}{4}e^{-jw})^2} + \frac{8}{(1 - \frac{1}{2}e^{-jw})} + \frac{-4}{(1 - \frac{1}{4}e^{-jw})}$$

From lookup table;

$$y(t) = \left[-2(n+1) \left(\frac{1}{4}\right)^n + 8 \left(\frac{1}{2}\right)^n - 4 \left(\frac{1}{4}\right)^n \right] u[n]$$

5.

$$Y(e^{jw}) = H_1(e^{jw})X(e^{jw}) + H_2(e^{jw})X(e^{jw})$$

$$Y(e^{jw}) = X(e^{jw}) [H_1(e^{jw}) + H_2(e^{jw})]$$

$$\text{Using Lookup Table; } H_1(e^{jw}) = \frac{1}{1 - \frac{1}{3}e^{-jw}}$$

$$\text{Also; } \frac{5e^{-jw} - 12}{e^{-2jw} - 7e^{-jw} + 12} = \frac{A}{e^{-jw} - 4} + \frac{B}{e^{-jw} - 3}$$

After the calculations, we find that: $A = 8, B = -3$

$$\text{We also know: } H_1(e^{jw}) + H_2(e^{jw}) = H(e^{jw})$$

$$\frac{1}{1 - \frac{1}{3}e^{-jw}} + H_2(e^{jw}) = \frac{8}{e^{-jw} - 4} + \frac{-3}{e^{-jw} - 3} == \frac{-2}{1 - \frac{1}{4}e^{-jw}} + \frac{1}{1 - \frac{1}{3}e^{-jw}}$$

$$H_2(e^{jw}) = \frac{-2}{1 - \frac{1}{4}e^{-jw}}$$

From lookup table;

$$h_2[n] = -2 \left(\frac{1}{4}\right)^n u[n]$$

```

6.
1 import numpy as np
2 import matplotlib.pyplot as plt
3
4 N = 10**3
5 def x(n): # defining x[n]
6     if n < 0:
7         return (1/2)**(-n)
8     return 1/2**(n)
9
10 w_axis = np.linspace(-20, +20, 600)
11 def X(w): # defining x[e^(jw)]
12     return np.sum([x(n) * np.exp(1j * w * -n) for n in range(-N,N+1)])
13
14 result = [X(w) for w in w_axis]
15
16 magnitudes = []
17 phases = []
18
19 for i in result:
20     re = i.real
21     im = i.imag
22     magnitude = np.sqrt(re**2 + im**2)
23     magnitudes.append(magnitude)
24     # print(f"i:{i} - real:{re} - imaginary:{im} - magnitude is:{magnitude}") // debugging
25     if re == 0:
26         if im == 0:
27             phases.append(0)
28         elif im > 0:
29             phases.append(np.pi / 2)
30         else:
31             phases.append(-np.pi / 2)
32     else:
33         phase = np.arctan(im/re)
34         phases.append(round(phase,3))
35
36 fig = plt.figure(figsize=(12, 5))
37
38 w1 = w_axis
39 w2 = w_axis
40
41 plt.subplot(1, 2, 1)
42 plt.plot(w1, magnitudes, color='r', label='Magnitude')
43 plt.xlabel('w-axis')
44 plt.ylabel('X(e^(jw))-axis')
45 plt.title('Magnitude')
46 plt.grid(True)
47 plt.legend()
48
49 plt.subplot(1, 2, 2)
50 plt.plot(w2, phases, color='b', linestyle='--', label='Phase')
51 plt.xlabel('w-axis')
52 plt.ylabel('X(e^(jw))-axis')
53 plt.title('Phase')
54 plt.grid(True)
55 plt.legend()
56
57 plt.tight_layout()
58 plt.show()

```




Figure 2: Plotting of the given signals

From the figure, we can observe that the phase of the Fourier transform of the given signal is always zero. This means the corresponding Fourier Transform is always a real number.