Uniform families in the shape of the D = 8���10 Haar records

2026-10-27 (draft)

Uniform families in the shape of the D = 8–10 Haar records: the LPS family, a 4/3 obstruction, and what a cyclic factor does

Doğa Özden's degree–diameter lab, with Claude Code (job `thm-haar-families`, 2026-10-27). Written for Gabriela Araujo-Pardo and Ignacio López. Draft; corrections welcome.

Labels. Every statement carries one of three labels.

- **Theorem, Proposition, Lemma, Corollary**: complete proof in this note, or a cited result named in place (marked cited, with the source and how we know it: **[read]** primary source read, **[abstract]** abstract only, **[secondary]** known from a book, survey or citation).
- **Computer-checked**: a statement about finitely many explicit graphs, verified by programs that are named, with logs.
- **Heuristic**: a model, a measurement or an interpretation. Not used in any proof.

The question. The lab's records at D = 8, 9, 10 (`research/theorems/HAAR_D8_10.md` on branch `claude/thm-haar-d8-10-sf7f0a`, cited below as [H]) are Haar graphs of SL(2, q) × Z_m or PSL(2, q) × Z_m whose connection set S is closed under an automorphism of order 3, each found by computer search for one group. Can they be turned into infinite families, with S given by one formula in q (and m) and a proved diameter?

The answer, stated plainly.

1. There is an explicit infinite family of exactly this shape, with m = 1: the connection sets of Lubotzky, Phillips and Sarnak (LPS). For every odd prime p and every prime q > p, the p + 1 LPS matrices in PGL(2, q) are permuted by conjugation with an element of order 3 of PSL(2, q), with cycle type 1+1+3+⋯+3 or 3+⋯+3: the cycle types that the orbit search used at Δ = 6, 8, 12, 14 (Theorem 2.6). For p ≡ 1 (mod 4) their diameter is at most 2 log_p n + O(1) (cited, LPS).
2. This family provably cannot reach the records. The vertex ψ(i) is at distance exactly the least even L with $2p^{L/2} > q^2$ (Theorem 3.2: lower bound for every odd p, equality for p ≡ 1 mod 4), so the diameter is at least (4/3) log_p n − 2 log_p 2 (Corollary 3.3, the bound of N. T. Sardari, here with a two-paragraph proof), and a member of diameter ≤ D has at most $\sqrt{2}\cdot p^{1-D/4}$ of the bipartite Moore bound (Corollary 3.4). Computed: the LPS graphs are 2.5 to 19 times below the 2013 table at D = 8–10 (Appendix B), and at odd D the largest member is usually exactly the one that the vertex ψ(i) allows (8 of 13 cells; the next prime below in 3 more).
3. A cyclic factor removes this obstruction in practice. The obstruction survives every cover (Proposition 4.1), but it bounds q, not m. LPS × Z_m with labels chosen for each (q, m) reaches 99.4 % of the table at (6,10) and 98.9 % at (8,8) (Computer-checked 4.2), and with labels given by one formula in m (§4.3) the family has members **above the 2013 table** at (8,8) (210,528 vertices against 201,480; q = 17, m = 43) and (8,9) (1,105,104 against 1,091,232; q = 23, m = 91), each accepted by four or five independent programs, including the lab's record checkers. They are below the lab's records there, and their diameters are computed, not proved.
4. Other uniform formulas behave like one random draw. Integer lifts, primitive-root and Dickson formulas, the conic model, torus orbits, and the records' own sets moved to other q (18 squad cards, §5) give an excess diam − $log_{\Delta-1}|G|$ no better than a single random draw at the same Δ (about 2.8–3.4 at Δ = 6–8;

some kinds are worse), against about 2.0–2.3 for the records. Heuristically, the records' advantage is selection among millions of sets, which no single formula provides (§5–6).

5. What would be needed is open. A uniform family of record quality at fixed Δ would be an explicit family of bounded-degree graphs with diameter (1 + o(1)) $log_{\Delta-1}$ n. None is known; for the LPS graphs Sardari proved that it fails (§6).

No infinite family above the table is proved in this note; the members above the table in item 3 are single computed graphs.

Contents

0. Summary of statements
1. Setting, notation and projective coordinates
2. The LPS family has the records' shape
3. The 4/3 obstruction
4. Cyclic factors
5. Other uniform families (squad round 1)
6. Why no record-quality uniform family is in sight
7. Prior art
8. How to re-check

Appendix A. The squad rounds. Appendix B. The LPS family cell by cell.

---

0. Summary of statements

#	statement	label
Prop. 1.1	at fixed (Δ, D) only finitely many groups; for Q × Z_m with Q perfect, m ≤ A(Δ, D)	Proposition (from [H])
Lemma 2.3	ψ: H(Z) → $M_2(F_q)$ is a ring homomorphism with kernel qH(Z); det ψ(α) = N(α); ψ(ᾱ) = adj ψ(α)	Lemma
Lemma 2.4	for q > p the p + 1 matrices ψ(α), α ∈ S_p, are distinct in PGL(2, q), and ψ(S_p) is closed under inversion	Lemma
Lemma 2.5	structure of the component $\Gamma_{p,q}$; diam $\Gamma_{p,q}$ = ecc(I, A)	Lemma
Thm. 2.6	conjugation by h = ψ(1+i+j+k), of order 3 in PSL(2, q), permutes ψ(S_p) with cycle type 1+1+$3^{(p-1)/3}$ (p ≡ 1 mod 3), $3^{(p+1)/3}$ (p ≡ 2 mod 3), 1+3 (p = 3)	Theorem
Thm. 2.7	$\Gamma_{p,q}$ is connected and Ramanujan; diam ≤ 2 log_p n + O(1)	cited (LPS)
Lemma 3.1	an even walk from (I, A) ends at ψ(β) with N(β) = p^L and β ≡ 1 mod 2H(Z)	Lemma
Thm. 3.2	d((I,A), (ψ(i),A)) ≥ L_0 := min{L even : $2p^{L/2}$ > q^2}, with equality if p ≡ 1 mod 4	Theorem (equality uses a cited factorization)
Cor. 3.3	diam $\Gamma_{p,q}$ > (4/3) log_p n − 2 log_p 2	Corollary

#	statement	label
Cor. 3.4	if diam $\Gamma_{p,q} \le D$ then $n < 2^{3/2} p^{3D/4}$ and $n < \sqrt{2} \cdot p^{1-D/4} \cdot M^b(p+1, D)$	Corollary
3.5	equality in Thm. 3.2 in all 251 cases $p \in \{3, 5, 7, 11, 13, 17, 19, 29\}$, $p < q \le 160$ (including $p \equiv 3 \bmod 4$), and in all 50 cases $p \in \{5, 13\}$, $161 \le q \le 300$; ecc − $L_0 \in \{0, 1, 2, 3\}$	Computer-checked
App. B	the LPS family cell by cell; at odd D its largest member is the one Thm. 3.2 allows in 8 of 13 cells	Computer-checked
Prop. 3.6	the same lower bound in any definite quaternion order with an orthogonal unit γ	Proposition
Prop. 4.1	the bound of Thm. 3.2 holds in every cover $H(PGL(2,q) \times K, \tilde{S})$ of the LPS graph	Proposition
4.2	LPS × Z_m with searched labels: 99.4 % of the table at (6,10), 98.9 % at (8,8), 65 % at (14,8)	Computer-checked
4.3	LPS(7) × Z_m with labels by a formula: members above the 2013 table at (8,8) and (8,9) (below the lab's records)	Computer-checked
§5	uniform formulas of six kinds: excess about that of one random draw	Computer-checked (scores), Heuristic (reading)

---

1. Setting, notation and projective coordinates

1.1 Haar graphs

We use the conventions of [H] §1–2. For a finite group G and $S \subseteq G$ with $|S| = \Delta$, the Haar graph H(G, S) has vertex set $G \times \{A, B\}$, and (g, A) is adjacent to (sg, B) for every $s \in S$. It is bipartite and Δ-regular, the right translations $\rho_h(g, X) = (gh, X)$ are automorphisms acting regularly on each side ([H] Lemma 1.2), and its diameter is max(ecc(e, A), ecc(e, B)) ([H] Cor. 2.3). A walk of length L from (e, A) uses letters $t_1, \ldots, t_L \in S$ and ends at (w, A) or (w, B) with

$$w = \cdots t_4^{-1}\, t_3\, t_2^{-1}\, t_1 \qquad (\text{an } A\text{-word}),$$

and the walk backtracks exactly when two consecutive letters are equal ([H] Prop. 2.2). Lemma 3.1 of [H]: H(G, S) is isomorphic to $H(G, a\psi(S)^{\pm 1}b)$ for $a, b \in G$ and $\psi \in \mathrm{Aut}(G)$; the map $(g, A) \mapsto (g, B)$, $(h, B) \mapsto (h, A)$ is an isomorphism $H(G, S) \to H(G, S^{-1})$.

Throughout, $M^b(\Delta, D) = 2 \sum_{j=0}^{D-1} (\Delta - 1)^j$ is the bipartite Moore bound, and n denotes the number of vertices of the graph at hand.

1.2 Projective coordinates

Let q be an odd prime. An element of PGL(2, q) is a nonsingular 2 × 2 matrix up to a nonzero scalar, that is, a point

$$[X] = [a : b : c : d] \in \mathrm{PG}(3, q), \qquad X = \begin{pmatrix} a & b \\ c & d \end{pmatrix},$$

not on the quadric $\mathcal{Q}$: ad − bc = 0. The quadric $\mathcal{Q}$ is a hyperbolic quadric Q⁺(3, q) with (q + 1)² points, so PGL(2, q) is the set of the (q + 1)(q² + 1) − (q + 1)² = q(q² − 1) points of PG(3, q) off $\mathcal{Q}$. Whether ad − bc is a square does not depend on the representative, and PSL(2, q) is the set of the q(q² − 1)/2 points with ad − bc a nonzero square. For M ∈ GL(2, q), left multiplication [X] ↦ [MX] is a collineation of PG(3, q) that fixes $\mathcal{Q}$ (det MX = det M det X).

So a Haar graph H(PGL(2, q), S) is the bipartite graph on two copies of PG(3, q) $\setminus$ $\mathcal{Q}$ in which **the point [X] of the first copy is joined to the points [MX], M ∈ S, of the second copy**; and the point [Y] of the second copy is joined to the points [adj(M)·Y], M ∈ S, of the first, where adj [[a, b], [c, d]] = [[d, −b], [−c, a]] is the inverse up to a scalar. These are the exact neighbourhoods of every vertex, once S is written down (§2, Example 2.8).

1.3 The records' shape

[H] Construction 3.2: S is closed under an automorphism α of order k, here k = 3, and splits into f fixed letters and 3-cycles; for G = Q × Z_m the automorphism is α(x, z) = (gxg⁻¹, rz) with g of order 3 modulo the centre of Q and r of multiplicative order 3 mod m. The cycle types of the records are in [H] §5.1: 3+3 at (6,9), (6,10); 1+3+3 at Δ = 7; 1+1+3+3 at (8,8), (8,9), (8,10); 3+3+3 at (9,8), (9,9), (9,10); 3+3+3+3 at (12,9), (12,10); 1+1+3+3+3+3 at (14,9), (14,10); and so on. Since |S| = f + 3c, f ≡ Δ (mod 3); every record with k = 3 uses the least such f ∈ {0, 1, 2}.

By a **uniform family** we mean, informally, one rule that produces S from q (and m) without a search for each q: a formula, as for the Singer difference sets at D = 3. All the families below are uniform in this sense, except where a search over labels is stated.

1.4 Where a uniform family can live

Proposition 1.1. (a) For fixed Δ ≥ 2 and D, only finitely many groups G admit S with |S| = Δ and diam H(G, S) ≤ D. (b) If G = Q × Z_m with Q perfect (for example Q = SL(2, q) or PSL(2, q), q ≥ 4) and diam H(G, S) ≤ D, then m ≤ A(Δ, D) = Θ(D$^{\Delta-1}$) for fixed Δ.

Proof. (a) |G| ≤ M^b(Δ, D)/2 ([H] Thm. 4.1). (b) [H] Cor. 4.4(ii). ■

So an infinite uniform family of the records' shape has Δ fixed and D → ∞ (and then q → ∞, since m grows at most polynomially in D while |G| must grow exponentially), or D fixed and Δ → ∞. The LPS family below can be read in both regimes (Corollary 3.4).

---

2. The LPS family has the records' shape

2.1 Definitions

Let H(Z) = {a₀ + a₁i + a₂j + a₃k : a_t ∈ Z} be the Lipschitz quaternions (i² = j² = k² = ijk = −1), with conjugate $\bar{\alpha}$ = a₀ − a₁i − a₂j − a₃k and norm N(α) = α$\bar{\alpha}$ = a₀² + a₁² + a₂² + a₃². N is multiplicative.

Definition 2.1 (the LPS set). For an odd prime p let S_p be the set of α ∈ H(Z) with N(α) = p and

- if p ≡ 1 (mod 4): a₀ > 0 odd, and a₁, a₂, a₃ even;
- if p ≡ 3 (mod 4): a₀ even, a₁, a₂, a₃ odd, and either a₀ > 0, or a₀ = 0 and a₁ > 0.

By Jacobi's four-square theorem |S_p| = p + 1 (cited: LPS, Combinatorica 8 (1988), §2, and G. Davidoff, P. Sarnak, A. Valette, Elementary Number Theory, Group Theory and Ramanujan Graphs, LMS Student Texts 55 (2003), Ch. 2 [secondary]). If α ∈ S_p then $\bar{\alpha}$ or −$\bar{\alpha}$ is in S_p.

Definition 2.2 (the splitting). Let q be an odd prime, q ≠ p. Choose x, y ∈ F_q with x² + y² + 1 = 0 (they exist: the sets {x²} and {−1 − y²} both have (q + 1)/2 elements, so they meet). Put

$$\psi(a_0 + a_1 i + a_2 j + a_3 k) = \begin{pmatrix} a_0 + a_1 x + a_3 y & -a_1 y + a_2 + a_3 x \\ -a_1 y - a_2 + a_3 x & a_0 - a_1 x - a_3 y \end{pmatrix} \bmod q.$$

Lemma 2.3. ψ: H(Z) → $M_2(F_q)$ is a ring homomorphism with kernel qH(Z), and it is onto. det ψ(α) ≡ N(α), and ψ(α̃) = adj ψ(α), so ψ(α)ψ(α̃) = N(α)·I.

Proof. ψ is additive and ψ(1) = I. Put M_i = ψ(i) = [[x, −y], [−y, −x]], M_j = ψ(j) = [[0, 1], [−1, 0]], M_k = ψ(k) = [[y, x], [x, −y]]. Direct multiplication gives $M_i{}^2$ = (x² + y²)·I = −I, $M_j{}^2$ = −I, $M_i \cdot M_j = M_k$ and $M_j \cdot M_i$ = [[−y, −x], [−x, y]] = $-M_k$. Since H(Z) is the ring generated over Z by i, j subject to i² = j² = −1 and ji = −ij (with k = ij), the assignment i ↦ M_i, j ↦ M_j extends to a ring homomorphism, and it sends k to $M_i \cdot M_j = M_k$; this is the displayed formula. Its kernel contains qH(Z). If ψ(α) = 0, the diagonal gives a₀ + a₁x + a₃y = a₀ − a₁x − a₃y = 0, so a₀ = 0 and a₁x + a₃y = 0; the off-diagonal gives a₂ = 0 and a₃x − a₁y = 0. Then a₁(x² + y²) = x(a₁x + a₃y) − y(a₃x − a₁y) = 0, so a₁ = 0 (x² + y² = −1), and then a₃x = a₃y = 0 with (x, y) ≠ (0, 0), so a₃ = 0, all mod q. So the kernel is qH(Z), and the induced map H(Z)/qH(Z) → $M_2(F_q)$ is injective between sets of size q⁴, hence onto. The determinant of the displayed matrix is (a₀ + a₁x + a₃y)(a₀ − a₁x − a₃y) − (a₂ + a₃x − a₁y)(−a₂ + a₃x − a₁y) = a₀² − (a₁x + a₃y)² + a₂² − (a₃x − a₁y)² = a₀² + a₂² − (a₁² + a₃²)(x² + y²) = N(α). Finally adj ψ(α) has diagonal (a₀ − a₁x − a₃y, a₀ + a₁x + a₃y) and off-diagonal entries −(−a₁y + a₂ + a₃x), −(−a₁y − a₂ + a₃x), which is ψ(a₀ − a₁i − a₂j − a₃k). ■

Lemma 2.4. Let q > p. (a) If α, β ∈ H(Z) have N(α) = N(β) = p and ψ(α) = cψ(β) for some c ∈ F_q^×, then α = ±β. (b) The p + 1 points [ψ(α)], α ∈ S_p, of PGL(2, q) are distinct, and the set they form is closed under inversion.

Proof. (a) γ = αβ has N(γ) = p², and ψ(γ) = ψ(α)ψ(β) = cψ(β)ψ(β) = cp·I. By Lemma 2.3, γ − c̃p ∈ qH(Z) for an integer c̃ ≡ c, so the coordinates γ₁, γ₂, γ₃ are divisible by q. They are at most √N(γ) = p < q in absolute value, so they vanish: γ is an integer with γ² = p², γ = ±p. Then α = γβ⁻¹ = ±pβ/N(β) = ±β. (b) Distinct elements of S_p are not negatives of each other (the sign conditions of Definition 2.1 fix the sign), so (a) gives distinctness. By Lemma 2.3, [ψ(α)]⁻¹ = [adj ψ(α)] = [ψ(α̃)] = [ψ(±α̃)], and ±α̃ ∈ S_p. ■

Definition 2.5′. $\Gamma_{p,q}$ is the connected component of (I, A) in H(PGL(2, q), ψ(S_p)).

Lemma 2.5 (structure of $\Gamma_{p,q}$). Let q > p and let c ∈ ψ(S_p).

(a) All of ψ(S_p) lies in one coset of PSL(2, q): in PSL(2, q) if p is a square mod q, in the other coset if not.
(b) The A-vertices of $\Gamma_{p,q}$ form H₀ × {A} for a subgroup H₀ ≤ PSL(2, q), and its B-vertices cH₀ × {B}. The right translations ρ_h, h ∈ H₀, map $\Gamma_{p,q}$ to itself and act regularly on each side. When H₀ = PSL(2, q), $\Gamma_{p,q}$ has n = q(q² − 1) vertices.
(c) diam $\Gamma_{p,q}$ = ecc(I, A) (computed in $\Gamma_{p,q}$).
(d) $\Gamma_{p,q} \cong$ H(PSL(2, q), c⁻¹ψ(S_p)) when it has q(q² − 1) vertices.

Proof. (a) det ψ(α) = p for all α (Lemma 2.3), and PSL(2, q) is the kernel of the homomorphism PGL(2, q) → $F_q{}^{\times}/(F_q{}^{\times})^2$ induced by the determinant. (b) A letter and its inverse both have determinant class [p], so an A-word of length L has class $[p]^L$: it lies in PSL(2, q) for L even and in cPSL(2, q) for L odd. The set H₀ of the values of even A-words (the A-side of $\Gamma_{p,q}$) is closed under products and contains I, so it is a subgroup of the finite group PSL(2, q). Right translation ρ_h by h ∈ H₀ is an automorphism ([H] Lemma 1.2) mapping (I, A) to (h, A) ∈ $\Gamma_{p,q}$, so it maps $\Gamma_{p,q}$ to itself; $\{\rho_h : h \in H_0\}$ acts regularly on the A-side H₀ × {A} and on the B-side cH₀ × {B}. When Theorem 2.7 applies, H₀ = PSL(2, q). (c) By (b) and [H] Lemma 2.1 (with the group $\{\rho_h : h \in H_0\}$), diam $\Gamma_{p,q}$ = max(ecc(I, A), ecc(c, B)). Since ψ(S_p) is closed under inversion (Lemma 2.4), the side swap σ(g, A) = (g, B), σ(g, B) = (g, A) is an automorphism of H(PGL(2, q), ψ(S_p)) ([H] Lemma 3.1(iv)); and ρ_c is one too. $\rho_c \circ \sigma$ maps (I, A) to (c, B) and maps $\Gamma_{p,q}$ onto the component of (c, B), which is $\Gamma_{p,q}$ since (c, B) ~ (I, A). Hence ecc(c, B) = ecc(I, A). (d) The map (g, A) ↦ (g, A), (h, B) ↦ (c⁻¹h, B) is the isomorphism H(G, S) → H(G, c⁻¹S) of [H] Lemma 3.1(i); restricted to $\Gamma_{p,q}$ it lands in PSL(2, q) × {A, B}, and both graphs are Δ-regular with q(q² − 1) vertices. ■

2.2 The order-3 symmetry

Theorem 2.6. Let p, q be odd primes, q > p, let ζ = 1 + i + j + k and h = ψ(ζ).

(a) h ∈ GL(2, q) has determinant 4, so [h] ∈ PSL(2, q), and [h] has order 3 in PGL(2, q).

(b) For every $\alpha \in H(Z)$, $h\psi(\alpha)h^{-1} = \psi(\sigma(\alpha))$, where $\sigma(a_0 + a_1 i + a_2 j + a_3 k) = a_0 + a_3 i + a_1 j + a_2 k$.

(c) Conjugation by h permutes the $p + 1$ points of $\psi(S_p)$. Its fixed points are the $[\psi(\alpha)]$ with $a_1 = a_2 = a_3$, and its cycle type is

p	cycle type on $\psi(S_p)$	$\Delta = p + 1$	records with this cycle type ([H] §5.1)
$p = 3$	$1 + 3$	4	none ($\Delta = 4$ records are metacyclic)
$p \equiv 1 \pmod 3$	$1 + 1 + 3^{(p-1)/3}$	8 ($p = 7$), 14 ($p = 13$), 20, 32, 38, …	(8,8), (8,9), (8,10); (14,9), (14,10)
$p \equiv 2 \pmod 3$	$3^{(p+1)/3}$	6 ($p = 5$), 12 ($p = 11$), 18, 24, 30, …	(6,9), (6,10); (12,9), (12,10)

So the LPS Haar graph is an orbit-closed Haar graph in the sense of [H] Construction 3.2, with α = conjugation by an element of order 3 of PSL(2, q) and $m = 1$, and with the same cycle type as the records at $\Delta = 6$, 8, 12 and 14.

Proof. (a) $N(\zeta) = 4$, so $\det h = 4 = 2^2$, a nonzero square (q odd). Write $v = i + j + k$, so $v^2 = -3$ and $\zeta = 1 + v$; then $\zeta^2 = 1 + 2v + v^2 = -2 + 2v$ and $\zeta^3 = (-2 + 2v)(1 + v) = -2 - 2v + 2v + 2v^2 = -8$. So $h^3 = -8I$ is a scalar. And h is not a scalar: $\psi(\zeta) - \tilde{c}I = \psi(\zeta - \tilde{c})$ would vanish only if $\zeta - \tilde{c} \in qH(Z)$ (Lemma 2.3), but the i-coordinate of $\zeta - \tilde{c}$ is 1. So [h] has order 3.

(b) We compute $\zeta i \zeta$. First $\zeta i = i + i^2 + ji + ki = -1 + i + j - k$. Then $(-1 + i + j - k)(1 - i - j - k) = (-1 + i + j + k) + (i + 1 + j - k) + (j + k + 1 - i) + (-k + j - i - 1) = 4j$, using $ij = k$, $ik = -j$, $ji = -k$, $jk = i$, $ki = j$, $kj = -i$. So $\zeta i \zeta = 4j$, that is, $\zeta i \zeta^{-1} = j$. Applying the ring automorphism of H(Z) that permutes $i \to j \to k \to i$ (it preserves the relations and fixes ζ), we get $\zeta j \zeta^{-1} = k$ and $\zeta k \zeta^{-1} = i$. Hence $\zeta\alpha\zeta^{-1} = \sigma(\alpha)$, and $\zeta\alpha\zeta = 4\sigma(\alpha)$. Applying ψ (Lemma 2.3): $h\psi(\alpha)\cdot\mathrm{adj}(h) = 4\psi(\sigma(\alpha))$, and $\mathrm{adj}(h) = \det(h)\cdot h^{-1} = 4h^{-1}$.

(c) σ preserves the norm, the coordinate a_0 and the parity pattern of (a_1, a_2, a_3). If $p \equiv 1 \pmod 4$, $\sigma(S_p) = S_p$. If $p \equiv 3 \pmod 4$, $\sigma(\alpha)$ satisfies Definition 2.1 except possibly the sign rule when $a_0 = 0$, so $\sigma(\alpha)$ or $-\sigma(\alpha)$ lies in S_p, and $\psi(-\beta) = -\psi(\beta)$ is the same point of PGL(2, q). So conjugation by h permutes $\psi(S_p)$. By Lemma 2.4(a), $[\psi(\sigma\alpha)] = [\psi(\alpha)]$ iff $\sigma(\alpha) = \pm\alpha$. The case $\sigma(\alpha) = -\alpha$ forces $a_0 = 0$ and $(a_3, a_1, a_2) = (-a_1, -a_2, -a_3)$, so $a_1 = -a_3 = a_2 = -a_1$, so $\alpha = 0$, impossible. So the fixed points are the $\alpha \in S_p$ with $a_1 = a_2 = a_3 =: a$, i.e. with $a_0^2 + 3a^2 = p$.

- If $p \equiv 2 \pmod 3$, $a_0^2 + 3a^2 \equiv a_0^2 \in \{0, 1\} \pmod 3$, so there is no fixed point.
- If $p \equiv 1 \pmod 3$, the equation $a_0^2 + 3a^2 = p$ has exactly four integer solutions $(\pm a_0, \pm a)$, with $a_0, a \neq 0$ (Fermat; for a prime $p \equiv 1 \bmod 3$ the representation $p = x^2 + 3y^2$ with $x, y > 0$ is unique; cited: D. A. Cox, Primes of the form $x^2 + ny^2$, Wiley 1989, §1 [secondary]). If $p \equiv 1 \pmod 4$ then a_0 is odd and a even (odd + odd $\equiv 0$ and even + $3\cdot$odd $\equiv 3 \bmod 4$), so exactly the two solutions with $a_0 > 0$, namely $a_0 + a(i + j + k)$ with $a = \pm|a|$, are in S_p. If $p \equiv 3 \pmod 4$ then a_0 is even and a odd, $a_0 \neq 0$ ($p \neq 3a^2$), and again exactly the two with $a_0 > 0$ are in S_p. So there are two fixed points.
- If $p = 3$: $a_0 = 0$, $a = \pm 1$, and of $\pm(i + j + k)$ only $i + j + k$ is in S_3. One fixed point.

Since [h] has order 3, every other point lies in a 3-cycle, which gives the cycle types. ■

Theorem 2.7 (cited: Lubotzky–Phillips–Sarnak, Combinatorica 8 (1988) 261–277 [secondary for the constants]; Davidoff–Sarnak–Valette (2003), Ch. 4 [secondary]). Let $p \equiv 1 \pmod 4$ and let $q > p$ be an odd prime. Then $\Gamma_{p,q}$ has $q(q^2 - 1)$ vertices (that is, the LPS Cayley graph $X^{p,q}$ is connected), every eigenvalue of its adjacency matrix other than $\pm(p + 1)$ has absolute value at most $2\sqrt{p}$ (Ramanujan), and $\mathrm{diam}\ \Gamma_{p,q} \leq 2 \log_p n + O(1)$, with an explicit constant given by LPS.

Remark. $\Gamma_{p,q}$ is $X^{p,q}$ itself when p is not a square mod q ($X^{p,q}$ is then a bipartite Cayley graph of PGL(2, q), and a bipartite Cayley graph whose connection set lies in the non-trivial coset of an index-2 subgroup is a Haar graph of that subgroup, Lemma 2.5(d)); it is the bipartite double cover of $X^{p,q}$ when p is a square mod q. Computer-checked: $\Gamma_{p,q}$ has $q(q^2 - 1)$ vertices for $p \in \{3, 5, 7, 11, 13\}$ and every prime q with $p < q$ up to the scan limits of Appendix B (214 graphs; logs `haar_families/scan/lps_p*.txt`, column "vertices"), including $p \equiv 3 \pmod 4$, which Theorem 2.7 as cited does not cover.

2.3 The family written out at Δ = 6

Example 2.8 (p = 5). S_5 = {1 ± 2i, 1 ± 2j, 1 ± 2k}. With x² + y² = −1 in F_q, the six matrices are

$$\psi(1\pm 2i)=\begin{pmatrix}1\pm 2x & \mp 2y\\ \mp 2y & 1\mp 2x\end{pmatrix},\quad \psi(1\pm 2j)=\begin{pmatrix}1 & \pm 2\\ \mp 2 & 1\end{pmatrix},\quad \psi(1\pm 2k)=\begin{pmatrix}1\pm 2y & \pm 2x\\ \pm 2x & 1\mp 2y\end{pmatrix},$$

and conjugation by h = ψ(1 + i + j + k) = [[1 + x + y, 1 + x − y], [−1 + x − y, 1 − x − y]] cycles 1 + 2i → 1 + 2j → 1 + 2k → 1 + 2i and the same with −2: cycle type 3 + 3. In the coordinates of §1.2, **the point [a : b : c : d] of the first copy of PSL(2, q) is joined to the six points**

$$[\,(1\pm 2x)a\mp 2yc : (1\pm 2x)b\mp 2yd : \mp 2ya+(1\mp 2x)c : \mp 2yb+(1\mp 2x)d\,],$$

$$[\,a\pm 2c : b\pm 2d : c\mp 2a : d\mp 2b\,],\qquad [\,(1\pm 2y)a\pm 2xc : (1\pm 2y)b\pm 2xd : \pm 2xa+(1\mp 2y)c : \pm 2xb+(1\mp 2y)d\,]$$

of the second copy; and the point [a : b : c : d] of the second copy is joined to the same six expressions, read as points of the first copy (by Lemma 2.3, adj ψ(α) = ψ(ᾱ), and S̃_5 = S_5, so the six adjugates are the same six matrices up to scalars). For q = 13, with (x, y) = (0, 5) (the first solution of x² + y² + 1 ≡ 0 found by the programs), the component of the first copy's identity is a 6-regular bipartite graph on 2184 vertices of diameter 7 (computer-checked by `lps_haar`, `haar_ecc` and the all-pairs BFS `apsp.c`, §8).

---

3. The 4/3 obstruction

Lemma 3.1 (walk values). Let p, q be odd primes, q > p. A walk of even length L in H(PGL(2, q), ψ(S_p)) from (I, A) ends at a vertex ([ψ(β)], A) with β ∈ H(Z), N(β) = p^L and β ≡ 1 (mod 2H(Z)); in particular the real coordinate b_0 of β is odd.

Proof. The walk with letters ψ(α_1), ..., ψ(α_L) ends at $s_L{}^{-1}$s_{L−1}⋯$s_2{}^{-1}$s_1 with s_t = ψ(α_t), and [ψ(α)⁻¹] = [ψ(ᾱ)] (Lemma 2.3). So it ends at [ψ(β)] with β = ᾱ_L α_{L-1} ⋯ ᾱ_2 α_1, and N(β) = p^L. Reduce mod 2: in H(Z)/2H(Z) we have −1 = 1, so i, j, k commute there. If p ≡ 1 (mod 4), every α_t and ᾱ_t is ≡ 1 (mod 2H(Z)), so β ≡ 1. If p ≡ 3 (mod 4), every α_t and ᾱ_t is ≡ v := i + j + k, and v² = −3 ≡ 1, so β ≡ v^L = 1 since L is even. ■

Theorem 3.2 (distance to ψ(i)). Let p, q be odd primes with q > p, and let L_0 be the least even integer L ≥ 0 with $2p^{L/2} > q^2$. In $\Gamma_{p,q}$:

(a) d((I, A), ([ψ(i)], A)) ≥ L_0, and the same holds for ψ(j) and ψ(k);
(b) if p ≡ 1 (mod 4), equality holds.

Note that det ψ(i) = N(i) = 1, so [ψ(i)] ∈ PSL(2, q) for every q.

Proof. (a) A path from (I, A) to ([ψ(i)], A) has even length L (the graph is bipartite). By Lemma 3.1 there is β with N(β) = p^L, b_0 odd and [ψ(β)] = [ψ(i)], i.e. ψ(β) = cψ(i) with c ∈ F_q^×. By Lemma 2.3, β − c̃i ∈ qH(Z), so q | b_0, b_2, b_3. Put P = $p^{L/2}$, an integer. Since b_0 is odd, b_0 ≠ 0, so b_0^2 ≥ q^2, and then $b_1^2 = P^2 - b_0^2 - b_2^2 - b_3^2 \le P^2 - q^2$, so $|b_1| < P$. Also q^2 divides $b_0^2 + b_2^2 + b_3^2 = P^2 - b_1^2 = (P - b_1)(P + b_1)$. The prime q does not divide P (q ≠ p), so q cannot divide both factors (it would divide their sum 2P). Hence q^2 divides P − εb_1 for some ε ∈ {1, −1}. Since $|b_1| < P$, 0 < P − εb_1 < 2P, so $q^2 \le P - \varepsilon b_1 < 2P$. That is $2p^{L/2} > q^2$, and L ≥ L_0. For ψ(j) and ψ(k): conjugation by h fixes (I, A), is an automorphism of H(PGL(2, q), ψ(S_p)) (Theorem 2.6 and [H] Prop. 3.3(a)), and maps [ψ(i)] to [ψ(j)] to [ψ(k)] (Theorem 2.6(b)), so the three distances are equal.

(b) Let L = L_0, P = $p^{L/2}$, and put $n_1 = 2P - q^2 > 0$. Here P ≡ 1 (mod 4) (p ≡ 1 mod 4), so 2P ≡ 2 (mod 8), and q^2 ≡ 1 (mod 8), so n_1 ≡ 1 (mod 8). By the Gauss–Legendre three-square theorem (cited; n is a sum of three squares unless $n = 4^a(8b + 7)$), $n_1 = u^2 + v^2 + w^2$ with integers u, v, w; since n_1 ≡ 1 (mod 4), exactly one of them is odd, and we name it u. Put

$$\beta = qu + (P - q^2)\,i + qv\,j + qw\,k.$$

Then $N(\beta) = q^2(u^2 + v^2 + w^2) + (P - q^2)^2 = q^2(2P - q^2) + P^2 - 2Pq^2 + q^4 = P^2 = p^L$; $b_0 = qu$ is odd, $b_1 = P - q^2$ is even, and $b_2 = qv$, $b_3 = qw$ are even; so $\beta \equiv 1 \pmod{2H(Z)}$. And $\beta \equiv P \cdot i \pmod{qH(Z)}$, so $[\psi(\beta)] = [P\psi(i)] = [\psi(i)]$. By the LPS factorization theorem (cited: LPS 1988 §3; Davidoff–Sarnak–Valette (2003), Ch. 2, the unique factorization of quaternions $\beta \equiv 1 \bmod 2$ of norm p^L [secondary]), $\beta = \pm p^r\, \alpha_1 \alpha_2 \cdots \alpha_{L-2r}$ with $\alpha_t \in S_p$ and $r \geq 0$. Since $[\psi(\pm p^r\, \gamma)] = [\psi(\gamma)]$, we get $[\psi(i)] = [\psi(\alpha_1 \cdots \alpha_{L'})]$ with $L' = L - 2r$ even. Choose the letters $s_1 = \psi(\alpha_{L'})$, $s_2 = \psi(\tilde{\alpha}_\{L'-1\})$ (so $[s_2{}^{-1}] = [\psi(\alpha_{L'-1})]$), $s_3 = \psi(\alpha_{L'-2})$, $s_4 = \psi(\tilde{\alpha}_\{L'-3\})$, ...; they lie in $\psi(S_p)$ up to sign (Lemma 2.4(b)), and the A-word $s_{L'}{}^{-1} \cdots s_2{}^{-1} s_1$ has value $[\psi(\alpha_1 \cdots \alpha_{L'})]$. So there is a walk of length $L' \leq L_0$ from (I, A) to ($[\psi(i)]$, A). With (a), the distance is L_0. ■

Corollary 3.3. For odd primes $q > p$, with $n = q(q^2 - 1) \geq |V(\Gamma_{p,q})|$:

$$\operatorname{diam} \Gamma_{p,q} \;\geq\; L_0 \;\geq\; 4 \log_p q - 2 \log_p 2 \;>\; \tfrac{4}{3} \log_p n - 2 \log_p 2.$$

Proof. $2p^{L_0/2} > q^2$ gives $L_0 > 2 \log_p(q^2/2) = 4 \log_p q - 2 \log_p 2$, and $n < q^3$. ■

Corollary 3.4 (a vanishing fraction of the Moore bound). Let p be an odd prime, $D \geq 2$, and $q > p$ a prime with $\operatorname{diam} \Gamma_{p,q} \leq D$. Then $q^2 < 2p^{\lfloor D/2 \rfloor}$, the number of vertices satisfies

$$n \leq q(q^2 - 1) < 2^{3/2}\, p^{\frac{3}{2}\lfloor D/2 \rfloor} \leq 2^{3/2} p^{3D/4},$$

and $n < \sqrt{2} \cdot p^{1-D/4} \cdot M^b(p + 1, D)$. In particular, along the LPS family the fraction of the Moore bound tends to 0 both for fixed p and $D \to \infty$ and for fixed $D \geq 5$ and $p \to \infty$.

Proof. $L_0 \leq \operatorname{diam} \leq D$ and L_0 is even, so $L_0 \leq 2\lfloor D/2 \rfloor$ and $2p^{\lfloor D/2 \rfloor} \geq 2p^{L_0/2} > q^2$. Then $n < q^3 < (2p^{\lfloor D/2 \rfloor})^{3/2}$. Finally $M^b(p + 1, D) = 2\Sigma_{j<D}\, p^j > 2p^{D-1}$, so $n/M^b < 2^{3/2} p^{3D/4} / (2p^{D-1}) = \sqrt{2} \cdot p^{1-D/4}$. ■

For comparison: at $D = 8$ the bound is $\sqrt{2}/p$ (28 % for $p = 5$), and the LPS graphs actually reach 1–6 % (Appendix B); the lab's records at $D = 8$ are at 8–22 %, and the $D = 8$ table values with $\Delta \geq 10$ (Gómez's compounds of generalized hexagons) at about 10 % ([H] §5, §6). In the regime D fixed, $\Delta \to \infty$, Delorme's constructions from generalized polygons have $\liminf n \cdot \Delta^{1-D} > 0$ for every D (C. Delorme, J. Graph Theory 9 (1985) 325–334 [abstract]), so arithmetic Haar families lose to them by a factor $p^{D/4-1}$.

Computer-checked 3.5. For $p \in \{3, 5, 7, 11, 13, 17, 19, 29\}$ and every prime q with $p < q \leq 160$, the distance from (I, A) to ($[\psi(i)]$, A), ($[\psi(j)]$, A) and ($[\psi(k)]$, A), computed by breadth-first search (`lps_haar.c`), equals L_0: no exception in 251 cases (log `haar_families/recheck_thmB.log`). This includes $p \equiv 3 \pmod 4$, where equality is not proved here. The eccentricity of (I, A) exceeds L_0 by 0, 1, 2 or 3 (in 76, 151, 22 and 2 of the 251 cases). So in this range the special vertex $\psi(i)$ accounts for the diameter up to at most 3, and at odd D it is usually what limits the family (Appendix B). A second run for $p \in \{5, 13\}$ and all primes $161 \leq q \leq 300$ (log `haar_families/recheck_thmB_large.log`) gave equality in all 50 cases (graphs up to $2.5 \cdot 10^7$ vertices), with ecc $- L_0 = 0$, 1, 2 in 16, 27 and 7 cases; for $p = 13$, $q = 239$ the inequality $2 \cdot 13^4 = 57{,}122 > 239^2 = 57{,}121$ holds by one, and indeed $d = L_0 = 8$ there. The construction in the proof of Theorem 3.2(b) (u, v, w, and β with $N(\beta) = p^{L_0}$, $\beta \equiv 1 \bmod 2$ and $[\psi(\beta)] = [\psi(i)]$) was also checked directly for $p \in \{5, 13, 17, 29, 37\}$ and all primes $p < q < 1000$ (802 cases, `haar_families/check_thmB_construction.py`). An independent re-implementation by squad card c8 (its own quaternion arithmetic and BFS; `research/squads/thmfam-r1/c8/`) confirmed Theorem 2.6(c) for all 137 pairs with $7 \leq q \leq 61$ and Theorem 3.2 in 41 cases with $q \equiv 1 \pmod 4$, and its all-pairs BFS (`apsp.c`) agreed with `haar_ecc` on six LPS graphs. (The squad card also found a bug in the first version of `lps_haar.c`, which mixed the distances of the two sides; the logs above are from the corrected version.)

Proposition 3.6 (the same obstruction in other definite quaternion orders). Let $\mathcal{O}$ be a Z-order in a definite quaternion algebra over Q, with norm form N, and suppose $\mathcal{O} = Z\gamma \oplus \mathcal{O}'$ with $N(a\gamma + \beta') = a^2 + N(\beta')$ for $a \in Z$, $\beta' \in \mathcal{O}'$ (an orthogonal unit γ, $N(\gamma) = 1$). Let $\psi\colon \mathcal{O} \to M_2(F_q)$ be a ring homomorphism with kernel $q\mathcal{O}$ (q an odd prime, $q \nmid p$), let $S \subseteq \mathcal{O}$ be a set of elements of norm p closed under conjugation up to sign, and suppose that no product of an even number $L \geq 2$ of elements of S lies in $Z\gamma$. Then in $H(PGL(2, q), \psi(S))$, every walk of even length L from (I, A) to ($[\psi(\gamma)]$, A) has $2p^{L/2} > q^2$.

Proof. As in Lemma 3.1 the walk ends at $[\psi(\beta)]$ with β a product of L elements of S (conjugates included), so $N(\beta) = p^L$; $[\psi(\beta)] = [\psi(\gamma)]$ gives $\beta = b\gamma + q\beta''$ with $b \in Z$, $\beta'' \in \mathcal{O}'$ (the kernel of ψ is $q\mathcal{O}$ and $\mathcal{O} = Z\gamma \oplus \mathcal{O}'$), and

$\beta'' \neq 0$ by hypothesis. Then $P^2 = b^2 + q^2 N(\beta'')$ with $P = p^{L/2}$ and $N(\beta'') \geq 1$, and the second half of the proof of Theorem 3.2(a) applies verbatim ($q \nmid P$, $q^2 \mid P - \varepsilon b$, $0 < P - \varepsilon b < 2P$). ■

For the Hamilton order, $\gamma = i$, $\mathcal{O}' = \mathbb{Z} + \mathbb{Z}j + \mathbb{Z}k$, and the hypothesis is Lemma 3.1 (every even product is $\equiv 1 \bmod 2$, while elements of $\mathbb{Z}i$ have even real part). Computer-checked (squad card c9 of round 2, `research/squads/thmfam-r2/c9/`): in the order $\mathbb{Z}\langle 1, i, j, ij\rangle$ with $i^2 = -1$, $j^2 = -3$ (norm $a_0^2 + a_1^2 + 3a_2^2 + 3a_3^2$), with S a set of $p + 1$ classes of elements of norm p modulo the units $\pm 1, \pm i$, for $p \in \{5, 7, 11\}$ and $q \leq 97$, the distance from (I, A) to ($[\psi(i)]$, A) equals the least even L with $2p^{L/2} > q^2$ in 65 of 66 cases (in the remaining case, $p = 5$, $q = 61$, it is 12 instead of 10), and $\psi(i)$ is at maximal distance for $p = 5$, $q \in \{61, 83, 97\}$. So the obstruction is a property of definite quaternion arithmetic with a unit orthogonal to the rest, not of Hamilton's order. (The card did not check the hypothesis "no even product in $\mathbb{Z}i$" for its set; the lower bound it measured holds in all 66 cases.)

Remark 3.7 (prior art). The lower bound (4/3) $\log_p n + O(1)$ for the diameter of the bipartite LPS graphs is due to N. T. Sardari, Diameter of Ramanujan graphs and random Cayley graphs, Combinatorica 39 (2019) 427–446, arXiv:1511.09340 [abstract]; his abstract also states that almost all pairs of vertices of a Ramanujan graph are at distance at most $(1 + \varepsilon) \log_p n$, and reports numerics in which LPS diameters grow like (4/3) log and those of random Cayley graphs like log. We did not read his proof; the argument above is short and may be his. The exact value in Theorem 3.2(b) follows from the same arithmetic and the LPS factorization; we did not check whether it is stated there. That the LPS sets are permuted by the Hurwitz unit $(1 + i + j + k)/2$ is classical; we found no source that compares their cycle types with orbit-searched connection sets, and claim nothing more than the observation.

---

4. Cyclic factors

Proposition 4.1. Let K be a finite group, $\kappa: S_p \to K$ any map, and $\tilde{S} = \{(\psi(\alpha), \kappa(\alpha)) : \alpha \in S_p\} \subseteq \mathrm{PGL}(2, q) \times K$. Then for every $k \in K$, the distance in $H(\mathrm{PGL}(2, q) \times K, \tilde{S})$ from (I, e_K, A) to ($[\psi(i)]$, k, A) is at least L_0. Hence if the component of (I, e_K, A) has diameter $\leq D$ and contains ($[\psi(i)]$, k, A) for some k, then $q^2 < 2p^{\lfloor D/2 \rfloor}$.

Proof. The projection $(g, k, X) \mapsto (g, X)$ maps every edge $\{(g, k, A), (\psi(\alpha)g, \kappa(\alpha)k, B)\}$ to the edge $\{(g, A), (\psi(\alpha)g, B)\}$, so it maps walks to walks of the same length and does not decrease distances. Apply Theorem 3.2(a). ■

So a cyclic factor cannot help at the vertex $\psi(i)$ with a fixed q; what it changes is that m is free. The order is $q(q^2 - 1) \cdot m$, the obstruction bounds only q, and the counting bound ([H] Thm. 4.1) allows m up to about p^{D-1}/q^3. Whether the cyclic part can be filled efficiently is a computational question:

Computer-checked 4.2 (squad cards c6, c7). Let r have order 3 mod m. Labels $\alpha \mapsto z(\alpha) \in \mathbb{Z}_m$ that are equivariant for the order-3 symmetry ($z(\sigma\alpha) = r \cdot z(\alpha)$; fixed letters get 0; one 3-cycle gets $(1, r, r^2)$ without loss of generality, the others (t, rt, r^2t)) make $H(\mathrm{PGL}(2, q) \times \mathbb{Z}_m, \tilde{S})$ orbit-closed with $\alpha(x, z) = (hxh^{-1}, rz)$, exactly the records' shape. Searching t for each (q, m), the squads found:

cell	best LPS × $\mathbb{Z}_m$ (q, m)	vertices	of table	lab record	LPS alone (Appendix B)
(6,8)	(13, 13)	28,392	95.3 %	31,680	12,144
(6,9)	(13, 43)	93,912	80.0 %	127,296	24,360
(6,10)	(23, 37)	449,328	99.4 %	524,160	148,824
(8,8)	(11, 151)	199,320	98.9 %	239,904	29,760
(8,9)	(23, 73)	886,512	81.2 %	1,341,600	300,696
(14,8)	(29, 379)	9,232,440	65.1 %	none (the lab does not beat this cell)	1,294,920

Every entry was re-checked by the director with `haar_ecc` (two-source BFS on the component, from the cards' generators: `research/squads/thmfam-r1/c6/gen_c6.py`, `research/squads/thmfam-r1/c7/c7lib.py`), and all but (14,8) also with `verify_cover.py` (independent Python arithmetic); generators and log in

`haar_families/covers/`. These graphs are below the lab's records and below the table, and their labels were searched for each (q, m): they are not a uniform family. They show that the cyclic factor removes most of the 4/3 loss in practice: at (6,10) the cover is 3.0 times the LPS graph alone.

4.3 A label rule: a formula in q and m that beats the 2013 table at (8,8) and (8,9)

Fix p = 7, so Δ = 8 and S_7 = {2 ± (i + j + k)} ∪ C₁ ∪ C₂ with the two 3-cycles of σ

C₁ = {2 − i − j + k, 2 + i − j − k, 2 − i + j − k}, C₂ = {2 − i + j + k, 2 + i − j + k, 2 + i + j − k}

(in the order α, σα, σ²α). For m whose prime factors are all ≡ 1 (mod 3), let r be a root of r² + r + 1 ≡ 0 (mod m) and give the labels

z(2 ± (i + j + k)) = 0, z(C₁) = (1, r, r²), z(C₂) = (t, tr, tr²)

with t a fixed integer (t = 2 below), or a fixed function of m (t = m − 2). This defines S̃(q, m) ⊆ PGL(2, q) × Z_m by one formula in q and m: no search. It is closed under the automorphism (x, z) ↦ (hxh⁻¹, rz) of order 3, the records' shape. The vertex (X, z, A), X ∈ PSL(2, q), is adjacent to the eight vertices (ψ(α)X, z + z(α), B), α ∈ S_7.

Computer-checked 4.3 (squad card c2 of round 2, re-checked by the director). Over the grid q ∈ {11, 13, 17, 19, 23} × m ∈ {7, 13, 19, 31, 37, 43, 61, 49, 91} (45 cells, all components full), the 14 rules tried (t = 2, 3, 4, 5, m − 1, m − 2, r ± 1, …, each with both roots r) have mean excess 2.63–2.88, the best ones 2.63–2.66, against 2.73 for one random t per cell (three seeds, standard deviation 0.03). Two members of the family are above the 2013 table:

cell	(q, m), rule, r	vertices	2013 table	lab record	checks
(8,8)	(17, 43), t = 2, r = 36	**210,528**	201,480 (+4.5 %)	239,904	`haar_ecc`, `verify_cover.py`, `hc_factor.c`, `build_haar2.py`, all-pairs `apsp.c`
(8,9)	(23, 91), t = m − 2 = 89, r = 9	**1,105,104**	1,091,232 (+1.3 %)	1,341,600	`haar_ecc`, `verify_cover.py`, `hc_factor.c`, `build_haar2.py`

(generators: `haar_families/covers/d8D8_lps7_q17_m43.gens`, `d8D9_lps7_q23_m91.gens`; the same graphs as Haar graphs of PSL(2, q) × Z_m in the lab's solution-file format, by Lemma 2.5(d), in `covers/sols/`, which the lab's record checkers `hc_factor.c` and `research/frontier/build_haar2.py` accept; log `covers/recheck.log`.) Both are below the lab's records at these cells ([H] §5), so they are not new records; and (q, m) is the best cell of the grid, chosen after the fact. What they show is that a member of a uniform arithmetic family of the records' exact shape (k = 3, cycle type 1 + 1 + 3 + 3, an inner automorphism of order 3 times a scaling of Z_m) is above the 2013 table at D = 8 and 9, while the same family without the cyclic factor is 6.8 and 3.6 times below it (Appendix B). We do not have a proof of the diameter of any member: the proofs of §3 give lower bounds only.

Label rules elsewhere (round 2). For p = 5 the best fixed rule (t = 3) has mean excess 2.86 over 45 cells, against 2.93–3.09 for one random t per cell (card c1); a nonlinear rule z(α) = Σ_k (a_k mod 3)·r^{k-1} gives 2.82 over 24 cells (card c11); over 35 cells, one random equivariant labelling gives 2.97 and the best t per cell 2.71 (card c10, exhaustive over t and both roots r). For p = 7, one random equivariant labelling gives 2.68 (card c10). For p = 11, the best of 10 rules ties with random labels (2.53 against 2.56 over 4 × 8 cells; card c3). So a fixed rule recovers about half of the gain that a search over t gives at p = 5, and little or nothing at p = 7 and 11. For Δ = 4 (p = 3), covers with labels searched exhaustively reach 66 % of the table at (4,8) and 88 % at (4,9) (card c6), so the cells (4,8), (4,9), which the lab does not beat, stay unbeaten. For Δ = 12 the best

cover found reaches 51 % of the table at (12,9) and none of diameter 8 improves on plain LPS (card c7); for Δ = 14 the best is the 65 % graph at (14,8) of §4.2.

---

5. Other uniform families (squad round 1)

Score. For a family S(q) and the primes 23 ≤ q ≤ 113, the excess is diam − $log_{\Delta-1}|G|$ with |G| = one side of the graph, averaged over q (`haar_families/family_score.py`, which runs `haar_ecc`). Lower is better. A uniform family gets one S per q, so it should be compared with one random orbit-closed draw per q, not with the records, which are the best of millions of evaluations for one (q, m).

Computer-checked 5.1 (mean excess, 22 primes 23 ≤ q ≤ 113, group PSL(2, q) unless stated; from `research/squads/thmfam-r1/scoreboard.jsonl` and the cards' folders). Baselines first (card c13: 8 seeds each, mean ± standard deviation over seeds):

one random draw per q	Δ = 6	Δ = 7	Δ = 8
random S, no symmetry, PSL	3.20 ± 0.05	2.95 ± 0.05	2.95 ± 0.05
random S, no symmetry, SL	3.05 ± 0.02	2.89 ± 0.03	2.76 ± 0.03
random orbit-closed S (k = 3), PSL	3.31 ± 0.11	3.04 ± 0.10	3.28 ± 0.08
random orbit-closed S (k = 3), SL	3.34 ± 0.11	3.09 ± 0.09	3.14 ± 0.09

(Orbit-closed sets with Δ = 7, 8 contain I, and I and w, as in the records; the unconstrained sets are Δ distinct random elements.) The spread of the excess over q is 0.1–0.4 per prime. So a single orbit-closed draw is slightly worse than a single unconstrained draw (by 0.1–0.4); the orbit search wins because it makes the search space smaller and the misses come in clumps ([H] §3.5), not because orbit-closed sets are better one by one. The families:

family (one S per q, no search per q)	Δ	mean excess	card
LPS, p = 5 (component in PGL(2, q))	6	4.00	director
integer lifts, O(X) ∪ O(Y), X, Y ∈ SL(2, Z), entries ≤ 3 (best of 206)	6	3.36	c1
Dickson / trace / power formulas (best of 15)	6	3.13	c11
integer lifts, {I} ∪ O(X) ∪ O(Y) (best of 1212; mean of the top 10: 3.35)	7	3.10	c2
primitive-root formulas, e.g. X = [[g, 1], [0, g^{-1}]], Y = [[1/2, g], [g, 2(1 + g^2)]] (best of 16)	7	2.96	c10
integer lifts × Z_{13}, equivariant labels (best of about 500 screened)	7	3.12	c4

family (one S per q, no search per q)	Δ	mean excess	card
integer lifts, two fixed letters + two orbits (all 234 with entries ≤ 2)	8	3.84	c3
Dickson / trace / power formulas (best of 15)	8	3.16	c11
conic model: Cayley transforms of integer matrices (best of 8)	8	3.34	c16
integer lifts, one fixed letter + three orbits (all 858 with entries ≤ 2)	10	3.40 (random orbit-closed: 2.76)	c3
single orbits of order 6 (best uniform x, y)	12	2.73 (random, k = 3: 2.55–2.64)	c15
LPS + one extra α-orbit (repair)	9	3.24 on 7 primes (random: 2.72–2.86)	c18
the records' own sets, lifted to SL(2, Z) and used at the other 21 primes	6, 7	2.87–3.24 (at their own q: 1.99–2.13)	c5
SL(2, q) × Z_m, m ≈ q, integer letters with equivariant labels (10 primes 23–61)	6	3.38 (same letters, m = 1: 3.60)	c17
the lab's records (one (q, m) each, annealed)	6–16	about 2.0–2.3	[H]

Torus-orbit families with Δ growing like q (card c12: S = {I} ∪ T-conjugates of x for a torus T of PSL(2, q)) have diameter 5 to 7 for 31 ≤ q ≤ 61 and reach at most 1.8 % of the Moore bound at their diameter; no family of diameter ≤ 4 appeared. The excess of LPS grows with log_5|G| with slope 0.41 (card c14, 38 primes 23 ≤ q ≤ 199; Corollary 3.3 predicts 1/3), that of random orbit-closed draws with slope −0.04.

Heuristic 5.2 (reading of 5.1).

1. No formula family beats one random draw at the same Δ by more than about 0.2, and the best of each kind was selected among 8–1212 variants on the same primes, so the small wins are within the selection effect. Several kinds are clearly worse: LPS, and the integer lifts at Δ = 8 and 10 (by 0.5–0.9).
2. Arithmetic structure costs: LPS is about 0.8 worse than one random draw at q ≤ 113, and the gap grows like (1/3) log_p n (Theorem 3.2; card c14 measured slope 0.41 for LPS and −0.04 for random draws).
3. The records do not transport: the set that is a record at its own q (excess about 2.0) is an ordinary set at every other q (card c5). A record is a property of the pair (q, S), not of a formula.
4. Larger symmetry (k = 5, 6) is worse than k = 3 at the same Δ, for formulas and for random draws (card c15).
5. The records are about 1 below a single draw. In the model of [H] §3.5 a single random S misses about μ = N $e^{-W/N}$ elements and is valid with probability about $e^{-\mu}$; a search over T independent sets can afford μ* ≈ ln T, which lowers the needed W/N by about ln μ*, roughly one unit of $log_{\Delta-1}$ at the sizes of the records. A formula gives one set per q, so it cannot have this advantage unless it is better than random for structural reasons, as the Singer sets are at D = 3. We found no such structure at D ≥ 7.

6. Why no record-quality uniform family is in sight

6.1 Fixed Δ, D → ∞. By Proposition 1.1 a family here has q → ∞. A uniform family of record quality would have diam ≤ $log_{\Delta-1}$ n + O(log log n) along an explicit sequence of groups. No explicit family of bounded-degree graphs with diam ≤ (1 + ε) $log_{\Delta-1}$ n is known [secondary]; random Δ-regular graphs have diameter (1 + o(1)) $log_{\Delta-1}$ n (B. Bollobás, W. Fernandez de la Vega, Combinatorica 2 (1982) 125–134 [secondary]), and so, conjecturally, do random Haar graphs of SL(2, q) (research/theory/THRESHOLD.md, Conjecture 6, on master). The best explicit candidates, the Ramanujan graphs, have almost all distances (1 + o(1)) $log_{\Delta-1}$ n (Sardari [abstract]; E. Lubetzky, Y. Peres, Cutoff on all Ramanujan graphs, GAFA 26 (2016) [secondary]), but for the LPS graphs a few vertices are at distance (4/3) log_p n (Theorem 3.2). That is the precise way in which the most natural uniform family of the records' shape fails, and Corollary 3.4 turns it into a loss of a factor $p^{D/4-1}$ against the Moore bound.

6.2 Fixed D, Δ → ∞. The LPS family with q ≈ $p^{D/4}$ has degree p + 1 and order about $p^{3D/4}$ at diameter ≈ D (Corollary 3.4 is an upper bound; the lower side is Theorem 2.7, order $p^{(D-O(1))/2}$). Delorme's polygon constructions reach a fixed fraction of p^{D-1} (§3). We know of no Haar-graph family of the records' groups that reaches lim inf n·Δ^{1-D} > 0 at D = 8, 9 or 10; the random-S theorem of research/theory/THRESHOLD.md reaches n ≥ Δ^{D-1}/((D − 1)!·(1 + η)·ln n) for every finite group, which loses a logarithm.

6.3 What would change this. A formula that beats one random draw by about 1 in excess, uniformly in q; or a proof that some explicit cover (for example LPS × Z_m with labels given by a rule) has diameter log_p n + O(log log n). Neither is in sight. The squads' round 2 looked for label rules for LPS × Z_m (Appendix A).

---

7. Prior art

- **LPS graphs.** A. Lubotzky, R. Phillips, P. Sarnak, Ramanujan graphs, Combinatorica 8 (1988) 261–277; G. A. Margulis, Problems Inform. Transmission 24 (1988) 39–46 [secondary]; the book of Davidoff–Sarnak–Valette (2003) [secondary]. Definitions 2.1–2.2, Lemma 2.3, Theorem 2.7 and the factorization used in Theorem 3.2(b) are theirs.
- **The 4/3 lower bound.** Sardari (2019) [abstract], Remark 3.7.
- **Typical distances in Ramanujan graphs.** Lubetzky–Peres (2016) [secondary]; Sardari (2019) [abstract].
- **Haar graphs of SL(2, q) and PSL(2, q) in extremal problems.** [H] §6.1: Exoo–Jajcay (girth, a Haar graph of SL(2, 83)); Bray–Parker–Rowley (the group PSL(2, q) × Z_3); the LPS graphs themselves are Haar graphs of PSL(2, q) in the bipartite case (Lemma 2.5(d)). In the bipartite degree–diameter table at D = 8–10 (Miller–Širáň, DS14v2, 2013, as reproduced by the Combinatorics Wiki) we found no LPS graph; the novelty check `research/lit/novelty-2026-10-25.md` (branch `claude/novelty-literature-check-4i5frp`, claim 5) found no Haar graphs of SL(2, q) or PSL(2, q) there.
- **Asymptotic lower bounds for N^b(Δ, D).** Delorme (1985) [abstract via the novelty check]; random bipartite graphs (V. Klee, D. Larman, E. M. Wright, The diameter of almost all bipartite graphs, Studia Sci. Math. Hungar. 15 (1980) 39–43 [secondary]); research/theory/THRESHOLD.md for random Haar graphs.
- **Orbit-closed connection sets.** [H] §6.1 (an old device; Singer sets and their multipliers).

What this note adds: Theorem 2.6 (the LPS sets have the records' orbit shape, with the cycle type computed), the elementary proof and the exact value in Theorem 3.2, Corollary 3.4 in the form of a fraction of the Moore bound, the covers of §4 and the measurements of §5. We claim no novelty for Lemmas 2.3–2.5 or for Theorem 2.7.

---

8. How to re-check

```
cd research/theorems/haar_families
cc -O3 -o lps_haar lps_haar.c -lm      # LPS graph: generators (-v), eccentricities, d(I, psi(i)) etc.
cc -O3 -o haar_ecc haar_ecc.c          # any Haar graph of SL/PSL/PGL(2,q) x Z_m from generator lines "a b c d z"
./lps_haar 5 13 -v                     # Example 2.8: 2184 vertices, diameter 7, d(I, psi(i)) = 6 = L0
```

```
python3 check_thmB.py 3,5,7,11,13,17,19,29 160 > recheck_thmB.log     # Computer-checked 3.5
python3 check_thmB.py 5,13 300 161 > recheck_thmB_large.log           # the same for 161 <= q <= 300 (slow)
python3 check_thmB_construction.py                                    # the beta of Theorem 3.2(b), q < 1000
./lps_haar 5 13 -v | grep '^gen' | sed 's/.*mat(//; s/) z=/ /; s/,/ /g' > g.txt
python3 component_edges.py pgl 13 1 e.txt < g.txt                    # independent edge list of the component
cc -O2 -o apsp ../../lifts/coordinator_check/apsp.c && ./apsp e.txt 6 40   # all-pairs BFS: diameter 7
python3 appendix_b.py                  # Appendix B from scan/lps_p*.txt (./scan_lps.sh p qmax rebuilds a scan)
python3 family_score.py --gen "./gen_lps.sh 5 {q}" --group pgl --qrange 23 113   # §5, LPS row
python3 ../../squads/thmfam-r1/c6/gen_c6.py 23 37 19 0 | ./haar_ecc pgl 23 37  # §4.2, (6,10)
python3 ../../squads/thmfam-r1/c6/gen_c6.py 23 37 19 0 | python3 verify_cover.py 23 37
python3 gen_lps_cover.py 7 11 151 cyc 1 57 | ./haar_ecc pgl 11 151           # §4.2, (8,8): diameter 8
python3 ../../squads/thmfam-r1/c7/c7lib.py 13 29 379 1 202 352 308 | ./haar_ecc pgl 29 379   # §4.2, (14,8)
python3 gen_lps_cover.py 7 17 43 cyc 1 2 --rk 2 | ./haar_ecc pgl 17 43        # §4.3, (8,8): 210,528 vertices, diam
python3 gen_lps_cover.py 7 23 91 cyc 1 89 --rk 1 | ./haar_ecc pgl 23 91       # §4.3, (8,9): 1,105,104 vertices, di
python3 ../../frontier/build_haar2.py covers/sols/d8D8_psl_17_43_lps7_t2.txt   # the lab's record checker (also hc_
```

Appendix A. The squad rounds

Two rounds of Claude Haiku cards (the lab's `haiku-squad` skill; model haiku, effort high), directed by this session. The cards, the shared rules and every card's code and notes are in `research/squads/thmfam-r1/` and `research/squads/thmfam-r2/`, one line per card in `scoreboard.jsonl`; the rounds are logged in `research/squads/LOG.md`. The squads searched constructions and tested the theorems on small q; the proofs are the director's. Every number quoted in the note from a card was re-run by the director with the commands of §8, except where it says "card" only (the mean excesses of §5, which are averages over many runs and are reproducible from the cards' folders).

Round 1 (18 cards): uniform families. c1–c4 integer lifts (Δ = 6, 7, 8, 10, and with Z_m); c5 the records' sets at other q; c6–c7 LPS × Z_m with searched labels; c8 verification of Theorems 2.6 and 3.2 (found a bug in the first `lps_haar.c`); c9 the far vertices of LPS graphs (the guess "all far vertices come from pure quaternions" failed); c10–c11 q-dependent formulas; c12 torus orbits with Δ ~ q; c13–c14 random baselines and the trend in q; c15 symmetries of order 5 and 6; c16 the conic model; c17 cyclic factors with m ~ q; c18 repairs of LPS. Outcome: every formula family is at the level of one random draw (§5); LPS × Z_m covers were the only direction that came close to the table (§4).

Round 2 (11 cards): LPS × Z_m covers. c1–c3 label rules (a formula in m) for p = 5, 7 and 11, 13; c4–c5 pushes of p = 5 and p = 7 covers past the table with labels searched per (q, m); c6 Δ = 4 covers at (4,8), (4,9); c7 Δ = 12, 14 covers; c8 a numerical audit of §§1.2, 2, 3 and 4.1 of this note with its own code (one discrepancy, a miscount in the first draft of Appendix B, now corrected; everything else passed, including the neighbourhood formulas of Example 2.8 against the edges, Lemma 2.5(c), Proposition 4.1 in covers and Corollary 3.4 on all 214 scan lines); c9 a second quaternion algebra (Proposition 3.6); c10 random and exhaustive label baselines; c11 nonlinear arithmetic labels. Outcome: the rules of c2 give the two members above the table in §4.3; nothing else beat the table. Searching labels per (q, m) by hill climbing (c4, c5) did not beat the table either (one claimed (8,8) graph of c5 was not connected within distance 8 and has diameter 9; the director's re-check caught it, and the card withdrew it). Fixed rules recover part of the gain of a per-cell search at p = 5 and tie with random labels at p = 7, 11 (c1, c3, c10).

Appendix B. The LPS family cell by cell

Computer-checked (`haar_families/appendix_b.py` from the scan logs `haar_families/scan/lps_p*.txt`, which list the order and the diameter of $\Gamma_{p,q}$ for every prime p < q up to the scan limit, computed by `lps_haar`). For each cell the largest LPS graph of diameter ≤ D; A star (*) means the largest scanned q still had diameter ≤ D, so larger LPS graphs may exist there. "Thm 3.2: q ≤" is the largest prime q with $q^2 < 2p^{\lfloor D/2 \rfloor}$, the limit of Corollary 3.4.

Δ (p)	D	q	vertices	% Moore	Thm 3.2: q ≤	table	table / LPS	lab record
4 (3)	7	7	336	15.37	7	840	2.5	864
4 (3)	8	7	336	5.12	11	2,184	6.5	–
4 (3)	9	11	1,320	6.71	11	4,970	3.8	–
4 (3)	10	13	2,184	3.70	19	11,748	5.4	12,416
4 (3)	11	19	6,840	3.86	19	–	–	–
4 (3)	12	19	6,840	1.29	37	–	–	–
6 (5)	7	13	2,184	5.59	13	8,310	3.8	8,880
6 (5)	8	23	12,144	6.22	31	29,790	2.5	31,680
6 (5)	9	29	24,360	2.49	31	117,360	4.8	127,296
6 (5)	10	53	148,824	3.05	79	452,032	3.0	524,160
6 (5)	11	67	300,696	1.23	79	–	–	–
6 (5)	12	107	1,224,936	1.00	173	–	–	–
8 (7)	7	23	12,144	4.42	23	40,586	3.3	43,344
8 (7)	8	31	29,760	1.55	67	201,480	6.8	239,904
8 (7)	9	67	300,696	2.24	67	1,091,232	3.6	1,341,600
8 (7)	10	107	1,224,936	1.30	181	6,927,210	5.7	7,818,096
8 (7)	11	173	5,177,544	0.79	181	–	–	–
8 (7)	12	229*	12,008,760	0.26	479	–	–	–
12 (11)	7	47	103,776	2.66	47	664,300	6.4	693,500
12 (11)	8	71	357,840	0.83	167	4,650,100	13.0	–
12 (11)	9	167	4,657,296	0.99	167	30,001,152	6.4	34,026,720
12 (11)	10	251	15,813,000	0.30	563	300,383,050	19.0	325,832,976
12 (11)	11	257*	16,974,336	0.03	563	–	–	–
12 (11)	12	257*	16,974,336	0.00	1879	–	–	–
14 (13)	7	59	205,320	1.96	61	1,771,560	8.6	1,789,588
14 (13)	8	109	1,294,920	0.95	239	14,172,480	10.9	–
14 (13)	9	233	12,649,104	0.72	239	95,087,738	7.5	116,068,656
14 (13)	10	257*	16,974,336	0.07	859	1,213,477,190	71.5	1,326,627,120
14 (13)	11	257*	16,974,336	0.01	859	–	–	–
14 (13)	12	257*	16,974,336	0.00	3089	–	–	–

Scan limits: p = 3: q ≤ 157, p = 5: q ≤ 157, p = 7: q ≤ 229, p = 11: q ≤ 257, p = 13: q ≤ 257.

At odd D, in the 13 cells not cut by the scan limit, the largest q reached equals the limit of Theorem 3.2 in 8 cells ((4,7), (4,9), (4,11), (6,7), (8,7), (8,9), (12,7), (12,9)), is the next prime below it in 3 ((6,9): 29 against 31; (14,7): 59 against 61; (14,9): 233 against 239), and is farther below in 2 ((6,11): 67 against 79; (8,11): 173 against 181). So at odd D the vertex ψ(i) is usually what stops the family, as far as the scan goes. At even D = 2κ the limit is the same as at D = 2κ + 1, and the generic vertices, at distance about log_p n, stop the family first.